

Nonabelian homotopical cohomology, higher fiber bundles with connection, and their σ -model QFTs

January 14, 2009

Abstract

Nonabelian cohomology can be regarded as a generalization of group cohomology to the case where both the group itself as well as the coefficient object are allowed to be generalized to ∞ -groupoids or even to general ∞ -categories. Cocycles in nonabelian cohomology in particular represent higher principal bundles (gerbes) – possibly equivariant, possibly with connection – as well as the corresponding *associated* higher vector bundles.

We formulate nonabelian cohomology and its classification of fiber bundles in a general context of enriched homotopy theory, and discuss general issues such as lifting and extension problems. We list examples and applications with enrichment over higher categories which describe higher principal bundles and higher vector bundles, possibly equivariant, possibly with connection.

Building on this we propose, expanding on considerations in [15, 42, 6], a systematic ∞ -functorial formalization of the σ -model quantum field theory associated with a given nonabelian cocycle regarded as the background field for a brane coupled to it. We define propagation in these σ -model QFTs and recover central aspects of *groupoidification* [1, 2].

In a series of examples we show how this formalization reproduces familiar structures in σ -models with finite target spaces such as Dijkgraaf-Witten theory and the Yetter model. Applications to σ -models with smooth target spaces is developed elsewhere [26].

Contents

1	Introduction	4
2	Enriched homotopy theory	5
3	Nonabelian homotopical cohomology	5
3.1	Pointed objects	8
3.2	Universal fiber bundles	9
3.3	Cocycles and bundles	11
3.4	Sections and homotopies	12
3.5	Extension and lifting problem	15
3.5.1	Extension	15
3.5.2	Lifting	17
3.5.3	Local semi-trivializations	17
4	Examples and Applications	17
4.1	Homotopical contexts	17
4.2	Closed monoidal homotopical categories	18
4.3	Pointed objects	18
4.4	Monoid of loops	18
4.5	Universal bundles	18
4.6	Bundles	18
4.7	Lifting problems	19
4.8	Extension problem	19
4.9	Equivariance on semi-total spaces	19
5	Quantization of nonabelian cocycles to σ-models	19
5.1	σ -Models	19
5.2	Branes and bibranes	20
5.3	Quantum propagation	22
6	Examples and applications	23
6.1	General examples	23
6.1.1	Ordinary vector bundles	23
6.1.2	Group algebras and category algebras from bibrane monoids	24
6.1.3	Monoidal categories of graded vector spaces from bibrane monoids	24
6.1.4	Twisted vector bundles	25
6.1.5	2-Hilbert spaces	26
6.1.6	The path integral	26
6.2	Dijkgraaf-Witten model: target space \mathbf{BG}_1	26
6.2.1	The 3-cocycle	27
6.2.2	Chern-Simons theory	28
6.2.3	Transgression of DW theory to loop space	28
6.2.4	The Drinfeld double modular tensor category from DW bibranes	30
6.2.5	The DW path integral	31
6.3	Yetter-Martins-Porter model: target space \mathbf{BG}_2	31

A	ω-Categories and their Homotopy Theory	31
A.1	Shapes for ∞ -cells	32
A.2	ω -Categories	33
A.3	ω -Groupoids	34
A.4	Cosimplicial ω -categories	34
A.5	Monoidal biclosed structure on ω Categories	35
A.6	Model structure on ω Categories	36

1 Introduction

A σ -model should, quite generally, be an n -dimensional quantum field theory which is canonically associated with the geometric structure given by a connection on a bundle whose fibers are n -categories – for instance a (higher) gerbe with connection.

For example for $n = 1$ a line bundle with connection over a Riemann manifold gives rise to the ordinary quantum mechanics of a charged particle. For $n = 2$ a line bundle gerbe with connection over a Lie group G gives rise to the 2-dimensional quantum field theory known as the WZW-model. For $n = 3$ a Chern-Simons 2-gerbe with connection over BG gives rise to Chern-Simons QFT.

The natural conceptual home of these higher connections appearing here is *differential nonabelian cohomology* [26], a joint generalization of sheaf cohomology, group cohomology and nonabelian group cohomology. One arrives at this general notion of cohomology for instance by first generalizing the coefficients of sheaf cohomology from complexes of abelian groups, via crossed complexes of groupoids and their equivalent ∞ -groupoids [8], to general ∞ -categories, and secondly by generalizing the domain spaces via orbifolds, hypercovers and their equivalent ∞ -groupoids also to general ∞ -categories. Therefore nonabelian cocycles are cocycles *on* ∞ -categories with coefficients *in* ∞ -categories. Moreover, when suitably interpreted such a cocycle is nothing but an ∞ -functor from its domain to its coefficient object, hence a rather fundamental concept. In one way or other it is well known that such cocycles in particular classify fiber bundles – possibly equivariant – whose fibers are higher categories.

In a series of articles [3, 35, 36, 37] (see also [23]) it was shown for low n that by *internalizing* this notion of ∞ -functorial nonabelian cocycles from the category of plain sets into a category of generalized *smooth spaces*, it yields a good notion of generalized *differential cohomology*: if the domain ∞ -category is taken to be the smooth fundamental ∞ -groupoid of a smooth space, then smooth ∞ -functors out of it provide a higher dimensional notion of *parallel transport* and characterize higher connections on higher fiber bundles.

Indeed, regarding an ∞ -functorial cocycle as a parallel transport functor generally provides a useful heuristic for the sense in which generalized cocycles are nothing but ∞ -functors from their domain to their coefficient object, even if there is no smooth structure and no connection around: the ∞ -functorial cocycles characterizing for instance a fiber bundle is the *fiber-assigning functor* which to each point in base space assigns the fiber sitting over that point, to each morphism in base space (be it a jump along an orbifold action, or a jump between points in the fiber of a Čech cover, or indeed a smooth path in base space) the corresponding morphisms between the fibers over its endpoints, and similarly for higher morphisms.

From this perspective much can already be learned from and achieved in finite approximations to full smooth differential cocycles. A central example is Dijkgraaf-Witten theory as a finite version of Chern-Simons theory: while Chern-Simons theory is a σ -model governed by a differential 3-cocycle on BG – in the smooth context – usually addressed as the *Chern-Simons 2-gerbe* –, Dijkgraaf-Witten theory is diagrammatically the same setup, but now internal to ∞ -categories internal to **Sets**: the space BG is replaced by the finite groupoid **BG** with one object and Hom-set the finite group G . So among other things, ∞ -functorial nonabelian cohomology, which treats group cocycles and higher bundles/higher gerbes intrinsically on the same footing, gives a precise formalization of the way in which finite group models such as Dijkgraaf-Witten theory are related to their smooth cousins such as Chern-Simons theory.

For that reason it is worthwhile to study the ∞ -functorial nonabelian cohomology perspective on finite group σ -models before adding the further technical complication of working internal to smooth spaces. While discussion of differential nonabelian cohomology in the context of smooth spaces is in preparation in [26], here we develop some concepts and their applications in the simpler context of plain sets.

In sections 3 and 5 we set up the central concepts which we use to formalize the notion of a σ -model associated with a (differential) nonabelian cocycle. In particular we formalize in this context the notion of higher sections and higher spaces of states as indicated in [15, 42], and generalize to corresponding notions of branes and bibranes [17]. In section 5 we then go through a list of examples and applications illustrating these concepts.

2 Enriched homotopy theory

There are two major well-developed 1-categorical tools for handling models for (directed) spaces and higher (directed) homotopies, i.e. for ∞ - or ω -categories: these are *enriched category theory* and *model category theory*. A comprehensive treatment is obtained from the combination of the two, known as *enriched homotopy theory* or *homotopy coherent category theory*.

For enriched category theory we rely on the canonical textbook [20]. For homotopy theory we mainly make use of the seminal article [7] and hence mostly require less structure than in full model category theory. The systematic study of enriched homotopy theory is much younger: we adopt the point of view of [33], which follows the textbooks [12, 18].

In section 3 we consider higher fiber bundles – possibly equivariant, possibly with connection – in a generic homotopical context without committing ourselves to a concrete model for ∞ -categories or ω -categories, aiming to come close to requiring a necessary minimum of structural prerequisites. The idea is to stipulate that an object \mathbf{A} in a category of higher structures should be

1. a *generalized space* locally modeled on objects in a locally small category S ;
2. and equipped with a consistent notion of *homotopy* between maps into it.

The first point we read as implying that \mathbf{A} is characterized by maps of test-objects in S into it, making it a presheaf on S . The second point then suggests that this presheaf takes values in a homotopical category \mathcal{V} and that S is \mathcal{V} -enriched such that there are \mathcal{V} -internal spaces of morphisms. The solution to this requirement suggested by [33] is to take \mathcal{V} to be a *closed monoidal homotopical category* so that $\mathcal{C} := [S^{\text{op}}, \mathcal{V}]$ is \mathcal{V} -enriched and becomes *\mathcal{V} -enriched homotopical* after choosing suitable local extensions of the weak equivalences in \mathcal{V} to \mathcal{C} . The nice consequence of these natural assumptions is that the homotopy category $\text{Ho}_{\mathcal{C}}$ of \mathcal{C} is naturally $\text{Ho}_{\mathcal{V}}$ -enriched while itself homotopical in a \mathcal{V} -enriched sense and thus retains information about higher homotopies and their weak inverses. This should make it an accurate enriched 1-categorical model for an ∞ -category of ∞ -categories modeled on S .

Two complementary useful perspectives on objects in such homotopically enriched presheaf categories \mathcal{C} are familiar: as generalized homotopical spaces and as ∞ -stacks. For instance if $S = \text{Diff}$ is the site of smooth manifolds, then a higher structure *probeable* by mapping objects in S into it may be a Lie groupoid \mathbf{A} , a higher structure whose smoothness is modeled by how test-objects in Diff are mapped into it. By instead regarding the assignment to an object $X \in \text{Diff}$ of collections $[X, \mathbf{A}] \xrightarrow{\text{Yoneda}} \mathbf{A}(X)$ of maps from X into \mathbf{A} as primary, it appears instead equivalently as the *differentiable stack* on S presented by the Lie groupoid.

More generally, for the choice $\mathcal{V} := \text{SimplicialSets}$ it is well known [41, 40] that the $\text{Ho}_{\mathcal{V}}$ -enriched homotopy category $\text{Ho}_{\mathcal{C}}$ for $\mathcal{C} := [S^{\text{op}}, \mathcal{V}]$ the \mathcal{V} -enriched category of simplicial presheaves (with weak equivalences the local weak equivalences of SimplicialSets) is an enriched 1-categorical model of ∞ -stacks on S . In our examples and applications in section 4 we find it useful to choose for \mathcal{V} categories of globular (instead of simplicial) n -categories equipped with the *folk model structure* [14, 9]; such as to make contact to results in [37]. The general considerations in section 3 are independent of all such choices of concrete realizations of \mathcal{V} and \mathcal{C} .

3 Nonabelian homotopical cohomology

We place ourselves in the context of derived \mathcal{V} -enriched category theory for \mathcal{V} a *closed monoidal homotopical category* and consider a *\mathcal{V} -enriched homotopical category* \mathcal{C} as described in [33].

Recall (sections 15 and 16 in [33]) that this means that \mathcal{V} is a category equipped with a choice of weak equivalences compatible with its closed monoidal structure, and in particular that on the Sets -enriched category $\mathcal{C}_0 := \mathcal{V}\text{-Cat}(I, \mathcal{C})$ underlying the \mathcal{V} -enriched category \mathcal{C} there is a \mathcal{V} -valued hom-functor $\text{hom} : \mathcal{C}_0 \times \mathcal{C}_0 \rightarrow \mathcal{V}$ compatible with the action of \mathcal{V} on \mathcal{C} by powers $[-, -]$ and by copowers \otimes which determines

the \mathcal{V} -enrichment of \mathcal{C} by $\mathcal{C}(\mathbf{X}, \mathbf{A}) \simeq \text{hom}(\mathbf{X}, \mathbf{A})$ for all objects \mathbf{Y}, \mathbf{A} in \mathcal{C}_0 . Its right-derived functor $\mathbb{R}\text{hom}$

$$\begin{array}{ccc} \mathcal{C}_0^{\text{op}} \times \mathcal{C}_0 & \xrightarrow{\text{hom}} & \mathcal{V} \\ \downarrow & & \downarrow \\ \text{Ho}_{\mathcal{C}_0}^{\text{op}} \times \text{Ho}_{\mathcal{C}_0} & \xrightarrow{\mathbb{R}\text{hom}} & \text{Ho}_{\mathcal{V}} \end{array}$$

similarly induces the $\text{Ho}_{\mathcal{V}}$ -enriched category $\text{Ho}_{\mathcal{C}}$ whose underlying **Sets**-enriched category is the homotopy category $\text{Ho}_{\mathcal{C}_0}$ of \mathcal{C}_0 (proposition 16.2 in [33]).

In such a setup cohomology identifies with the hom-objects in the homotopy categories, and algorithms for computation of cohomology are algorithms for computation of the right derived (internal) hom-functors:

Definition 3.1 (cohomology) For \mathcal{C} a \mathcal{V} -enriched homotopical category of the form $[S^{\text{op}}, \mathcal{V}]$ or $\text{Sh}(S, \mathcal{V})$, we say for \mathbf{A} any object of \mathcal{C} and X a representable object that $H(X, \mathbf{A}) := \text{Ho}_{\mathcal{C}_0}(X, \mathbf{A})$ is the cohomology of X with coefficients in \mathbf{A} . The generalized elements $c : I \rightarrow Z(X, \mathbf{A})$ of the \mathcal{V} -object $Z(X, \mathbf{A}) := \text{Ho}_{\mathcal{C}}(X, \mathbf{A})$ are the cocycles on X with coefficients in \mathbf{A} . Homotopies between these are the coboundaries.

More generally $\text{Ho}_{\mathcal{C}}(-, \mathbf{A})$ can be interpreted as computing equivariant cohomology and generalizations thereof:

Definition 3.2 (equivariant and relative cohomology) For $i : X \hookrightarrow \mathbf{X}$ a monomorphism in \mathcal{C}_0 it is often useful to address $H^{\mathbf{X}}(X, \mathbf{A}) := \text{Ho}_{\mathcal{C}_0}(\mathbf{X}, \mathbf{A})$ as equivariant cohomology on X , where the kind of equivariance is controlled by i . Conversely the kernel of i^* for any $i : X \rightarrow \mathbf{X}$ is relative cohomology $H_X(\mathbf{X}, B)$ on \mathbf{X} relative to X .

In terms of $\text{Ho}_{\mathcal{C}}$ the familiar grading on cohomology is entirely implicit in the grading that the coefficient object \mathbf{A} carries for cases that \mathcal{V} may be interpreted as a category of graded or higher structures. To guarantee a consistent interpretation for which this is the case, we impose the additional condition that \mathcal{V} be a category with interval object.

Definition 3.3 (category with interval object) A category with interval object is a closed monoidal category with tensor unit I the terminal object – which we write $I \simeq \text{pt}$ – and equipped with an internal co-category of the form $\text{pt} \begin{array}{c} \xrightarrow{\sigma} \\ \xrightarrow{\tau} \end{array} \mathcal{I}$. We say \mathcal{I} is the interval object.


In particular for \mathcal{V} a category with interval object this means that

- there is a diagram of the form

$$\begin{array}{ccccc} & & \mathcal{I} \sqcup_{\text{pt}} \mathcal{I} & & \\ & \nearrow & & \nwarrow & \\ \mathcal{I} & & & & \mathcal{I} \\ \sigma \nearrow & \tau \nwarrow & & \sigma \nearrow & \tau \nwarrow \\ \text{pt} & & \text{pt} & & \text{pt} \\ & \searrow & & \swarrow & \\ & & \mathcal{I} & & \end{array}$$

with c co-associative and co-unital in the obvious sense;

- for every object B in a category \mathcal{C} enriched and powered over \mathcal{V} such that $[I, B] \simeq B$, there are canonical morphisms $B \xrightarrow{i} [I, B] \xrightarrow{d_0 \times d_1} B \times B$ and a composition morphism



which makes $[I, B]$ a category internal to \mathcal{C}_0 .

Definition 3.4 (directed and undirected objects) An object B in a category with interval object is undirected if $B \xrightarrow{i} [I, B]$ is a weak equivalence.

On undirected or *groupoidal* objects the computation of cohomology is usually easier, in particular if these arrange themselves into a *category of fibrant objects* in the sense of [7].

Definition 3.5 (compatible fibrant objects) A homotopical category with interval \mathcal{I} has a compatible structure of a category of fibrant objects if it is equipped with the structure of a category of fibrant objects in the sense of [7] such that the weak equivalences of both structures coincide and such that for every object B the object $[I, B]$ with its canonical structure morphisms is a path object for B .

Since in a category of fibrant objects the morphisms $B \xrightarrow{i} [I, B]$ are weak equivalences, the objects of such a category are *undirected*.

An important structure present in a category of fibrant objects is the factorization lemma.

Lemma 3.6 (factorization lemma [7]) In a category of fibrant objects for every morphism $\mathbf{X} \longrightarrow \mathbf{A}$

there is a diagram such that the acyclic fibration p has a section $\sigma : \mathbf{X} \rightarrow \mathbf{Y}$.

It follows that for fixed \mathbf{A} the functor $\text{hom}(-, \mathbf{A}) : \mathcal{C}_0^{\text{op}} \rightarrow \mathcal{V}$ is *homotopical* (respects weak equivalences) and hence already coincides on objects with its right-derived functor if it sends acyclic fibrations to weak equivalences. The objects \mathbf{A} for which this is true at least for acyclic fibrations $\pi : \mathbf{Y} \xrightarrow{\simeq} X$ over representables X are called ∞ -stacks. For each such π the condition that $\text{hom}(-, \mathbf{A})$ be homotopical is that

$$\text{hom}(\pi, \mathbf{A}) : \mathbf{A}(X) \simeq \text{hom}(X, \mathbf{A}) \xrightarrow{\simeq} \text{hom}(\mathbf{Y}, \mathbf{A})$$

is a weak equivalence. This is the descent condition on \mathbf{A} and $\text{hom}(\mathbf{Y}, \mathbf{A})$ is the \mathcal{V} -object of descent data of \mathbf{A} along π . Conversely, this means that the passage from \mathcal{C}_0 to $\text{Ho}\mathcal{C}_0$ is to be addressed as ∞ -stackification.

Finally, sometimes we need to consider sub-categories of \mathcal{C}_0 which are *pointed*:

Definition 3.7 (pointed category) A category is pointed if it has a *0-object*: an initial object isomorphic to a terminal object.

A \mathcal{V} -enriched homotopical category \mathcal{C} with interval object is our general situation. A compatible structure of fibrant objects puts us in an undirected *groupoidal* context. Pointedness further puts us in the context of an *abelian* category. In the following let $\mathcal{P}_0 \subset \mathcal{F}_0 \subset \mathcal{C}_0$ be inclusions of full subcategories, with \mathcal{F}_0 a category of fibrant objects compatible with the homotopical structure induced \mathcal{C}_0 and \mathcal{P}_0 a pointed compatible category of fibrant objects.

Only for some applications, such as for discussion of the *extension problem* do we need to assume the full structure of a model category.

3.1 Pointed objects

In *abelian* homotopy theory one considers (for instance section 4 of [7]) the case that in the homotopical category exists an initial object isomorphic to the terminal one. This makes all objects uniquely *pointed* in the sense of the following definition and allows to define *loop group* objects of all objects.

To retain a nonabelian (directed) setup we generalize this to the case where there need not be an initial object isomorphic to the terminal object by instead considering objects equipped with a specified point. This leads to loop *monoids* and will allow us in particular to discuss not only higher principal bundles but also higher vector bundles.

Definition 3.8 (pointed object) *The category of pointed objects in \mathcal{C}_0 is the under-category $\mathcal{C}_0 \backslash \text{pt}$: a pointed object is a morphism $\text{pt} \xrightarrow{\text{pt}_F} F$ and a morphism of pointed objects is a morphism $f : E \rightarrow F$ making the diagram*

$$\begin{array}{ccc} & \text{pt} & \\ \text{pt}_E \swarrow & & \searrow \text{pt}_F \\ E & \xrightarrow{f} & F \end{array}$$

commute.

Lemma 3.9 *The identity $\text{Id} : \text{pt} \rightarrow \text{pt}$ is both the terminal as well as the initial object of $\mathcal{C}_0 \backslash \text{pt}$.*

This says that $\mathcal{C}_0 \backslash \text{pt}$ is pointed. In the following we consider operations on pointed objects but in all of \mathcal{C}_0 , which itself need not be pointed.

Definition 3.10 (kernel and cokernel) *The kernel $\ker(f)$ of a morphism $f : A \rightarrow B$ into a pointed object B is the pullback*

$$\begin{array}{ccc} \ker(f) & \longrightarrow & \text{pt} \\ \downarrow & \lrcorner & \downarrow \text{pt}_B \\ A & \xrightarrow{f} & B \end{array} .$$

The cokernel $\text{coker}(f)$ of a morphism $f : B \rightarrow A$ is the pushout

$$\begin{array}{ccc} B & \xrightarrow{f} & A \\ \downarrow & & \downarrow \\ \text{pt} & \xrightarrow{\text{pt}_{\text{coker}(f)}} & \text{coker}(f) \end{array} .$$

The cokernel is naturally a pointed object $\text{pt} \xrightarrow{\text{pt}_{\text{coker}(f)}} \text{coker}(f)$ as indicated, the kernel is naturally pointed if A is a pointed object and f a morphism of pointed objects, with the point $\text{pt} \xrightarrow{\text{pt}_{\ker(f)}} \ker(f)$ given by the

universal dashed morphism in

$$\begin{array}{ccc}
 \text{pt} & & \\
 \downarrow \text{pt}_{\ker(f)} & \searrow & \\
 \ker(f) & \longrightarrow & \text{pt} \\
 \downarrow & \lrcorner & \downarrow \text{pt}_F \\
 A & \xrightarrow{f} & B \\
 \uparrow \text{pt}_B & & \\
 \text{pt} & &
 \end{array}$$

The kernel of a fibration of pointed objects is called its fiber.

If we have an ambient structure of a model category then by replacing limits and colimits in the above with their homotopy coherent versions, we obtain the homotopy kernel $\text{hoker}(f)$ and the homotopy cokernel $\text{hocoker}(f)$.

Definition 3.11 (monoid of loops) *The monoid of loops $\Omega_{\text{pt}}B$ of a pointed object $\text{pt} \xrightarrow{\text{pt}_B} B$ is the fiber of $[\mathcal{I}, B] \xrightarrow{d_0 \times d_1} B \times B$ with $B \times B$ equipped with its canonical point $\text{pt} \xrightarrow{\text{pt}_B \times \text{pt}_B} B \times B$, i.e. the pullback*

$$\begin{array}{ccc}
 \Omega_{\text{pt}}B & \longrightarrow & \text{pt} \\
 \downarrow & \lrcorner & \downarrow \text{pt}_B \times \text{pt}_B \\
 [\mathcal{I}, B] & \xrightarrow{d_0 \times d_1} & B \times B
 \end{array}$$

equipped with the structure of a monoid induced from the structure of a co-category on \mathcal{I} .

Lemma 3.12 *In the case that \mathcal{V} is a category of fibrant objects with initial object isomorphic to the terminal one, the image of $\Omega_{\text{pt}}B$ in the homotopy category is the loop group object considered in section 4 of [7].*

3.2 Universal fiber bundles

Definition 3.13 (universal fiber bundles) *For $\text{pt} \xrightarrow{\text{pt}_B} B$ a pointed object, the corresponding universal B-bundle is the morphism $p : \mathbf{E}_{\text{pt}}B \longrightarrow B$ with $\mathbf{E}_{\text{pt}}B$ the kernel of $[\mathcal{I}, F] \xrightarrow{d_0} F$ and p the composite*

$$\begin{array}{ccc}
 \mathbf{E}_{\text{pt}}B & \longrightarrow & \text{pt} \\
 \downarrow & \lrcorner & \downarrow \text{pt}_B \\
 B^{\mathcal{I}} & \xrightarrow{d_0} & B \\
 \downarrow d_1 & & \\
 B & &
 \end{array}$$

Lemma 3.14 *The fiber of the universal B-bundle $p : \mathbf{E}_{\text{pt}}B \rightarrow B$ is the monoid of loops $\Omega_{\text{pt}}B$.*

Proof. Evidently

$$\begin{array}{ccc}
 \Omega_{\text{pt}}B & \longrightarrow & B^{\mathcal{I}} \\
 \downarrow & \lrcorner & \downarrow d_0 \times d_1 \\
 \text{pt} & \xrightarrow{\text{pt}_B \times \text{pt}_B} & B \times B
 \end{array}
 \Leftrightarrow
 \begin{array}{ccc}
 \Omega_{\text{pt}}B & \longrightarrow & \mathbf{E}_{\text{pt}}B \\
 \downarrow & \lrcorner & \downarrow p \\
 \text{pt} & \xrightarrow{\text{pt}_B} & B
 \end{array}$$

□

Notice that

Definition 3.15 *There is a natural action $\rho : \mathbf{E}_{\text{pt}} \times \Omega_{\text{pt}} B \rightarrow \mathbf{E}_{\text{pt}} B$ induced from the co-category structure on \mathcal{I} .*

Lemma 3.16 *This action is a morphism of bundles*

$$\begin{array}{ccc} \mathbf{E}_{\text{pt}} B \times \Omega_{\text{pt}} B & \xrightarrow{\rho} & \mathbf{E}_{\text{pt}} B \\ & \searrow p \circ p_1 & \swarrow p \\ & & B \end{array}$$

Proposition 3.17 *If \mathcal{V} is compatibly a category of fibrant objects, then for every object B the universal B -bundle*

1. *is an acyclic fibration over the point, $\mathbf{E}_{\text{pt}} B \xrightarrow{\simeq} \text{pt}$;*
2. *sits in a sequence $\Omega_{\text{pt}} B \xrightarrow{i} \mathbf{E}_{\text{pt}} B \xrightarrow{p} B$ with p a fibration and i the fiber p .*

Proof. Recall from [7] that in a category of fibrant objects the maps $d_0, d_1 : [\mathcal{I}, B] \xrightarrow{\simeq} B$ out of a path object are acyclic fibrations and that acyclic fibrations are preserved under pullback. Hence from

$$\begin{array}{ccc} \mathbf{E}_{\text{pt}} & \xrightarrow{\simeq} & \text{pt} \\ \downarrow & \lrcorner & \downarrow \text{pt}_B \\ B^{\mathcal{I}} & \xrightarrow{d_0} & B \\ \downarrow & \lrcorner & \downarrow \\ B & & B \end{array}$$

p (curved arrow from \mathbf{E}_{pt} to B)

we find the acyclic fibration from $\mathbf{E}_{\text{pt}} B$ to pt .

That p is a fibration is a special case of the proof of the “factorization lemma” in [7]:

Using that pullbacks of fibrations are again fibrations, we obtain for all fibrant objects C and D that projections out of their product are fibrations

$$\begin{array}{ccc} C \times D & \xrightarrow{\text{pr}_2} & D \\ \text{pr}_1 \downarrow & & \downarrow \\ C & \longrightarrow & \text{pt} \end{array}$$

and for all morphisms $f : C \rightarrow D$ that the top left vertical morphisms in the double pullback square

$$\begin{array}{ccc} C \times_D D^{\mathcal{I}} & \longrightarrow & D^{\mathcal{I}} \\ \text{id} \times d_1 \downarrow & & \downarrow d_0 \times d_1 \\ C \times D & \xrightarrow{f \times \text{Id}} & D \times D \\ \text{pr}_1 \downarrow & & \downarrow \text{pr}_1 \\ C & \xrightarrow{f} & D \end{array} \quad \cdot$$

d_0 (curved arrow from $D^{\mathcal{I}}$ to D)

is a fibration. Since composites of two fibrations are fibrations, it follows that p in

$$\begin{array}{ccc} C \times_D D^{\mathcal{I}} & \longrightarrow & D^{\mathcal{I}} \\ \text{id} \times d_1 \downarrow & \searrow & \downarrow d_0 \times d_1 \\ C \times D & \xrightarrow{f \times \text{Id}} & D \times D \xrightarrow{\text{pr}_2} D \end{array}$$

p (curved arrow from $D^{\mathcal{I}}$ to D)

is a fibration. Taking f to be $\text{pt} \xrightarrow{\text{pt}_B} B$ this yields the desired statement for p .

Finally, that i is the kernel of p is lemma ??.

□

3.3 Cocycles and bundles

Now consider a subcategory $\mathcal{F}_0 \subset \mathcal{C}_0$ of fibrant objects. By the central theorem in [7] morphisms in the homotopy category $\text{Ho}_{\mathcal{F}_0}$ are represented already by single spans whose left leg is an acyclic fibration.

Definition 3.18 (anamorphisms) An anamorphism or cocycle on X with values in A is a span

$$\begin{array}{ccc} \hat{X} & \longrightarrow & A \\ & \downarrow \simeq & \\ & X & \end{array} .$$

Lemma 3.19 *Anamorphisms have a consistent composition induced by pullback*

$$\begin{array}{ccccc} g_1^* \hat{B} & \longrightarrow & \hat{B} & \xrightarrow{g_2} & C \\ \downarrow \simeq & & \downarrow \simeq & & \\ \hat{A} & \xrightarrow{g_1} & B & & \\ \downarrow \simeq & & & & \\ A & & & & \end{array} .$$

which is associative and unital up to isomorphism of spans.

Definition 3.20 (bundles obtained from cocycles) Given a cocycle $X \xleftarrow{\simeq} \hat{X} \xrightarrow{g} B$ into a pointed object $\text{pt} \xrightarrow{\text{pt}_B} B$ the corresponding B -bundle $p : g^* \mathbf{E}_{\text{pt}} B \longrightarrow X$ is the pullback

$$\begin{array}{ccc} g^* \mathbf{E}_{\text{pt}} B & \longrightarrow & \mathbf{E}B \\ \downarrow & & \downarrow \\ p \left(\hat{X} \right) & \xrightarrow{g} & B \\ \downarrow \simeq & & \\ X & & \end{array}$$

This bundle inherits an action $\rho : (g^* \mathbf{E}_{\text{pt}} B) \times \Omega_{\text{pt}} B \rightarrow g^* \mathbf{E}_{\text{pt}} B$ of the monoid of loops from the commutativity of

$$\begin{array}{ccccc} g^* \mathbf{E}B \times \Omega_{\text{pt}} B & \longrightarrow & \mathbf{E}B \times \Omega_{\text{pt}} B & \xrightarrow{\rho} & \mathbf{E}B \\ \downarrow p_1 & & \downarrow p_1 & & \downarrow \\ g^* \mathbf{E}B & \longrightarrow & \mathbf{E}B & & \\ \downarrow & & \downarrow & & \swarrow \\ \hat{X} & \xrightarrow{g} & B & & \end{array}$$

Lemma 3.21 *This induced action is still a morphism of bundles*

$$\begin{array}{ccc}
 (g^*\mathbf{E}_{\text{pt}}B) \times \Omega_{\text{pt}}B & \xrightarrow{\rho} & g^*\mathbf{E}_{\text{pt}}B \\
 \searrow_{p \circ p_1} & & \swarrow_p \\
 & X &
 \end{array}$$

Definition 3.22 (fiber bundle) *We say the morphism $P \longrightarrow X$ equipped with an action of $\Omega_{\text{pt}}B$ is a B -fiber bundle if there is a cocycle $X \xleftarrow{\simeq} \hat{X} \xrightarrow{g} B$ and a weak equivalence $g^*\mathbf{E}_{\text{pt}}B \xrightarrow{\simeq} P$ respecting the $\Omega_{\text{pt}}B$ -action on both sides.*

Proposition 3.23 (fiber bundle trivializes over itself) *In the category \mathcal{F}_0 of fibrant objects, every fiber bundle $P \rightarrow X$ becomes trivializable when pulled back along itself.*

Proof. For $X \xleftarrow{\simeq} \hat{X} \longrightarrow B$ a cocycle characterizing the bundle $P \rightarrow X$ we obtain the pullback diagram

$$\begin{array}{ccccc}
 P \xleftarrow{\simeq} g^*\mathbf{E}_{\text{pt}}B & \xrightarrow{\quad} & \mathbf{E}_{\text{pt}}B & \twoheadrightarrow & \text{pt} \\
 \downarrow & & \downarrow & & \downarrow \\
 & & g^*B^{\mathcal{I}} & \xrightarrow{\quad} & B^{\mathcal{I}} \xrightarrow{d_0} B \\
 & & \downarrow \simeq & & \downarrow d_1 \\
 X \xleftarrow{\simeq} \hat{X} & \xrightarrow{g} & B & &
 \end{array}$$

The cocycle g pulled back to P is represented by the morphism from $g^*\mathbf{E}_{\text{pt}}B$ to the B at the bottom. The right part of the diagram says that this is homotopic to a map factoring through the point. \square

Definition 3.24 (associated bundle) *Let $g^*\mathbf{E}_{\text{pt}}B \rightarrow X$ be a B -bundle as above and let $\rho : B \rightarrow F$ be a morphism in \mathcal{C}_0 to a pointed object $\text{pt} \xrightarrow{\text{pt}_F} F$ not necessarily fibrant. Then we call ρ a representation of B on F and call the pullback*

$$\begin{array}{ccccc}
 g^*\rho^*\mathbf{E}_{\text{pt}}F & \longrightarrow & \rho^*\mathbf{E}_{\text{pt}}F & \longrightarrow & \mathbf{E}_{\text{pt}}F \\
 \downarrow & & \downarrow & & \downarrow \\
 \hat{X} & \xrightarrow{g} & B & \xrightarrow{\rho} & F \\
 \downarrow & & & & \\
 X & & & &
 \end{array}$$

the associated bundle, associated by ρ to P .

So in particular $\rho^*\mathbf{E}_{\text{pt}}F$ is the F bundle ρ -associated to the universal B -bundle.

3.4 Sections and homotopies

One way to think of a section of an ω -bundle is as a morphism from a certain trivial ω -bundle into it. The following formalizes this and then provides reformulations of this notion which are useful later on in section 5.

Definition 3.25 (section) A section σ of an F -cocycle $\hat{X} \xrightarrow{\nabla} F$ is a directed homotopy from the trivial F -cocycle with fiber pt_F into ∇ .

$$\Gamma(\nabla) := \left\{ \begin{array}{ccc} & \hat{X} & \\ \swarrow & & \searrow \text{Id} \\ \text{pt} & \overset{\sigma}{=} & \hat{X} \\ \searrow \text{pt}_F & & \swarrow \nabla \\ & F & \end{array} \right\}$$

Proposition 3.26 If the F -cocycle ∇ is ρ -associated to a B -cocycle $\hat{X} \xrightarrow{g} B$ then sections of ∇ are equivalently lifts of g through $\rho^* \mathbf{E}_{\text{pt}F} \rightarrow B$

$$\Gamma(\nabla) \simeq \left\{ \begin{array}{ccc} & \rho^* \mathbf{E}_{\text{pt}F} & \\ \swarrow \sigma & & \downarrow \\ \hat{X} & \xrightarrow{g} & B \end{array} \right\}.$$

Proof. First rewrite

$$\left\{ \begin{array}{ccc} & \hat{X} & \\ \swarrow & & \downarrow g \\ \text{pt} & \overset{\sigma}{=} & B \\ \searrow \text{pt}_F & & \downarrow \rho \\ & F & \end{array} \right\} \simeq \left\{ \begin{array}{ccc} & \text{pt} & \xrightarrow{\text{pt}_F} F \\ \swarrow & & \uparrow d_0 \\ \hat{X} & \overset{\sigma}{\dashrightarrow} & [\mathcal{I}, F] \\ \searrow g & & \downarrow d_1 \\ & B & \xrightarrow{\rho} F \end{array} \right\}$$

using the characterization of right (directed) homotopies by the (directed) path object $[\mathcal{I}, F]$. Using the universal property of $\mathbf{E}_{\text{pt}F}$ as a pullback this yields

$$\dots \simeq \left\{ \begin{array}{ccc} & \mathbf{E}_{\text{pt}F} & \\ \swarrow \sigma & & \downarrow \\ \hat{X} & \xrightarrow{g} B & \xrightarrow{\rho} F \end{array} \right\} \simeq \left\{ \begin{array}{ccc} & \rho^* \mathbf{E}_{\text{pt}F} & \\ \swarrow \sigma & & \downarrow \\ \hat{X} & \xrightarrow{g} & B \end{array} \right\}.$$

□

A third way to think about sections comes from observing that since a directed homotopy between

two cocycles $\hat{X} \begin{array}{c} \xrightarrow{g_1} \\ \parallel \eta \\ \xrightarrow{g_2} \end{array} F$ is given by a morphism $\hat{X} \xrightarrow{\eta} [\mathcal{I}, F]$ it can itself be regarded as an $[\mathcal{I}, F]$ -cocycle.

Definition 3.27 (universal $F^{\mathcal{I}}$ -bundle) The object $[\mathcal{I}, F]$ is naturally equipped with the point $\text{pt}_{[\mathcal{I}, F]}$ defined by

$$\begin{array}{ccc} \text{pt} \xrightarrow{\text{pt}_F} F & \longrightarrow & [\mathcal{I}, F] \\ \text{pt}_{[\mathcal{I}, F]} \curvearrowright & & \end{array} \Rightarrow \begin{array}{ccc} \text{pt} \xrightarrow{\text{pt}_{[\mathcal{I}, F]}} [\mathcal{I}, F] & \xrightarrow{d_0 \times d_1} & F \times F \\ \text{pt}_F \times \text{pt}_F \curvearrowright & & \end{array} .$$

We write $\mathbf{E}_{\text{pt}}([\mathcal{I}, F]) \longrightarrow [\mathcal{I}, F]$ for the corresponding universal $[\mathcal{I}, F]$ -bundle according to definition 3.13.

Notice the commutativity of the diagram

$$\begin{array}{ccccc} \mathbf{E}_{\text{pt}}([\mathcal{I}, F]) & \longrightarrow & [\mathcal{I}, [\mathcal{I}, F]] & \xrightarrow{[\mathcal{I}, d_i]} & [\mathcal{I}, F] \\ \downarrow & & \downarrow d_0 & & \downarrow d_0 \\ \text{pt} & \xrightarrow{\text{pt}_{[\mathcal{I}, F]}} & [\mathcal{I}, F] & \xrightarrow{d_i} & F \\ & \text{pt}_F \curvearrowright & & & \end{array}$$

for $i = 0$ and $i = 1$. The right square commutes by the functoriality of $[\mathcal{I}, -]$, the left square and the bottom triangle by definition 3.27 of the universal $[\mathcal{I}, F]$ -bundle.

Definition 3.28 Let $\mathbf{E}_{\text{pt}} F \xleftarrow{\mathbf{E}_{\text{pt}} d_0} \mathbf{E}_{\text{pt}}([\mathcal{I}, F]) \xrightarrow{\mathbf{E}_{\text{pt}} d_1} \mathbf{E}_{\text{pt}} F$ be the universal morphisms induced from the commutativity of the outermost rectangle of the above diagram in view of the universal property of $\mathbf{E}F$ as a pullback.

Proposition 3.29 The morphism $\mathbf{E}d_0 \times \mathbf{E}d_1$ covers the morphism $[\mathcal{I}, F] \xrightarrow{d_0 \times d_1} F \times F$ of base spaces:

$$\begin{array}{ccccc} \mathbf{E}_{\text{pt}} F & \xleftarrow{\mathbf{E}d_0} & \mathbf{E}([\mathcal{I}, F]) & \xrightarrow{\mathbf{E}d_1} & \mathbf{E}_{\text{pt}} F \\ \downarrow & & \downarrow & & \downarrow \\ F & \xleftarrow{d_0} & [\mathcal{I}, F] & \xrightarrow{d_1} & F \end{array} .$$


Proof. By inspection of the above commuting diagram. □

This is sometimes called a concordance of bundles.

In total this yields for every homotopy of F -cocycles a span of the corresponding bundles.

Definition 3.30 (associated span of bundles) For $\begin{array}{ccc} & \hat{X} & \\ \iota \nearrow & \Downarrow & \searrow \nabla_1 \\ \hat{Q} & & F \\ \iota' \searrow & \Downarrow & \nearrow \nabla_2 \\ & \hat{Y} & \end{array}$ a directed homotopy of F -cocycles, we


say that the rear of the joint pullback diagram




is the associated span of ω -bundles.

From definition 3.11 notice the following observation:

Lemma 3.31 *In the case that the homotopy in question is a section*



associated span of bundles is of the form




with $\Omega_{\text{pt}} F$ the monoid of loops from

definition 3.11.

Remark on groupoidification. In the case that \mathcal{F}_0 is a category of ∞ -groupoids, this realizes a section of an associated bundle as an ∞ -groupoid over the total space ∞ -groupoid of the associated ∞ -bundle, and equipped with a map to the ground ∞ -monoid. Since this is the description of vectors in the context of *groupoidification* [1, 2] it motivates the following definition.

Definition 3.32 (generalized sections of associated ω -bundles) *Given an F -bundle $V := \nabla^* \mathbf{E}_{\text{pt}} F$,*

its generalized sections are spans $|\Psi\rangle := \Omega_{\text{pt}} F \leftarrow \Psi \rightarrow V$ and its generalized co-sections are spans $\langle \Psi| :=$



3.5 Extension and lifting problem

3.5.1 Extension

Assume for the following that on \mathcal{C}_0 we have a full structure of a model category.

Consider extensions \underline{g} of morphisms g along morphisms i

$$\begin{array}{ccc} X & \xrightarrow{g} & A \\ \downarrow i & \nearrow \underline{g} & \\ \underline{X} & & \end{array}$$

Definition 3.33 (equivariant structure and flat connection) An extension through a morphism $i : A \hookrightarrow X$ which is an isomorphism from the points of A to the points of X

$$[\text{pt}, A] \xrightarrow{i_*} [\text{pt}, X]$$

is called an *i -equivariant structure* or an *i -flat connection*.

Theorem 3.34 (long exact sequence for extensions) For every object B for which there is a diagram

$$\begin{array}{ccc} B & & \\ \downarrow i & & \\ \mathbf{W}B & \xrightarrow{\cong} & \text{pt} \\ \downarrow p & & \\ \text{hocoker}(i) & & \end{array}$$

and for every extension $\iota : X \rightarrow \underline{X}$ there is a morphism $\delta : H(X, B) \rightarrow H_X(\underline{X}, \text{hocoker}(i))$ whose kernel is the image of ι^* . If furthermore i is the homotopy kernel of p then then image of δ is the kernel of $H_X(\underline{X}, \text{hocoker}(i)) \rightarrow H(X, \text{hocoker}(i))$ so that we get a semi-long exact sequence in cohomology

$$H_X(\underline{X}, B) \longrightarrow H(\underline{X}, B) \xrightarrow{\iota^*} H(X, B) \xrightarrow{\delta} H_X(\underline{X}, \text{hocoker}(i)) \longrightarrow H(\underline{X}, \text{hocoker}(i)) \longrightarrow \dots$$

Proof. (the idea)

The connecting homomorphism δ is obtained by first constructing a diagram

$$\begin{array}{ccccc} X & \xleftarrow{\cong} & \hat{X} & \xrightarrow{g} & B \\ \downarrow i & & \downarrow \exists i & \nearrow \exists F & \downarrow i \\ \underline{X} & \xleftarrow{\cong} & \hat{X} & \xrightarrow{\exists F} & \mathbf{W}B \\ & & \searrow \delta g & & \downarrow p \\ & & & & \text{hocoker}(i) \end{array}$$

and then setting $\delta g := p \circ F$. Here the existence of the cofibration \hat{i} follows from the general factorization property in a model category, while F is constructed as the dashed morphism in

$$\begin{array}{ccc} \hat{X} & \longrightarrow & B \\ \downarrow \hat{i} & & \downarrow i \\ \hat{X} & \xrightarrow{F} & \mathbf{W}B \\ \downarrow \hat{i} & \nearrow & \downarrow \cong \\ \hat{X} & \longrightarrow & \text{pt} \end{array}$$

which exists by the lifting property of cofibrations. By the property of a homotopy colimit δg is precisely the obstruction for the dashed lift to exist. That every cocycle in $H(\underline{X}, \text{coker}(i))$ which trivializes under i^* arises this way follows by using that $i = \text{hoker}(p)$... \square

Definition 3.35 (curvature and characteristic classes) Here F is the curvature, δg the characteristic classes of g for the extension along i .

3.5.2 Lifting

...

3.5.3 Local semi-trivializations

Proposition 3.23 states that every B -bundle trivializes over its own total space. If \hat{B} is part of an exact sequence $A \xrightarrow{i} \hat{B} \xrightarrow{p} B$, with i the homotopy kernel of p , then there is a relative version of this statement, which states that a \hat{B} bundle becomes equivalent to an A -bundle with a certain \hat{B} -equivariance on the total space of the underlying B -bundle.

Definition 3.36 (relative equivariance) For $\pi : P \rightarrow X$ a B -bundle we say a cocycle $P \dashrightarrow A$ on its total space is relatively equivariant with respect to the above sequence if

- there exists \hat{B} -cocycle \underline{g} on X
- and a homotopy $\pi^* \underline{g} \Rightarrow p^* g$
- and such that P is classified by $p^* g$

Proposition 3.37 (local semi-trivialization) \hat{B} -cohomology on X is in bijection with relatively equivariant A -cohomology on the underlying B -bundles.

Proof. Consider

$$\begin{array}{ccccccc}
 & & & & A & & \\
 & & & & \swarrow & \searrow & \\
 & & & & \hat{B} & & \\
 & & & & \swarrow & \searrow & \\
 P & \xleftarrow{\simeq} & g^* \mathbf{E}_{\text{pt}} B & \xrightarrow{\quad} & \mathbf{E}_{\text{pt}} B & \twoheadrightarrow & \text{pt} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 P//G & \xleftarrow{\simeq} & g^* [\mathcal{I}, B] & \xrightarrow{\quad} & [\mathcal{I}, B] & \xrightarrow{d_0} & B \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X & \xleftarrow{\simeq} & \hat{X} & \xrightarrow{g} & B & & \\
 & & & & \downarrow & & \\
 & & & & d_1 & &
 \end{array}$$

\square

4 Examples and Applications

4.1 Homotopical contexts

- $\mathcal{V} = \text{SimplicialSets}$: the theory of $\text{Ho}_{[\text{S}^{\text{op}}, \mathcal{V}]}$ developed in great detail in series of articles by Toën [?].

4.2 Closed monoidal homotopical categories

- Cat with standard tensor and folk model structure
- 2Cat with Gray tensor and Lack model structure
- ω Cat with Crans-Gray tensor and folk model structure (need to confirm some axioms of closed monoidal homotopical).
- ω Groupoids with Brown-Golasiski tensor and model structure (need to confirm some axioms of closed monoidal homotopical).

4.3 Pointed objects

- one-object ω -groupoids $\text{pt}! \xrightarrow{\exists!} \mathbf{BG}$
- category of vector spaces, point maps to ground field $\text{pt} \xrightarrow{\text{pt} \mapsto k} \text{Vect}_k$
- similarly for higher vector spaces,

4.4 Monoid of loops

1. for $\text{pt} \xrightarrow{\exists!} \mathbf{BG}$ we have $\Omega \mathbf{BG} = G$.
2. for $\text{pt} \xrightarrow{\text{pt} \mapsto k} \text{Vect}_k$ we have $\Omega_{\text{pt}} \text{Vect} = k$,
3. for $\text{pt} \longrightarrow 2\text{Vect}_k$ we have $\Omega_{\text{pt}} 2\text{Vect} = \text{Vect}_k$, etc.

4.5 Universal bundles

1. for $B = \mathbf{BG}$ and G a 1-group the universal B -bundle is $\mathbf{EG} := G//G$ is the action groupoid of G acting on itself and the sequence $G \longrightarrow \mathbf{EG} \longrightarrow \mathbf{BG}$ maps under nerve and topological realization to the universal G -bundle in its incarnation in topological spaces. This is discussed in [25] as a preparation for the following example.
2. for $B = \mathbf{BG}$ and G a 2-group or bi-group [4] shows that \mathbf{EG} is the action bigroupoid of G acting on itself. we show that \mathbf{EG} is the universal 2-bundle (sketched in [25]).

4.6 Bundles

- $\rho : \mathbf{BG} \rightarrow \text{Vect}$ a representation, then $\rho^* \mathbf{E}_{\text{pt}} \text{Vect}$ is the action groupoid (ρ -associated vector bundle of \mathbf{EG})
- pullback along G_1 -cocycle g weakly equivalent to G -principal bundle;
- pullback along G_2 -cocycle g weakly equivalent to G_2 -principal 2-bundle (Bartels, Baković, Wockel);
- combined pullback $g^* \circ \rho^*$ weakly equivalent to associated bundle

etc. pp.

4.7 Lifting problems

4.8 Extension problem

- $X \hookrightarrow X//G$: ordinary equivariance under G -action; connection homomorphism computes obstruction to having equivariant structure;
- $X \hookrightarrow \Pi(X)$: flat connection, connection homomorphism computes curvature (characteristic forms);

4.9 Equivariance on semi-total spaces

- for $\mathbf{BU}(1) \rightarrow \text{AUT}(U(1)) \rightarrow \mathbb{Z}_2$ this yields Jandl gerbes [34]
- for $\mathbf{BU}(1) \rightarrow \text{String}(G) \rightarrow G$ this yields String bundle gerbes on the total space of a G -bundle
- etc.

5 Quantization of nonabelian cocycles to σ -models

We want to think of a ρ -associated F -cocycle $\hat{X} \xrightarrow{\nabla} F$ and the corresponding F -bundle $\nabla^* \mathbf{E}_{\text{pt}} F \longrightarrow X$ as a *background field* (a generalization of an electromagnetic field) on X to which a higher dimensional *fundamental brane* – such as a *particle*, a *string* or a *membrane* – propagating on X may *couple*.

We now propose a formalization in the context of homotopical cohomology of what it means to *quantize* such a background field to obtain the corresponding σ -model quantum field theory as a functorial QFT (as described in [31] and references given there). Our constructions are motivated by and supposed to implement and generalize the considerations of [15, 42] and make contact with [24].

We define a notion of *parameter space* or *worldvolume* category and a notion of *background field* over a *target space* coming from an ω -bundle. This pair of data we call a σ -*model*. We show how this data induces a functor from the parameter space category to spans in ω -groupoids. Using methods from *groupoidification* [1, 2] we show that these spans represent linear maps which deserve to be addressed as *propagation* in the quantum field theory induced by the σ -model.

5.1 σ -Models

In the following $B := \mathbf{BG}$ denotes an object of \mathcal{F}_0 which we think of as modelling a one object ∞ -groupoid the automorphisms of whose single object form the ∞ -group G . At the moment this is just notation meant to be suggestive.

Definition 5.1 (background structure) A background structure for a σ -model is

- an ω -groupoid X called target space;
- an ω -group G , called the gauge group;

• a representation

$$\begin{array}{ccc} \rho^* \mathbf{E}G & \longrightarrow & \mathbf{E}_{\text{pt}} F \\ \downarrow & & \downarrow \\ \mathbf{B}G & \xrightarrow{\rho} & F \leftarrow \text{pt}_F \text{pt} \end{array} \quad \text{called the } \underline{\text{matter content}};$$

- an F -cocycle $\hat{X} \xrightarrow{\nabla} F$ ρ -associated to a G -principal cocycle on X , $\nabla : \hat{X} \xrightarrow{g} \mathbf{B}G \xrightarrow{\rho} F$,
 $\downarrow \simeq$
 X
 called the background field.

For brevity we shall indicate a background structure just as $(\hat{X} \xrightarrow{\nabla} F)$, leaving the choice of representation and the target space X weakly equivalent to the hypercover \hat{X} implicit.

Definition 5.2 (parameter space category) A parameter space ω -category Cob is a sub ω -category of $\text{Cospans}(\omega\text{Groupoids})$.

Definition 5.3 (σ -model) A σ -model is a pair consisting of a parameter space category and a background structure for a σ -model.

Definition 5.4 Given a σ -model with background structure $\nabla : \hat{X} \xrightarrow{\nabla} F$ and with parameter space Cob for every object $\Sigma \in \text{Cob}$ we say that

- $C(\Sigma) := [\Sigma, \hat{X}]$ is the space of fields over Σ ;
- The $[\Sigma, F]$ -cocycle $[\Sigma, \nabla] : [\Sigma, \hat{X}] \rightarrow [\Sigma, F]$ on the space of fields over Σ is the action functional over Σ .

Remark. One can identify $[\Sigma, \nabla]$ with the *transgression* of the cocycle g to the mapping space $[\Sigma, X]$. Examples showing that this canonical operation indeed reproduces the ordinary notion of transgression of cocycles are in [35, 36] and in our section 6.

In section 5.3 we construct for every σ -model its corresponding quantum field theory. This involves the notion of *branes* discussed in section 5.2.

5.2 Branes and bibranes

From the second part of definition 3.25 one sees that spaces spaces of sections of ω -bundles are given by certain morphisms between background fields pulled back to spans/correspondences of target spaces. From the diagrammatics this has an immediate generalization, which leads to the notion of *branes* and *bibranes*.

Definition 5.5 (branes and bibranes) A brane for a background structure $(\hat{X} \xrightarrow{\nabla} F)$ is a morphism

$\iota : \hat{Q} \rightarrow \hat{X}$ equipped with a section of the background field pulled back to \hat{Q} , i.e. a transformation


$$\begin{array}{ccc} & \hat{Q} & \\ \swarrow & & \searrow \iota \\ \text{pt} & \xrightarrow{V} & \hat{X} \\ \swarrow \text{pt}_F & & \searrow g \\ & F & \end{array} .$$

More generally, given two background structures $(\hat{X} \xrightarrow{\nabla} F)$ and $(\hat{X}' \xrightarrow{\nabla'} F)$, a bibrane between them


is a span $\begin{array}{ccc} & \hat{Q} & \\ \swarrow \iota & & \searrow \iota' \\ \hat{X} & & \hat{X}' \end{array}$ equipped with a transformation

$$\begin{array}{ccc} & \hat{Q} & \\ \swarrow \iota & & \searrow \iota' \\ \hat{X} & \xrightarrow{V} & \hat{X}' \\ \swarrow \nabla & & \searrow \nabla' \\ & F & \end{array} .$$

Bibranes may be composed –“fused” – along common background structures ($\hat{X} \xrightarrow{\nabla} F$): the composite or *fusion* of a bibrane V_1 on \hat{Q} with a bibrane V_2 on \hat{Q}' is the bibrane $V_1 \cdot V_2$ given by the diagram



Proposition 5.6 (composition of associated spans from fusion of bibranes) *The associated span of ω -groupoids corresponding, according to definition 3.30, to the fusion of two bibranes is the composition of the spans associated with each bibrane:*



Proof. By commutativity of pullbacks. □


Remark on groupoidification. Comparing with the remark above definition 3.32 we find that fusion of bibranes corresponds to composition of groupoidified linear maps.

If \hat{Q} carries further structure, the fused bibrane on $\hat{Q} \times_{\hat{X}'} \hat{Q}$ may be pushed down again to \hat{Q} .

Definition 5.7 Let B be a category enriched in the bicategory $\mathcal{V} := \text{Spans}(\omega\text{Categories})$ of spans in $\omega\text{Categories}$ and let F be an ω -category. Then the category of bibranes relative to B and F is given by:

- objects are background structures $\hat{X} \xrightarrow{\nabla} F$ for \hat{X} an object of B ;
- morphisms are bibranes on morphisms of B ;
- composition of morphisms is given by bibrane fusion followed by push-forward along the composition map in B .

A simple special case is a category $\hat{Q} \begin{smallmatrix} \xrightarrow{s} \\ \xrightarrow{t} \end{smallmatrix} \hat{X}$ internal to ω -groupoids, equivalently a monad in the bicategory of spans internal to $\omega\text{Groupoids}$, with composition operation the morphism of spans



Definition 5.8 (monoidal structure on bibranes) Given an internal category as above, and given a

background structure $\nabla : \hat{X} \rightarrow F$, the composite of two bibranes $\begin{array}{ccc} & \hat{Q} & \\ & \swarrow \searrow & \\ \hat{X} & \xrightarrow{V,W} & \hat{X} \\ & \swarrow \searrow & \\ & \hat{F} & \end{array}$ on \hat{Q} is the result of first

forming their composite bibrane on $\hat{Q} \times_{t,s} \hat{Q}$ and then pushing that forward along comp :

$$V \star W := \int_{\text{comp}} (s^*V) \cdot (t^*W).$$

Here for finite cases, which we concentrate on, push-forward is taken to be the right adjoint to the pullback in a proper context.

Remarks. Notice that branes are special cases of bibranes and that bibrane composition restricts to an action of bibranes on branes. Also recall that the sections of a cocycle on X are the same as the branes of this cocycle for $\iota = \text{Id}_X$.

The idea of bibranes was first formulated in [17] in the language of modules for bundle gerbes. We show in section 6.1.4 how this is reproduced within the present formulation. In its smooth L_∞ -algebraic version the idea also appears in [28].

5.3 Quantum propagation

Every σ -model with parameter space Cob and background structure $\hat{X} \xrightarrow{\nabla} F$ induces a functor

$$\exp\left(\int \nabla\right) : \text{Cob} \rightarrow \text{Spans}(\omega\text{Groupoids})$$

which sends

$$\exp\left(\int \nabla\right) : \begin{array}{c} \Sigma \\ \swarrow \iota \quad \searrow \tau \\ \Sigma_{\text{in}} \quad \Sigma_{\text{out}} \end{array} \mapsto \begin{array}{ccc} & [\Sigma, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma, F] & \\ \swarrow & & \searrow \\ [\Sigma_{\text{in}}, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma_{\text{in}}, \nabla] & & [\Sigma_{\text{out}}, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma_{\text{out}}, F] \end{array}$$

a morphism $\Sigma : \Sigma_{\text{in}} \rightarrow \Sigma_{\text{out}}$ in Cob to the span of ω -bundles associated to, definition 3.30, the bibrane on the span

$$\begin{array}{ccc} & [\Sigma, \hat{X}] & \\ \swarrow \iota^* & & \searrow \tau_* \\ [\Sigma, \hat{X}] & & [\Sigma, \hat{X}] \end{array}$$

which is induced by transgression of the background field:

$$\begin{array}{ccccccc} & & [\Sigma, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma, F] & & & & \\ & \swarrow & \downarrow & \searrow & & & \\ [\Sigma_{\text{in}}, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma_{\text{in}}, F] & & [\Sigma, \hat{X}] & & [\Sigma_{\text{out}}, \nabla]^* \mathbf{E}_{\text{pt}}[\Sigma_{\text{out}}, F] & & \\ & \swarrow \iota & \downarrow \mathbf{E}_{\text{pt}}[\Sigma, F] & \searrow \iota' & \downarrow \mathbf{E}_{\text{pt}}[\Sigma, F] & \searrow \iota'' & \\ & [\Sigma_{\text{in}}, \hat{X}] & \downarrow [\Sigma, \nabla] & & \downarrow [\Sigma, \nabla] & & [\Sigma_{\text{out}}, \hat{X}] \\ & \downarrow \mathbf{E}_{\text{pt}}[\Sigma_{\text{in}}, F] & \downarrow [\Sigma_{\text{in}}, \nabla] & \downarrow \mathbf{E}_{\text{pt}}[\Sigma_{\text{in}}, F] & \downarrow [\Sigma_{\text{out}}, \nabla] & \downarrow \mathbf{E}_{\text{pt}}[\Sigma_{\text{out}}, F] & \\ & [\Sigma_{\text{in}}, F] & \downarrow d_0 & & \downarrow d_1 & & [\Sigma_{\text{out}}, F] \end{array}$$

A state of the σ -model over Σ is a generalized section, definition 3.32 of $[\Sigma_a, \nabla]$ in $\mathcal{H}_a := \mathcal{H}([\Sigma_a, \nabla])$ and the propagation of along a morphism Σ is the map

$$\int_{\text{hom}(\Sigma, X)} \exp\left(\int \nabla\right) : \mathcal{H}_{\Sigma_{\text{in}}} \longrightarrow \mathcal{H}_{\Sigma_{\text{out}}}$$

induced by pull-push through the span $\exp(\int \nabla)(\Sigma)$.

An example is spelled out in section 6.1.6.

6 Examples and applications

We start with some simple applications to illustrate the formalism and then exhibit some useful constructions in the context of finite group quantum field theory.

6.1 General examples

6.1.1 Ordinary vector bundles

Let G be an ordinary group, hence a 1-group, and denote by $F := \text{Vect}$ the 1-category of vector spaces over some chosen ground field k . A linear representation ρ of G on a vector space V is indeed the same thing as a functor $\rho : \mathbf{BG} \rightarrow \text{Vect}$ which sends the single object of \mathbf{BG} to V .

The canonical choice of point $\text{pt}_F : \text{pt} \rightarrow \text{Vect}$ is the ground field k , regarded as the canonical 1-dimensional vector space over itself. Using this we find

- from definition ?? that the *ground ω -monoid* in this case is just the ground field itself, $K = k$,
- from definition ?? that the *universal Vect-bundle* is $\mathbf{E}_{\text{pt}} \text{Vect} = \text{Vect}_*$, the category of *pointed* vector spaces with $\text{Vect}_* \longrightarrow \text{Vect}$ the canonical forgetful functor;
- from definition ?? that the ρ -associated vector bundle to the universal G -bundle is $V//G \longrightarrow \mathbf{BG}$, where $V//G := (V \times G \xrightarrow[\rho]{p_1} V)$ is the *action groupoid* of G acting on V , the weak quotient of V by G ;
- From definition 3.25 that for $g : X \xrightarrow{q} \mathbf{BG}$ a cocycle describing a G -principal bundle and for V the corresponding ρ -associated vector bundle according to definition ??, that sections $\sigma \in \Gamma(V)$ are precisely sections of V in the ordinary sense.

6.1.2 Group algebras and category algebras from bibrane monoids

In its simplest version the notion of monoidal bibranes from section 5.2 reproduces the notion of *category algebra* $k[C]$ of a category C , hence also that of a *group algebra* $k[G]$ of a group G . Recall that the category algebra $k[C]$ of C is defined to have as underlying vector space the span of C_1 , $k[C] = \text{span}_k(C_1)$, where the product is given on generating elements $f, g \in C_1$ by

$$f \cdot g = \begin{cases} g \circ f & \text{if the composite exists} \\ 0 & \text{otherwise} \end{cases}$$

To reproduce this as a monoid of bibranes in the sense of section 5.2, take the category of fibers in the sense of section ?? to be $F = \text{Vect}$ as in section 6.1.1. Consider on the space (set) of objects, C_0 , the trivial line bundle given as an F -cocycle by $i : C_0 \longrightarrow \text{pt} \xrightarrow{\text{pt}_k} \text{Vect}$. An element in the monoid of bibranes for this trivial line bundle on the span given by the source and target map

$$\begin{array}{ccc} & C_1 & \\ s \swarrow & & \searrow t \\ C_0 & & C_0 \end{array}$$

is a transformation of the form $\begin{array}{ccc} & C_1 & \\ s \swarrow & & \searrow t \\ C_0 & \xrightarrow{V} & C_0 \\ i \swarrow & & \searrow i \\ & \text{Vect} & \end{array}$. In terms of its components this is canonically identified with

a function $V : C_1 \rightarrow k$ from the space (set) of morphisms to the ground field and every such function gives such a transformation. This identifies the C -bibranes with functions on C_1 .

Given two such bibranes V, W , their product as bibranes is, according to definition 5.8, the push-forward along the composition map on C of the function on the space (set) of composable morphisms

$$C_1 \times_{t,s} C_1 \rightarrow k$$

$$(\xrightarrow{f} \xrightarrow{g}) \mapsto V(f) \cdot W(g).$$

This push-forward is indeed the product operation on the category algebra.


6.1.3 Monoidal categories of graded vector spaces from bibrane monoids

The straightforward categorification of the discussion of group algebras in section 6.1.2 leads to bibrane monoids equivalent to monoidal categories of graded vector spaces.

Let now $F := 2\text{Vect}$ be a model for the 2-category of 2-vector spaces. For our purposes and for simplicity, it is sufficient to take $F := \mathbf{B}\text{Vect} \hookrightarrow 2\text{Vect}$, the 2-category with a single object, vector spaces as morphisms with composition being the tensor product, and linear maps as 2-morphisms. This can be regarded as the full sub-2-category of 2Vect on 1-dimensional 2-vector spaces. And we can assume $\mathbf{B}\text{Vect}$ to be strictified.

Notice from definition ?? that the ground ω -monoid in this case is the monoidal category $K = \text{Vect}$.

Then bibranes over G for the trivial 2-vector bundle on the point, i.e. transformations of the form



canonically form the category Vect^G of G -graded vector spaces. The fusion of such bibranes

reproduces the standard monoidal structure on Vect^G .

6.1.4 Twisted vector bundles

The ordinary notion of a brane in string theory is: for an abelian gerbe \mathcal{G} on target space X a map $\iota : Q \rightarrow X$ and a $\text{PU}(n)$ -principal bundle on Q whose lifting gerbe for a lift to a $U(n)$ -bundle is the pulled back gerbe $\iota^*\mathcal{G}$. Equivalently: a twisted $U(n)$ -bundle on Q whose twist is $\iota^*\mathcal{G}$. Equivalently: a gerbe module for $\iota^*\mathcal{G}$.


We show how this is reproduced as a special case of the general notion of branes from definition 5.5, see also [37].

The bundle gerbe on X is given by a cocycle $g : X \multimap \mathbf{B}BU(1)$. The coefficient group has a canonical representation $\rho : \mathbf{B}^2U(1) \rightarrow F := \mathbf{B}\text{Vect} \hookrightarrow 2\text{Vect}$ on 2-vector spaces (as in section 6.1.3) given by

$$\rho : \begin{array}{ccc} & \text{Id} & \\ \circlearrowleft & & \circlearrowright \\ \bullet & \in \mathbf{B}U(1) & \bullet \\ \circlearrowright & & \circlearrowleft \\ & \text{Id} & \end{array} \mapsto \begin{array}{ccc} & \mathbb{C} & \\ \circlearrowleft & & \circlearrowright \\ \bullet & \downarrow \cdot c & \bullet \\ \circlearrowright & & \circlearrowleft \\ & \mathbb{C} & \end{array} .$$

See also [37, 31].

By inspection one indeed finds that branes in the sense of diagrams



identified with twisted vector bundles on Q with twist given by the ι^*g : the naturality condition satisfied by the components of V is

$$\begin{array}{ccc} & \mathbb{C} & \\ & \swarrow \cdot c \quad \searrow \cdot c & \\ \mathbb{C} & \xrightarrow{\cdot c} & \mathbb{C} \\ \downarrow (\pi_1^* E)_y & \searrow \pi_{13}^* g_{\text{tw}}(y) & \downarrow \pi_3^* E_y \\ \mathbb{C} & \xrightarrow{\cdot c} & \mathbb{C} \end{array} \quad = \quad \begin{array}{ccc} & \mathbb{C} & \\ & \swarrow \cdot c \quad \searrow \cdot c & \\ \mathbb{C} & \xrightarrow{\pi_{12}^* g_{\text{tw}}(y)} & \mathbb{C} \\ \downarrow \pi_1^* E_y & \searrow \pi_{23}^* g_{\text{tw}}(y) & \downarrow (\pi_3^* E)_y \\ \mathbb{C} & \xrightarrow{\cdot c} & \mathbb{C} \\ \downarrow \pi_1^* E_y & \searrow \cdot g(y) & \downarrow \pi_3^* E_y \\ \mathbb{C} & \xrightarrow{\cdot c} & \mathbb{C} \end{array} ,$$

for all $y \in Y \times_X Y \times_X Y \times_X Y$ in the triple fiber product of a local-sections admitting map $\pi : Y \rightarrow X$ whose simplicial nerve Y^\bullet , regarded as an ω -category, provides the cover for the ω -anafunctor $X \xleftarrow{\simeq} Y^\bullet \xrightarrow{g} \mathbf{B}^2U(1)$ representing the gerbe. See [37] for details. $E \rightarrow Y$ is the vector bundle on the cover encoded by the transformation V . The above naturality diagram says that its transition function g_{tw} satisfies the usual cocycle

condition for a bundle only up to the twist given by the gerbe g : if $Y \rightarrow X$ is a cover by open subsets $Y = \sqcup_i U_i$, then the above diagram is equivalent to the familiar equation

$$(g_{\text{tw}})_{ij}(g_{\text{tw}})_{jk} = (g_{\text{tw}})_{ik} \cdot g_{ijk}.$$

In this functorial cocyclic form twisted bundles on branes were described in [30, 37].

6.1.5 2-Hilbert spaces

Let B be the category internal to spans in ω -categories given by all product spans in **Sets** with composition morphisms the canonical morphisms. Let $F = \mathbf{B}\text{Vect}$ as before. Then the 2-category of (B, F) -bibranses is the 2-category of 2-Hilbert spaces as in [6].

6.1.6 The path integral

We unwrap the notion of propagation in a σ -model form section 5.3 for the case that the background field is an ordinary vector bundle (with connection), i.e. for the case $F = \text{Vect}$. This can be regarded in terms of the quantization of the charged 1-particle as well as, after transgression, as the top-dimensional propagation in higher dimensional theories. We shall re-encounter this example in the discussion of Dijkgraaf-Witten theory in section 6.2.

Let for the present example the parameter space Cob consist just of a single edge

$$\text{Cob} = \left\{ \begin{array}{c} \Sigma := \{a \rightarrow b\} \\ \nearrow \quad \quad \quad \nwarrow \\ \Sigma_{\text{in}} := \{a\} \quad \quad \quad \Sigma_{\text{in}} := \{b\} \end{array} \right\}.$$

Recall from section 6.1.1 that for $F = \text{Vect}$ and $\rho : \mathbf{B}G \rightarrow \text{Vect}$ a linear representation, we have $\rho^* \mathbf{E}_{\text{pt}} F = V//G$ is the action groupoid of G acting on the representation space V .

Write $\nabla := \rho \circ g$ for the background field. It follows that the ω -bundle over X is given by the groupoid $\nabla^* \mathbf{E}_{\text{pt}} F$ with morphisms

$$(\nabla^* \mathbf{E}_{\text{pt}} F)_1 = \left\{ (x_1, v_1) \xrightarrow{\gamma} (x_2, v_2) \mid (x \xrightarrow{\gamma} y) \in X, v_1, v_2 \in V, v_2 = \rho(g(\gamma)) \right\}$$

with the obvious composition operation.

So a *state* in \mathcal{H}_{Σ_a} , a groupoid $v : \Psi \rightarrow \nabla^* V//G$ over $\nabla^* V//G$, is over each point $x \in X$ a groupoid over V . By the yoga of groupoid cardinality [1, 2] we can hence identify a state $v : \Psi \rightarrow \nabla^* V//G$ with a V -valued function on $\text{Obj}(X)$.

The objects of the transgressed background bundle $(\nabla^\Sigma)^* \mathbf{E}_{\text{pt}}(F^\Sigma)$ are the morphisms of $\nabla^* \mathbf{E}_{\text{pt}} F$.

The pull-push propagation map

$$\int_{\text{hom}(\Sigma, X)} \exp\left(\int \nabla\right) : \mathcal{H}_a \rightarrow \mathcal{H}_b$$

reproduces the path integral in this setup as described in [32].

6.2 Dijkgraaf-Witten model: target space $\mathbf{B}G_1$

Dijkgraaf-Witten theory [16] is the σ -model which in our terms is specified by the data

- target space $X = \mathbf{B}G$, the one-object groupoid corresponding to an ordinary 1-group G ;
- background field $\alpha : \mathbf{B}G \rightarrow \mathbf{B}^3U(1)$, a group 3-cocycle on G .

6.2.1 The 3-cocycle


Indeed, we can understand group cocycles precisely as ω -anafunctors $\mathbf{BG} \leftarrow^{\cong} Y \xrightarrow{\alpha} \mathbf{B}^n U(1)$. This is described in [8]. Here it is convenient to take Y to be essentially the free ω -category on the nerve of \mathbf{BG} , i.e. $Y := F(N(\mathbf{BG}))$, but with a few formal inverses thrown in to ensure that we have an acyclic fibration to \mathbf{BG} :

the 1-morphisms of Y are given by finite sequences of elements of G , its 2-morphisms are freely generated as pasting diagrams from 2-morphisms of the form $\left\{ \begin{array}{c} \bullet \\ \begin{array}{ccc} g \nearrow & & \searrow h \\ & \Downarrow & \\ \bullet & \xrightarrow{hg} & \bullet \end{array} \end{array} \right\}$ together with their formal

inverses. Its 3-morphisms are freely generated as pasting diagrams from 3-morphisms of the form

$$\left\{ \begin{array}{ccc} \begin{array}{ccc} \bullet & \xrightarrow{h} & \bullet \\ \uparrow & \Downarrow & \uparrow \\ g & & hg \\ \bullet & \xrightarrow{khg} & \bullet \end{array} & \xrightarrow{(g,h,k)} & \begin{array}{ccc} \bullet & \xrightarrow{h} & \bullet \\ \uparrow & \Downarrow & \uparrow \\ g & & kh \\ \bullet & \xrightarrow{khg} & \bullet \end{array} \end{array} \right\}$$

together with their formal inverses. Its 4-morphisms are freely generated from pasting diagrams of 4-morphisms of the form



together with their formal inverses.

The ω -functor $\alpha : Y \rightarrow \mathbf{B}^3U(1)$ has to send the generating 3-morphisms (g, h, k) to a 3-morphism in $\mathbf{B}^3U(1)$, which is an element $\alpha(g, h, k) \in U(1)$. In addition, it has to map the generating 4-morphisms between pasting diagrams of these 3-morphisms to 4-morphisms in $\mathbf{B}^3U(1)$. Since there are only identity 4-morphisms in $\mathbf{B}^3U(1)$ and since composition of 3-morphisms in $\mathbf{B}^3U(1)$ is just the product in $U(1)$, this says that α has to satisfy the equations

$$\forall g, h, k, l \in G : \alpha(g, h, k)\alpha(g, kh, l)\alpha(h, k, l) = \alpha(hg, k, l)\alpha(g, h, lk)$$

in $U(1)$. This identifies the ω -functor α with a group 3-cocycle on G . Conversely, every group 3-cocycle gives rise to such an ω -functor and one can check that coboundaries of group cocycles correspond precisely to transformations between these ω -functors. Notice that α uniquely extends to the additional formal inverses of cells in Y which ensure that $Y \xrightarrow{\cong} \mathbf{B}G$ is indeed an acyclic fibration. For instance the 3-cell

$$\left(\begin{array}{ccc} \bullet & \xrightarrow{h} & \bullet \\ \uparrow g & \swarrow \nearrow hg & \downarrow k \\ \bullet & \xrightarrow{khg} & \bullet \end{array} \xrightarrow{(g,h,k)'} \begin{array}{ccc} \bullet & \xrightarrow{h} & \bullet \\ \uparrow g & \swarrow \nearrow kh & \downarrow k \\ \bullet & \xrightarrow{khg} & \bullet \end{array} \right)$$

has to go to $\alpha(g, h, k)^{-1}$.

6.2.2 Chern-Simons theory

In this article we do not want to get into details of the discussion of ω -categories internal to smooth spaces, but in light of the previous section 6.2 it should be noted that in terms of nonabelian cocycles the appearance of Chern-Simons theory is formally essentially the same as that of Dijkgraaf-Witten theory:

if we take BG to be a smooth model of the classifying space of G -principal bundles, then a smooth cocycle $BG \dashrightarrow \mathbf{B}^3U(1)$, i.e. an ω -anafunctor internal to (suitably generalized) smooth spaces is precisely the cocycle for a 2-gerbe, i.e. a line 3-bundle. In nonabelian cohomology, the difference between group cocycles and higher bundles is no longer a conceptual difference, but just a matter of choice of target “space” ω -groupoid.

6.2.3 Transgression of DW theory to loop space

Proposition 6.1 *The background field α of Dijkgraaf-Witten theory transgressed according to definition 5.4 to the mapping space of parameter space $\Sigma := \mathbf{B}\mathbb{Z}$ – a combinatorial model of the circle –*

$$\tau_{\mathbf{B}\mathbb{Z}}\alpha := \text{hom}(\mathbf{B}\mathbb{Z}, \alpha)_1 : \Lambda G \rightarrow \mathbf{B}^2U(1)$$


is the groupoid 2-cocycle known as the twist of the Drinfeld double, as recalled for instance on the first page of [42]:

$$(\tau_{\mathbf{B}\mathbb{Z}}\alpha) : (x \xrightarrow{g} gxg^{-1} \xrightarrow{h} (hg)x(hg)^{-1}) \mapsto \frac{\alpha(x, g, h) \alpha(g, h, (hg)x(hg)^{-1})}{\alpha(h, gxg^{-1}, g)}.$$


Proof. According to definition A.13 the transgressed functor is obtained on 2-cells as the composition of ω -anafunctors $\mathbf{B}\mathbb{Z} \xrightarrow{(x,g)} \mathbf{B}G \xrightarrow{\alpha} \mathbf{B}^3U(1)$, given by

$$\begin{array}{ccc} (x, g, h)^*Y & \longrightarrow & Y \xrightarrow{\alpha} \mathbf{B}^3U(1) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathbf{B}\mathbb{Z} \otimes O([2]) & \xrightarrow{(x,g,h)} & \mathbf{B}G \end{array}$$


where (x, g, h) denotes a 2-cell in ΛG




which comes from a prism



in \mathbf{BG} . The 2-cocycle $\tau_{\mathbf{BZ}}\alpha$ evidently sends this to the evaluation of α on a 3-morphism in the cover Y filling this prism. One representation of such a 3-morphism, going from the back and rear to the top and front of this prism, is



Here the first step follows by 2-dimensional whiskering of the standard 3-morphism:



This manifestly yields the cocycle as claimed. \square

6.2.4 The Drinfeld double modular tensor category from DW bibranes

Let again $\rho : \mathbf{B}^2U(1) \rightarrow 2\mathbf{Vect}$ be the representation of $\mathbf{BU}(1)$ from section 6.1.3 and let $\tau_{\mathbf{BZ}\alpha} : \Lambda G \rightarrow \mathbf{B}^2U(1)$ be the 2-cocycle obtained in section 6.2.3 from transgression of a Dijkgraaf-Witten line 3-bundle on \mathbf{BG} and consider the ρ -associated 2-vector bundle $\rho \circ \tau_{\mathbf{BZ}\alpha}$ corresponding to that. Its sections according to definition 3.25 form a category $\Gamma(\tau_{\mathbf{BZ}\alpha})$.


Corollary 6.2 *The category $\Gamma(\tau_{\mathbf{BZ}\alpha})$ is canonically isomorphic to the representation category of the α -twisted Drinfeld double of G .*

Proof. Follows by inspection of our definition of sections applied to this case and using the relation established in 6.2.3 between nonabelian cocycles and the ordinary appearance of the Drinfeld double in the literature. \square

In the case that α is trivial, the representation category of the twisted Drinfeld double is well known to be a modular tensor category. We now show how the fusion tensor product on this category is reproduced from a monoid of bibranes on ΛG .


Consider any 2-group $\mathbf{BG}_2 := (G \times H \xrightarrow[\text{(Id}, \delta)}{p_1} G \longrightarrow \text{pt})$.

Pullback to the single object of \mathbf{BEZ} yields a canonical morphism from the *disk-space* $DG_2 := \text{hom}(\mathbf{BEZ}, \mathbf{BG}_2)$ to \mathbf{BG} , $p : DG_2 \rightarrow \mathbf{BG}$ which inherits from the 2-group the structure of a category internal to groupoids in that on the span



there is induced the structure of a monad from the horizontal composition in G_2 . Notice that DG_2 is very similar to but in general slightly different from the action groupoid $H//G$ obtained from the canonical action of G on H in a 2-group. Both coincide in the special case that $G_2 = \mathbf{EG}$, so that $H = G$. In this case the morphism p exhibits DG_2 as the action groupoid (as in section 6.1.1) of G acting on itself by the adjoint action.

For $\mathbf{BG} \rightarrow 2\mathbf{Vect}$ the trivial gerbe, the transformations



are representations of DG_2 on

vector spaces. In the case that $H = G$ and the boundary map is the identity we have $DG_2 = \Lambda G$, so that, by the above, bibranes on DG_2 become representations of ΛG .

One checks in this case that the fusion product of bibranes using the internal category structure on DG_2 according to 5.8 does reproduce the familiar fusion tensor product on representations of ΛG , hence of the Drinfeld double.

6.2.5 The DW path integral

Let S_{in} and S_{out} be two oriented surfaces and let V be an oriented 3-manifold with boundary $\partial V = S_{\text{in}} \sqcup \bar{S}_{\text{out}}$. Forming fundamental groupoids yields a co-span

$$\begin{array}{ccc} & \Sigma := \Pi_1(V) & \\ \nearrow & & \nwarrow \\ \Sigma_{\text{in}} := \Pi_1(S_{\text{in}}) & & \Sigma_{\text{out}} := \Pi_1(S_{\text{out}}) \end{array}$$

Notice that the space of fields on V in DW theory $\text{hom}(\Sigma, \mathbf{BG})$ is equivalent to the groupoid of G -principal bundles on V . This implies that pull-push quantum propagation in the sense of section 5.3 reproduces the right DW path integral.

6.3 Yetter-Martins-Porter model: target space \mathbf{BG}_2

The Yetter-Martins-Porter model is a σ -model with target space $X = \mathbf{BG}$ for G a 2-group.

Here, too, our quantization reproduces the right combinatorial path integral factor [?].

A ω -Categories and their Homotopy Theory

An ∞ -category is a combinatorial model for higher directed homotopies, a combinatorial model for a *directed space*. The fact that it is *directed* means that not all cells in this space are necessarily *reversible*. If they are, the ∞ -category is an ∞ -groupoid, a combinatorial model for an ordinary space.

There are various definitions of ∞ -categories and ∞ -groupoids [21]. Most of them model ∞ -categories as conglomerates of n -dimensional cells of certain shape, for all $n \in \mathbb{N}$, equipped with certain structure and certain properties.

Conglomerates of cells. A “conglomerate of n -dimensional cells of certain shape” technically means a presheaf on a category of basic cells.

Simplicial sets and $(\infty, 1)$ -categories. The most familiar example is the simplicial category Δ whose objects are the standard cellular simplices and presheaves on which are simplicial sets. A popular model for ∞ -groupoids are simplicial sets with the *Kan property*: *Kan complexes*. The Kan property can be interpreted as ensuring that for all adjacent simplices in the Kan simplicial set there exists a composite simplex and that for all simplices there exists a reverse simplex. Replacing the Kan property on simplicial sets by a slightly weaker property called the *weak Kan property* generalizes Kan complexes to a model of ∞ -categories called *weak Kan complexes* or *quasicategories* or $(\infty, 1)$ -*categories*: the weak Kan condition ensures just that for all n -simplices for $n \geq 2$ there exists a reverse simplex. A further weakening of the Kan condition such as to ensure only the existence of composites without any restriction on reversibility leads to a definition of weak ∞ -categories based on simplicial sets with extra properties proposed by Ross Street []. This is very general but also somewhat unwieldy.

Two other basic shapes of relevance besides simplices are globes and cubes.

Globular sets and ω -categories.

Cubical sets and n -fold categories.

∞ -Categories in terms of 1-categories. A general strategy to handle ∞ -categories in practice is to regard them as categories (i.e. 1-categories) with extra bells and whistles. This notably involves the tools of *enriched category theory* and of *model category theory*.

Enriched categories. The definition of a category *enriched* over a monoidal category \mathcal{V} [20] is like that of an ordinary category, but with the requirement that there is a *set* of morphisms between any two objects replaced by the requirement that there is an *object of C* for any two objects. If the enriching category \mathcal{V} is a category of higher structures, such as simplicial sets, \mathcal{V} -enriched categories are models for ∞ -categories. In practice the advantage of conceiving ∞ -categories as suitably enriched categories is that enriched category theory is a well-developed subject with a supply of powerful general tools.

Model categories. From the modern perspective, a model category (Quillen model category), is the 1-categorical truncation of an $(\infty, 1)$ -category, remembering which of the 1-morphisms retained used to be like isomorphisms, monomorphisms and epimorphisms up to higher coherent cells, in the original $(\infty, 1)$ -category: in a model category these special 1-morphisms are, respectively, called *weak equivalences*, *cofibrations* and *fibrations* and satisfy a couple of properties.

One says that model categories are *presentations* of $(\infty, 1)$ -categories in that they provide a convenient re-packaging of the information contained in an $(\infty, 1)$ -category in purely 1-categorical terms. In practical computation the model category structure on a 1-category is in particular used to generalize morphisms between given objects to morphisms between suitable weakly equivalent *replacements* of these objects.

Our approach. The ∞ -vector bundles which we want to describe are given by cocycles with values in ∞ -categories (of models for ∞ -vector spaces) which are not ∞ -groupoids and are not $(\infty, 1)$ -categories in that in general they have non-reversible cells in all degrees.

Among the simplicial models for ∞ -categories this would force one to use models such as Street’s weak ∞ -categories. This model, however, we find unwieldy for our applications.

Among the remaining choices of models for ∞ -categories for our developments in sections 3 and 5 we choose one which combines the “folk” model category structure [14] on $\omega\mathbf{Categories}$ with the enrichment of $\omega\mathbf{Categories}$ over itself [11]. For most considerations in section 5 and 6 this means effectively that we work in the 1-category $\omega\mathbf{Categories}$ while making use of the internal hom-functor and using the freedom to replace ω -categories by weakly equivalent replacements.

For handling $\omega\mathbf{Categories}$ the different shapes – globes, simplices, cubes – are useful for different purposes. Globular sets have the simplest boundary structure, simplicial sets provide powerful computational tools, cubical sets provide the important monoidal structure. In the following all three models of shapes are combined: following [8, 11] we conceive ω -categories as globular sets for general purposes and make use of their incarnation as cubical sets for describing their biclosed monoidal structure. Moreover, following [38, 8] we use cosimplicial ω -categories such as the orientals to pass between $\omega\mathbf{Categories}$ and $\mathbf{SimplicialSets}$, mostly for the purpose of constructing weakly equivalent replacements of ω -categories.

A.1 Shapes for ∞ -cells

Three types of basic shapes are used frequently: globes, simplices and cubes. These are modeled, respectively, by the globular category G , the simplicial category Δ and the cubical category C . These three categories have as objects the integers, $n \in \mathbb{N}$, thought of as the standard cellular n -globe G^n , the standard cellular n -simplex Δ^n and the standard cellular n -cube C^n , respectively. Morphisms are all maps between these standard cellular shapes which respect the cellular structure.

Definition A.1 (globular category) *The globular category G is the category whose objects are the integers \mathbb{N} and whose morphisms are generated from morphisms*

$$\sigma_n, \tau_n : [n] \rightarrow [n + 1]$$

subject to the relations

$$\begin{array}{ccc} [n] & \xrightarrow{\sigma_n} & [n+1] \\ \sigma_n \downarrow & & \downarrow \sigma_n \\ [n+1] & \xrightarrow{\tau_n} & [n+2] \end{array} \quad , \quad \begin{array}{ccc} [n] & \xrightarrow{\tau_n} & [n+1] \\ \tau_n \downarrow & & \downarrow \sigma_n \\ [n+1] & \xrightarrow{\tau_n} & [n+2] \end{array}$$

for all $n \in \mathbb{N}$.

Definition A.2 (simplicial category) *The simplicial category Δ is the full subcategory of Categories on categories which are freely generated from connected linear graphs. Equivalently, Δ is the category with totally ordered finite sets as objects and order-preserving maps as morphisms.*

Definition A.3 (cubical category) *The cubical category C is defined ... section 2 of [11]*

Definition A.4 (monoidal structure on the cubical category) *section 2 of [11]*

A.2 ω -Categories

Recall the following standard facts:

- The category **Sets** is symmetric monoidal with respect to the standard cartesian product.
- For \mathcal{V} a symmetric monoidal category, the category \mathcal{V} -Cat of \mathcal{V} -enriched categories is naturally itself symmetric monoidal.

Definition A.5 (strict globular n -category, [13]) *The category of 0-categories is $0\text{Categories} := \mathbf{Sets}$. For $n \in \mathbb{N}$, $n \geq 1$ the category of (“strict, globular”) n -categories is defined inductively as the category*

$$n\text{Categories} := (n-1)\text{Categories} - \text{Cat}$$

of categories enriched over $(n-1)\text{Categories}$.

One notices that for all $n \in \mathbb{N}$ there is a canonical inclusion $n\text{Categories} \hookrightarrow (n+1)\text{Categories}$.

Definition A.6 (ω -category, [39]) *The category of ω -categories is the direct limit over this chain of inclusions*

$$\omega\text{Categories} := \lim_{\rightarrow_{n \in \mathbb{N}}} n\text{Categories}.$$

Unwrapping this definition shows that ω -categories are globular sets equipped with compatible structures of a strict 2-category on all sub-globular sets of length two:

Definition A.7 (globular set) *A globular set S is a presheaf on the globular category G , i.e. a functor $S : G^{\text{op}} \rightarrow \mathbf{Sets}$.*

We write $S([n] \xrightarrow{\sigma_n, \tau_n} [n+1]) := S_{n+1} \xrightarrow{s_n, t_n} S_n$ and call S_n the set of n -globes, s_n the n -source map and t_n the n -target map of S . The identities $s_n \circ s_{n+1} = s_n \circ t_{n+1}$ and $t_n \circ s_{n+1} = t_n \circ t_{n+1}$, called the globular identities, ensure that for all $n, k \in \mathbb{N}$ there are unique maps $S_{n+k} \xrightarrow{s, t} S_n$ themselves satisfying analogous globular identities.

Proposition A.8 (ω -category, [38]) *An ω -category C is a globular set $C : G^{\text{op}} \rightarrow \mathbf{Sets}$ equipped for all $n, k \in \mathbb{N}$ the structure of a category extending $C_{n+k} \xrightarrow[s]{t} C_n$ such that this makes for all $n, k, l \in \mathbb{N}$*

$$C_{n+k+l} \xrightarrow[s]{t} C_{n+k} \xrightarrow[s]{t} C_n \text{ into a strict 2-category.}$$

The elements in C_k are called k -morphisms. The composition in $C_{n+k} \xrightarrow[t]{s} C_n$ is called composition of $n+k$ -morphisms along n -morphisms. A morphism between ω -categories, called an ω -functor, is a morphism of the underlying globular sets respecting all the additional structure.

Definition A.9 (standard globular globes) *The globular set G_n represented by $n \in \mathbb{N}$, $G_n := \text{Hom}_G(-, [n])$ is the standard globular globe. There is a unique structure of an ω -category on G_n . This yields co-globular ω -category G_\bullet , i.e. a functor $G^\bullet : G \rightarrow \omega\text{Categories}$.*

We also write

- $\emptyset :=: G^{-1} :=: \mathcal{I}^{-1}$ for the ω -category on the empty globular set (the initial object in $\omega\text{Categories}$);
- $\text{pt} :=: I :=: \mathcal{I}^0 = G^0 = \{\bullet\}$ for the ω -category with a single object and no nontrivial morphisms (the terminal object in $\omega\text{Categories}$ and the tensor unit with respect to the Crans-Gray tensor product \otimes described below);
- $\mathcal{I} :=: \mathcal{I}^1 :=: G^1 = \{a \longrightarrow b\}$ for the ω -category with two objects and a single nontrivial morphism connecting them.

The first few n -globes can be depicted as follows:

$$G^0 = \{d_0\} \xrightarrow[\tau_0: d_0 \mapsto d_0^+]{\sigma_0: d_0 \mapsto d_0^-} G^1 = \{d_0^- \xrightarrow{d_1} d_0^+\} \xrightarrow[\tau_1: d_1 \mapsto d_1^+]{\sigma_1: d_1 \mapsto d_1^-} G^2 = \{d_0^- \begin{array}{c} \xrightarrow{d_1^-} \\ \Downarrow d_2 \\ \xrightarrow{d_1^+} \end{array} d_0^+\} \xrightarrow[\tau_2: d_2 \mapsto d_2^+]{\sigma_2: d_2 \mapsto d_2^-} G^3 = \{d_1^- \begin{array}{c} \xrightarrow{d_0^-} \\ \xrightarrow{d_2^-} \\ \Downarrow d_3 \\ \xrightarrow{d_2^+} \\ \xrightarrow{d_0^+} \end{array} d_1^+\} .$$

A.3 ω -Groupoids

... ω -groupoids and crossed complexes...

A.4 Cosimplicial ω -categories

We can translate back and forth between simplicial sets and ω -categories by means of a fixed cosimplicial ω -category, i.e. a functor $O : \Delta \rightarrow \omega\text{Categories}$ from the simplicial category Δ : from any such we obtain an ω -nerve functor $N : \omega\text{Categories} \rightarrow \text{SimplicialSets}$ by

$$N(C) : \Delta^{\text{op}} \xrightarrow{O^{\text{op}}} \omega\text{Categories}^{\text{op}} \xrightarrow{\text{Hom}(-, C)} \text{Sets}$$

and its left adjoint $F : \text{SimplicialSets} \rightarrow \omega\text{Categories}$ given by the coend formula

$$F(S^\bullet) := \int^{[n] \in \Delta} S^n \cdot O([n]).$$

Ross Street defined such a cosimplicial ω -category called the orientals [38], for which $O([n])$ is the ω -category free on a single n -morphism of the shape of an n -simplex. To obtain more inverses, we can alternatively use the unorientals, for which $O([n])$ is the ω -category with n -objects, with 1-morphisms finite sequences of these objects, 2-morphisms finite sequences of such finite sequences, and so on.

A.5 Monoidal biclosed structure on ω Categories


The category $\omega\text{Categories}$ is equipped with the Crans-Gray tensor product [11], which is the extension to ω -categories of the tensor product on cubical sets which in turn is induced via Day convolution from the addition of natural numbers. This means that the Crans-Gray tensor product is dimension raising in a way analogous to the cartesian product on topological spaces:

for instance the tensor product of the interval ω -category $I = \{ a \longrightarrow b \}$ with itself is the ω -category free on a single directed square


$$I \otimes I = \left\{ \begin{array}{ccc} (a, a) & \longrightarrow & (a, b) \\ \downarrow & \swarrow & \downarrow \\ (b, a) & \longrightarrow & (b, b) \end{array} \right\}.$$

Moreover, $\omega\text{Categories}$ is biclosed with respect to this monoidal structure.

Definition A.10 (internal hom) For ω -categories C and D the ω -category $[C, D]$ is given by the globular set $\text{Hom}(G_{[-]} \otimes C, D) : G^{\text{op}} \rightarrow \text{Sets}$ on which the composition of k -morphisms along an l -morphism is defined as the image of the diagram which glues two standard k -globes along a common l -globe



under $\text{Hom}(G_{[-]} \otimes C, D)$:



Remarks. Notice that everything in this definition works by abstract nonsense – for instance that the contravariant Hom takes colimits to limits – except the existence of the maps $c_{k,l}$, which encodes genuine information about pasting of standard globes [10]. For instance $G_2 \sqcup_{G_0} G_2 = \left\{ \begin{array}{ccc} a & \begin{array}{c} \curvearrowright \\ \Downarrow \\ \curvearrowleft \end{array} & b \\ & & \end{array} \right\}$

while $G_2 \sqcup_{G_1} G_2 = \left\{ \begin{array}{ccc} a & \begin{array}{c} \curvearrowright \\ \Downarrow \\ \longrightarrow \\ \Downarrow \\ \curvearrowleft \end{array} & b \\ & & \end{array} \right\}$, where the right sides denote the free ω -categories on the indicated

pasting diagram [10].

Proposition A.11 *For every $C \in \omega\text{Categories}$ this extends to a functor $[C, -] : \omega\text{Categories} \rightarrow \omega\text{Categories}$ which is right adjoint to $- \otimes C : \omega\text{Categories} \rightarrow \omega\text{Categories}$.*

Of particular interest to us are the internal hom- ω -categories of the form $A^{\mathcal{I}} := [\mathcal{I}, A]$ which satisfy

$$\text{Hom}(I \otimes X, A) \simeq \text{Hom}(X, A^{\mathcal{I}}), \text{ where the set in question here is the set of } \underline{\text{lax transformations}} \quad X \begin{array}{c} \xrightarrow{f} \\ \Downarrow \eta \\ \xrightarrow{g} \end{array} A \Leftrightarrow$$

$$X \xrightarrow{\eta} A^{\mathcal{I}} \xrightarrow{d_0 \times d_1} A \times A \text{ or directed right homotopies between } \omega\text{-functors from } X \text{ to } A.$$

A.6 Model structure on $\omega\text{Categories}$

That the 1-category $\omega\text{Categories}$ is really an ∞ -structure itself is remembered by a *model category structure* carried by it, due to [14], with respect to which the acyclic fibrations or hypercovers $f : C \xrightarrow{\simeq} D$ are those ω -functors which are k -surjective for all $k \in \mathbb{N}$, meaning that the universal dashed morphism in

$$\begin{array}{ccc} C_{k+1} & \xrightarrow{f_{k+1}} & D_{k+1} \\ \downarrow \text{dashed} & \searrow & \downarrow \text{dashed} \\ (f_k \times f_k)^* C_{k+1} & \rightarrow & D_{k+1} \\ \downarrow s \times t & & \downarrow s \times t \\ C_k \times C_k & \xrightarrow{f_k \times f_k} & D_k \times D_k \end{array}$$

is epi, for all k . The weak equivalences $f : C \xrightarrow{\simeq} D$ are those ω -functors where these dashed morphisms become epi after projecting onto ω -equivalence classes of $(k+1)$ -morphisms.

Using this we define an ω -anafunctor from an ω -category X to an ω -category A to be a span

$$(g : X \dashrightarrow A) := \begin{array}{c} \hat{X} \xrightarrow{g} A \\ \downarrow \simeq \\ X \end{array}$$

whose left leg is a hypercover. (This terminology follows [22, 5].) One finds [7] that in the context of $\omega\text{Groupoids}$ such ω -anafunctors represent morphisms in the homotopy category $[g] \in \text{Ho}(X, A)$ which allows us to regard g as a cocycle in nonabelian cohomology on the ω -groupoid X with coefficients in the ω -groupoid A . Cocycles are regarded as distinct only up to refinements of their covers. This makes their composition by pullbacks

$$(X \dashrightarrow A \dashrightarrow A') := \begin{array}{ccc} g^* \hat{A} & \longrightarrow & \hat{A} \xrightarrow{r} A' \\ \downarrow \simeq & & \downarrow \simeq \\ X & \xrightarrow{g} & A \\ \downarrow \simeq & & \\ X & & \end{array}$$

well defined (noticing that acyclic fibrations are closed under pullback) and associative.

Definition A.12 We write \mathbf{Ho} for the corresponding category of ω -anafunctors,

$$\mathbf{Ho}(C, D) := \operatorname{colim}_{\hat{C} \in \operatorname{Hypercovers}(C)} \operatorname{Hom}(\hat{C}, D).$$

(This is to be contrasted with the true homotopy category \mathbf{Ho} , which is obtained by further dividing out homotopies.)

While cocycles in nonabelian cohomology are morphisms in \mathbf{Ho} , coboundaries should be morphisms between these morphisms. Hence \mathbf{Ho} is to be thought of as enriched over $\omega\mathbf{Categories}$.

Definition A.13 Define a functor $\operatorname{hom} : \mathbf{Ho}^{\operatorname{op}} \times \mathbf{Ho} \rightarrow \omega\mathbf{Categories}$ by $\operatorname{hom}(C, D) := F(\operatorname{Hom}(C \otimes O([\bullet]), D))$.

References

- [1] J. Baez, *Higher dimensional algebra VII: Groupoidification*, [<http://math.ucr.edu/home/baez/hda7.pdf>]
- [2] J. Baez, A. Hoffnung, C. Walker, *Groupoidification made easy*, [<http://math.ucr.edu/home/baez/groupoidification.pdf>]
- [3] J. Baez, and U. Schreiber, *Higher gauge theory*, Categories in Algebra, Geometry and Mathematical Physics, 7–30, Contemp. Math., **431**, Amer. Math. Soc., Providence, RI, 2007, [[arXiv:math/0511710v2](https://arxiv.org/abs/math/0511710v2)] [[math.DG](#)].
- [4] I. Baković, *Bigroupoid 2-torsors*, PhD thesis, Munich 2008.
- [5] T. Bartels, *2-Bundles*, [[arXiv:math/0410328](https://arxiv.org/abs/math/0410328)] [[math.CT](#)].
- [6] B. Bartlett, *On unitary 2-representations of finite groups and topological quantum field theory*, PhD thesis, Sheffield (2008)
- [7] K. Brown, *Abstract Homotopy Theory and Generalized Sheaf Cohomology*, Transactions of the American Mathematical Society, Vol. 186 (1973), 419-458
- [8] R. Brown, P. Higgins and R. Sivera, *Nonabelian algebraic topology*
- [9] R. Brown and M. Golasinski, *A model structure on the homotopy theory of crossed complexes*, Cahier Topologie Géom. Différentielle Catég. 30 (1) (1989) 61-82
- [10] S. Crans, *Pasting presentations for ω -categories*
- [11] S. Crans, *Pasting schemes for the monoidal biclosed structure on ω -Cat*
- [12] W. Dwyer, P. Hirschhorn, D. Kan, J. Smith, *Homotopy Limit Functors on Model Categories and Homotopical Categories*, volume 113 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2004
- [13] S. Eilenberg and G. M. Kelly, *Closed categories*, Proc. Conf. Categorical Algebra at La Jolla, 1965 (Springer Verlag, Berlin 1966) 421-562
- [14] Y. Lafont, F. Metayer, and K. Worytkiewicz, *A folk model structure on omega-cat*, [[arXiv:0712.0617](https://arxiv.org/abs/0712.0617)] [[math.CT](#)].
- [15] D. Freed, *Higher Algebraic Structures and Quantization*, Commun.Math.Phys. 159 (1994) 343-398 [[arXiv:hep-th/9212115](https://arxiv.org/abs/hep-th/9212115)]
- [16] D. Freed, *Chern-Simons Theory with Finite Gauge Group*, Commun.Math.Phys. 156 (1993) 435-472 [[arXiv:hep-th/9111004](https://arxiv.org/abs/hep-th/9111004)]

- [17] J. Fuchs, C. Schweigert, K. Waldorf, *Branes: Target Space Geometry for World Sheet topological Defects* J.Geom.Phys.58:576-598,2008 [[arXiv:hep-th/0703145](#)]
- [18] Mark Hovey, *Model Categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, 1999
- [19] J. Jardine, *Stacks and the homotopy theory of simplicial sheaves*, Homology, homotopy and applications, vol 3(2), 2001, pp. 361-384
- [20] G. M. Kelly, *Basic concepts of enriched category theory*, Reprints in Theory and Applications of Categories, No. 10, 2005
- [21] T. Leinster, *Higher Operads, Higher Categories*, London Mathematical Society Lecture Note Series, **298**, Cambridge University Press, Cambridge, 2004, [[arXiv:math/0305049](#)] [[math.CT](#)].
- [22] M. Makkai, *Avoiding the axiom of choice in general category theory*, J. Pure Appl. Algebra **108** (1996), no. 2, 109-173, [<http://www.math.mcgill.ca/makkai/anafun>].
- [23] J. Martins, R.Picken, *A Cubical Set Approach to 2-Bundles with Connection and Wilson Surfaces*, [[arXiv:0808.3964](#)]
- [24] J. Morton, *Categorified algebra and quantum mechanics*, Theory and Applications of Categories, Vol. 16, 2006, No. 29, pp 785-854. [[arXiv:math/0601458](#)]
- [25] D. Roberts and U. Schreiber, *The inner automorphism 3-group of a strict 2-group*, J. Homotopy Relat. Struct. **3** (2008) no. 1, 193-244, [[arXiv:0708.1741](#)] [[math.CT](#)].
- [26] Hisham Sati, U. Schreiber, Z. Škoda, D. Stevenson, *Differential nonabelian cohomology* in preparation [<http://www.math.uni-hamburg.de/home/schreiber/nactwist.pdf>]
- [27] H. Sati, U. Schreiber and J. Stasheff, *L_∞ -connections and applications to String- and Chern-Simons n -transport*, in *Recent Developments in QFT*, eds. B. Fauser et al., Birkhäuser, Basel (2008), [[arXiv:0801.3480](#)] [[math.DG](#)].
- [28] H. Sati, U. Schreiber, and J. Stasheff, *Fivebrane structures: topology*, [[arXiv:math/0805.0564](#)] [[math.AT](#)].
- [29] H. Sati, U. Schreiber, and J. Stasheff, *Twists of and by higher bundles, such as String and Fivebrane bundles*, in preparation.
- [30] U. S. *Quantum 2-states: Sections of 2-vector bundles*, talk at *Higher categories and their applications*, Fields Institute, Jan. 2007, [<http://www.math.uni-hamburg.de/home/schreiber/atd.pdf>]
- [31] U. Schreiber, *AQFT from n -extended FQFT* [[arXiv:0806.1079](#)]
- [32] U. S., *An exercise in groupoidification: The Path integral*, blog entry, [http://golem.ph.utexas.edu/category/2008/06/an_exercise_in_groupoidificati.html]
- [33] Michael Shulman, *Homotopy limits and colimits and enriched homotopy theory*, [[arXiv:math/0610194](#)]
- [34] Jandl gerbes
- [35] U. Schreiber and K. Waldorf, *Parallel transport and functors*, [[arXiv:0705.0452](#)] [[math.DG](#)].
- [36] U. Schreiber and K. Waldorf, *Smooth functors vs. differential forms*, [[arXiv:0802.0663](#)] [[math.DG](#)].
- [37] U. Schreiber and K. Waldorf, *Connections on nonabelian gerbes and their holonomy*, [[arXiv:0808.1923](#)] [[math.DG](#)].

- [38] R. Street, *The algebra of oriented simplexes*, J. Pure Appl. Algebra **49** (1987) 283-335.
- [39] R. Street, *Categorical and combinatorial aspects of descent theory*, Appl. Categ. Structures **12** (2004), no. 5-6, 537-576, [[arXiv:math/0303175](#)] [[math.CT](#)].
- [40] B. Toën, *Stacks and non-abelian cohomology*,
- [41] B. Toën, *Higher and derived stacks: a global overview*, [[arXiv:math/0604504](#)]
- [42] S. Willerton, *The twisted Drinfeld double of a finite group via gerbes and finite groupoids* [[arXiv:math/0503266](#)]