



Introduction to Topology -- 2

This page is a detailed introduction to basic [topological homotopy theory](#). We introduce the [fundamental group](#) of [topological spaces](#) and the concept of [covering spaces](#). Then we prove the [fundamental theorem of covering spaces](#), saying that they are equivalent to [permutation representations](#) of the fundamental group. This is a simple topological version of the general principle of [Galois theory](#) and has many applications. As one example application, we use it to prove that the [fundamental group of the circle is the integers](#).

Under construction.

main page: [Introduction to Topology](#)

previous chapter: [Introduction to Topology 1 -- Point-set topology](#)

this chapter: **Introduction to Topology 2 – Basic Homotopy Theory**

For introduction to more general and abstract [homotopy theory](#) see instead at [Introduction to Homotopy Theory](#).

Basic Homotopy Theory

1. Homotopy

[Fundamental group](#)

[Groupoids](#)

2. Covering spaces

[Lifting properties](#)

[Monodromy](#)

3. Examples

[Fundamental groups](#)

[Higher homotopy groups](#)

4. References

Context

Topology

Homotopy theory

In order to handle topological spaces, to compute their properties and to distinguish them, it turns out to be useful to consider not just continuous

variation within a topological space, i.e. [continuous functions](#) between topological spaces, but also continuous deformations of [continuous functions](#) themselves. This is the concept of [homotopy](#) (def. [1.2](#) below), and its study is called [homotopy theory](#). If one regards topological spaces with [homotopy classes](#) of continuous functions between them then their nature changes, and one speaks of [homotopy types](#) (remark [1.6](#) below).

Of particular interest are homotopies between [paths](#) in a topological space. If a [loop](#) in a topological space is homotopic to the constant loop, this means that it does not “wind around a hole” in the space. Hence the set of homotopy classes of loops in a topological space, which is a [group](#) under [concatenation of paths](#), detects crucial information about the global structure of the space, and hence is called the [fundamental group](#) of the space (def. [1.16](#)).

This same information turns out to be encoded in “continuously varying sets” over a topological space, hence in “[bundles](#) of [sets](#)”, called [covering spaces](#) (def. [2.1](#) below). As one moves around a loop, then the parameterized set comes back to itself up to a [bijection](#) called the [monodromy](#) of the loop. This encodes an [action](#) or [permutation representation](#) of the fundamental group. The [fundamental theorem of covering spaces](#) (prop. [2.25](#) below) says that covering spaces are equivalently characterized by their [monodromy representation](#) of the fundamental group. This is an incarnation of the general principle of [Galois theory](#) in [topological homotopy theory](#). Sometimes this allows to compute fundamental groups from behaviour of covering spaces, for instance it allows to prove that the [fundamental group of the circle is the integers](#) (prop. [3.1](#) below).

In order to formulate and prove these statements, it turns out convenient to do away with the arbitrary choice of basepoint that is involved in the definition of fundamental groups, and instead collect *all* homotopy classes of paths into a single structure, called the [fundamental groupoid](#) of a topological space (example [1.27](#) below) an example of a generalization of groups to [groupoids](#) (discussed below). The fundamental groupoid may be regarded as an [algebraic](#) incarnation of the [homotopy type](#) presented by a topological space, *up to level 1* (the [homotopy 1-type](#)).

The algebraic reflection of the full [homotopy type](#) of a topological space involves higher dimensional analogs of the [fundamental group](#) called the *higher homotopy groups*. We close with an outlook on these [below](#).

1. Homotopy

It is clear that for $n \geq 1$ the [Euclidean space](#) \mathbb{R}^n or equivalently the [open ball](#) $B_0^\circ(1)$ in \mathbb{R}^n is *not* [homeomorphic](#) to the [point space](#) $*$ (simply because there is not even a [bijection](#) between the underlying [sets](#)). Nevertheless, intuitively the n -ball is a “continuous deformation” of the point, obtained as the radius of the n -ball tends to zero.

This intuition is made precise by observing that there is a [continuous function](#) out

of the [product topological space](#) (this example) of the open ball with the [closed interval](#)

$$\eta: [0, 1] \times B_0^\circ(1) \rightarrow B_0^\circ(1)$$

which is given by rescaling:

$$(t, x) \mapsto t \cdot x.$$

This continuously interpolates between the open ball and the point, in that for $t = 1$ it restricts to the identity, while for $t = 0$ it restricts to the map constant on the origin.

We may summarize this situation by saying that there is a [diagram](#) of [continuous functions](#) of the form

$$\begin{array}{ccc} B_0^\circ(1) \times \{0\} & \xrightarrow{\exists!} & * \\ \downarrow & & \downarrow \text{const}_0 \\ [0, 1] \times B_0^\circ(1) & \xrightarrow{(t,x) \mapsto t \cdot x} & B_0^\circ(1) \\ \uparrow & \nearrow \simeq & \\ B_0^\circ(1) \times \{1\} & & \end{array}$$

Such “continuous deformations” are called [homotopies](#):

In the following we use this terminology:

Definition 1.1. ([topological interval](#))

The [topological interval](#) is

1. the [closed interval](#) $[0, 1] \subset \mathbb{R}^1$ regarded as a [topological space](#) in the standard way, as a [subspace](#) of the [real line](#) with its [Euclidean metric topology](#),
2. equipped with the [continuous functions](#)

1. $\text{const}_0 : * \rightarrow [0, 1]$

2. $\text{const}_1 : * \rightarrow [0, 1]$

which include the [point space](#) as the two endpoints, respectively

3. equipped with the (unique) [continuous function](#)

$$[0, 1] \rightarrow *$$

to the [point space](#) (which is the [terminal object](#) in [Top](#))

regarded, in summary, as a factorization

$$\nabla_* : * \sqcup * \xrightarrow{(\text{const}_0, \text{const}_1)} [0, 1] \rightarrow *$$

of the [codiagonal](#) on the point space, namely the unique continuous function ∇_* out of the [disjoint union space](#) $* \sqcup * \simeq \text{Disc}(\{0, 1\})$ ([homeomorphic](#) to the [discrete topological space](#) on two elements).

Definition 1.2. ([homotopy](#))

Let $X, Y \in \mathbf{Top}$ be two [topological spaces](#) and let

$$f, g : X \rightarrow Y$$

be two [continuous functions](#) between them.

A [\(left\) homotopy](#) from f to g , to be denoted



$$\eta : f \Rightarrow g,$$

is a [continuous function](#)

$$\eta : X \times [0, 1] \rightarrow Y$$

out of the [product topological space](#) (this example) of X the [topological interval](#) (def. 1.1) such that this makes the following [diagram](#) in \mathbf{Top} commute:

$$\begin{array}{ccc} 0 \times X & & \\ (\text{id}, \text{const}_0) \downarrow & \searrow f & \\ X \times [0, 1] & \xrightarrow{\eta} & Y \\ (\text{id}, \text{const}_1) \uparrow & \nearrow g & \\ \{1\} \times X & & \end{array}$$



graphics grabbed from J.
Tauber [here](#)

hence such that

$$\eta(-, 0) = f \quad \text{and} \quad \eta(-, 1) = g .$$

If there is a homotopy $f \Rightarrow g$ (possibly unspecified) we say that f is *homotopic* to g , denoted

$$f \sim_h g .$$

Proposition 1.3. ([homotopy is an equivalence relation](#))

Let $X, Y \in \mathbf{Top}$ be two [topological spaces](#). Write $\text{Hom}_{\mathbf{Top}}(X, Y)$ for the [set](#) of [continuous functions](#) from X to Y .

Then the relating of being [homotopic](#) (def. 1.2) is an [equivalence relation](#) on this set. The corresponding [quotient set](#)

$$[X, Y] := \text{Hom}_{\text{Top}}(X, Y) / \sim_h$$

is called the set of [homotopy classes](#) of continuous functions.

Moreover, this equivalence relation is compatible with [composition](#) of continuous functions:

For $X, Y, Z \in \text{Top}$ three topological spaces, there is a unique function

$$[X, Y] \times [Y, Z] \rightarrow [X, Z]$$

such that the following [diagram commutes](#):

$$\begin{array}{ccc} \text{Hom}_{\text{Top}}(X, Y) \times \text{Hom}_{\text{Top}}(Y, Z) & \xrightarrow{\circ_{X, Y, Z}} & \text{Hom}_{\text{Top}}(X, Z) \\ \downarrow & & \downarrow \\ [X, Y] \times [Y, Z] & \longrightarrow & [X, Z] \end{array} .$$

Proof. To see that the relation is [reflexive](#): A homotopy $f \Rightarrow f$ from a function f to itself is given by the function which is constant on the topological interval:

$$X \times [0, 1] \xrightarrow{\text{pr}_1} X .$$

This is continuous because [projections](#) out of [product topological spaces](#) are continuous, by the [universal property](#) of the [Cartesian product](#).

To see that the relation is [symmetric](#): If $\eta: f \Rightarrow g$ is a homotopy then

$$\begin{array}{ccccc} X \times [0, 1] & \xrightarrow{\text{id}_X \times (1 - (-))} & X \times [0, 1] & \xrightarrow{\eta} & X \\ (x, t) & \mapsto & (x, 1 - t) & \mapsto & \eta(x, 1 - t) \end{array}$$

is a homotopy $g \Rightarrow f$. This is continuous because $1 - (-)$ is a [polynomial](#) function, and [polynomials are continuous](#), and because [Cartesian product](#) and [composition](#) of continuous functions is again continuous.

Finally to see that the relation is [transitive](#): If $\eta_1: f \Rightarrow g$ and $\eta_2: g \Rightarrow h$ are two composable homotopies, then consider the “ X -parameterized [path concatenation](#)”

$$\begin{array}{ccc} X \times [0, 1] & \xrightarrow{\eta_2 \circ \eta_1} & X \\ (x, t) & \mapsto & \begin{cases} \eta_1(x, 2t) & | \ t \leq 1/2 \\ \eta_2(x, 2t - 1) & | \ t \leq 1/2 \end{cases} . \end{array}$$

To see that this is continuous, observe that $\{X \times [0, 1/2] \subset X, X \times [1/2, 1] \subset X\}$ is a [cover](#) of $X \times [0, 1]$ by [closed subsets](#) (in the [product topology](#)) and because $\eta_1(-, 2(-))$ and $\eta_2(-, 2(-) - 1)$ are continuous (being composites of Cartesian products of continuous functions) and agree on the intersection $X \times \{1/2\}$. Hence the continuity follows by [this example](#).

Finally to see that homotopy respects composition: Let

$$X \xrightarrow{f_1} Y \xrightarrow[f'_2]{f_2} Z \xrightarrow{f_3} W$$

be continuous functions, and let

$$\eta : f_2 \Rightarrow f'_2$$

be a homotopy. It is sufficient to show that then there is a homotopy of the form

$$f_3 \circ f_2 \circ f_1 \Rightarrow f_3 \circ f'_2 \circ f_1 .$$

This is exhibited by the following diagram

$$\begin{array}{ccccc}
 X & \xrightarrow{f_1} & Y & & \\
 (\text{id}_X, \text{const}_0) \downarrow & & (\text{id}_Y, \text{const}_0) \downarrow & \searrow f_2 & \\
 X \times [0, 1] & \xrightarrow{f_1 \times \text{id}_{[0,1]}} & Y \times [0, 1] & \xrightarrow{\eta} & Z \xrightarrow{f_3} W . \\
 (\text{id}_X, \text{const}_1) \uparrow & & (\text{id}_Y, \text{const}_1) \uparrow & \nearrow f'_2 & \\
 X & \xrightarrow{f_1} & Y & &
 \end{array}$$

Remark 1.4. (homotopy category)

Prop. 1.3 means that homotopy classes of continuous functions are the morphisms in a category whose objects are still the topological spaces.

This category (at least when restricted to spaces that admit the structure of CW-complexes) is called the classical homotopy category, often denoted

$$\text{Ho}(\text{Top}) .$$

Hence for X, Y topological spaces, then

$$\text{Hom}_{\text{Ho}(\text{Top})}(X, Y) = [X, Y]$$

Moreover, sending a continuous function to its homotopy class is a functor

$$\kappa : \text{Top} \longrightarrow \text{Ho}(\text{Top})$$

from the ordinary category Top of topological spaces with actual continuous functions between them.

Definition 1.5. (homotopy equivalence)

Let $X, Y \in \text{Top}$ be two topological spaces.

A continuous function

$$f : X \rightarrow Y$$

is called a homotopy equivalence if there exists

1. a continuous function the other way around,

$$g : Y \rightarrow X$$

2. homotopies (def. 1.2) from the two composites to the respective identity function:

$$f \circ g \Rightarrow \text{id}_Y$$

and

$$g \circ f \Rightarrow \text{id}_X .$$

We indicate that a continuous function is a homotopy equivalence by writing

$$X \xrightarrow{\simeq h} Y .$$

If there exists *some* (possibly unspecified) homotopy equivalence between topological spaces X and Y we write

$$X \simeq_h Y .$$

Remark 1.6. (homotopy equivalences are the isomorphisms in the homotopy category)

In view of remark 1.4 a continuous function f is a homotopy equivalence precisely if its image $\kappa(f)$ in the homotopy category is an isomorphism.

As an object of the homotopy category, a topological space is often referred to as a (strong) homotopy type. Homotopy types have a different nature than the topological spaces which *present* them, in that topological spaces that are far from being homeomorphic may still be equivalent as homotopy types.

Example 1.7. (homeomorphism is homotopy equivalence)

Every homeomorphism is a homotopy equivalence (def. 1.5).

Proposition 1.8. (homotopy equivalence is equivalence relation)

Being homotopy equivalent is an equivalence relation on the class of topological spaces.

Proof. This is immediate from remark 1.6 by general properties of categories and functors.

But for the record we spell it out. This involves the construction already used in the proof of prop. 1.3:

It is clear that the relation is reflexive and symmetric. To see that it is transitive

consider continuous functions

$$\begin{array}{ccc} X & \xrightarrow{f_1} & Y \xrightarrow{f_2} Z \\ & \xleftarrow{g_1} & \xleftarrow{g_2} \end{array}$$

and homotopies

$$\begin{array}{ll} g_1 \circ f_1 \Rightarrow \text{id}_X & f_1 \circ g_1 \Rightarrow \text{id}_Y \\ g_2 \circ f_2 \Rightarrow \text{id}_Y & f_2 \circ g_2 \Rightarrow \text{id}_Z . \end{array}$$

We need to produce homotopies of the form

$$(g_1 \circ g_2) \circ (f_2 \circ f_1) \Rightarrow \text{id}_X$$

and

$$(f_2 \circ f_1) \circ (g_1 \circ g_2) \Rightarrow \text{id}_Y .$$

Now the diagram

$$\begin{array}{ccccc} X & & \xrightarrow{f_1} & & Y \\ (\text{id}_X, \text{const}_0) \downarrow & & (\text{id}_Y, \text{const}_0) \downarrow & & \searrow g_2 \circ f_2 \\ X \times [0, 1] & \xrightarrow{f_1 \times \text{id}_{[0,1]}} & Y \times [0, 1] & \xrightarrow{\eta} & Y \xrightarrow{g_1} X, \\ (\text{id}_X, \text{const}_1) \uparrow & & (\text{id}, \text{const}_1) \uparrow & & \nearrow \text{id}_Y \\ X & & \xrightarrow{f_1} & & Y \end{array}$$

with η one of the given homotopies, exhibits a homotopy

$(g_1 \circ g_2) \circ (f_2 \circ f_1) \Rightarrow g_1 \circ f_1$. Composing this with the given homotopy $g_1 \circ f_1 \Rightarrow \text{id}_X$ gives the first of the two homotopies required above. The second one follows by the same construction, just with the labels of the functions exchanged. ■

Definition 1.9. ([contractible topological space](#))

A [topological space](#) X is called [contractible](#) if the unique [continuous function](#) to the [point space](#)

$$X \xrightarrow{\simeq h} *$$

is a [homotopy equivalence](#) (def. 1.5).

Remark 1.10. ([contractible topological spaces](#) are the [terminal objects](#) in the [homotopy category](#))

In view of remark 1.4, a topological space X is [contractible](#) (def. 1.9) precisely if its image $\kappa(X)$ in the [classical homotopy category](#) is a [terminal object](#).

Example 1.11. ([closed ball](#) and [Euclidean space](#) are [contractible](#))

Let $B^n \subset \mathbb{R}^n$ be the unit [open ball](#) or [closed ball](#) in [Euclidean space](#). This is [contractible](#) (def. 1.9):

$$p : B^n \xrightarrow{\simeq h} *$$

The homotopy inverse function is necessarily constant on a point, we may just as well choose it to go pick the origin:

$$\text{const}_0 : * \longrightarrow B^n .$$

For one way of composing these functions we have the [equality](#)

$$p \circ \text{const}_0 = \text{id}_*$$

with the [identity function](#). This is a homotopy by prop. 1.3.

The other composite is

$$\text{const}_0 \circ p = \text{const}_0 : B^n \longrightarrow B^n .$$

Hence we need to produce a homotopy

$$\text{const}_0 \Rightarrow \text{id}_{B^n}$$

This is given by the function

$$\begin{array}{ccc} B^n \times [0, 1] & \xrightarrow{\eta} & B^n \\ (x, t) & \mapsto & tx \end{array} ,$$

where on the right we use the multiplication with respect to the standard [real vector space](#) structure in \mathbb{R}^n .

Since the [open ball](#) is [homeomorphic](#) to the whole [Cartesian space](#) \mathbb{R}^n (this example) it follows with example 1.7 and example 1.3 that also \mathbb{R}^n is a contractible topological space:

$$\mathbb{R}^n \xrightarrow{\simeq h} *$$

In direct generalization of the construction in example 1.11 one finds further examples as follows:

Example 1.12. The following three [graphs](#)



(i.e. the evident [topological subspaces](#) of the [plane](#) \mathbb{R}^2 that these pictures indicate) are not [homeomorphic](#). But they are [homotopy equivalent](#), in fact they are each homotopy equivalent to the [disk](#) with two points removed, by the

homotopies indicated by the following pictures:



graphics grabbed from *Hatcher*

Fundamental group

Definition 1.13. (homotopy relative boundary)

Let X be a topological space and let

$$\gamma_1, \gamma_2 : [0, 1] \rightarrow X$$

be two paths in X , i.e. two continuous functions from the closed interval to X , such that their endpoints agree:

$$\gamma_1(0) = \gamma_2(0) \quad \gamma_1(1) = \gamma_2(1) .$$

Then a homotopy relative boundary from γ_1 to γ_2 is a homotopy (def. 1.2)

$$\eta : \gamma_1 \Rightarrow \gamma_2$$

such that it does not move the endpoints:

$$\eta(0, -) = \text{const}_{\gamma_1(0)} = \text{const}_{\gamma_2(0)} \quad \eta(1, -) = \text{const}_{\gamma_1(1)} = \text{const}_{\gamma_2(1)} .$$

Proposition 1.14. (homotopy relative boundary is equivalence relation on sets of paths)

Let X be a topological space and let $x, y \in X$ be two points. Write

$$P_{x,y}X$$

for the set of paths γ in X with $\gamma(0) = x$ and $\gamma(1) = y$.

Then homotopy relative boundary (def. 1.13) is an equivalence relation on $P_{x,y}X$.

The corresponding set of equivalence classes is denoted

$$\text{Hom}_{\pi_1(X)}(x, y) := (P_{x,y}X) / \sim .$$

Recall the operations on paths: path concatenation $\gamma_2 \cdot \gamma_1$, path reversion $\bar{\gamma}$ and constant paths

Proposition 1.15. (concatenation of homotopy relative boundary-classes of paths)

For X a topological space, then the operation of path concatenation descends to homotopy relative boundary equivalence classes, so that for all $x, y, z \in X$ there is a function

$$\begin{aligned} \text{Hom}_{\pi_1(X)}(x, y) \times \text{Hom}_{\pi_1(X)}(y, z) &\longrightarrow \text{Hom}_{\pi_1(X)}(x, z) \\ ([\gamma_1], [\gamma_2]) &\mapsto [\gamma_2] \cdot [\gamma_1] := [\gamma_2 \cdot \gamma_1] \end{aligned}$$

Moreover,

1. this composition operation is associative in that for all $x, y, z, w \in X$ and $[\gamma_1] \in \text{Hom}_{\pi_1(X)}(x, y)$, $[\gamma_2] \in \text{Hom}_{\pi_1(X)}(y, z)$ and $[\gamma_3] \in \text{Hom}_{\pi_1(X)}(z, w)$ then

$$[\gamma_3] \cdot ([\gamma_2] \cdot [\gamma_1]) = ([\gamma_3] \cdot [\gamma_2]) \cdot [\gamma_1]$$

2. this composition operation is unital with neutral elements the constant paths in that for all $x, y \in X$ and $[\gamma] \in \text{Hom}_{\pi_1(X)}(x, y)$ we have

$$[\text{const}_y] \cdot [\gamma] = [\gamma] = [\gamma] \cdot [\text{const}_x] .$$

3. this composition operation has inverse elements given by path reversal in that for all $x, y \in X$ and $[\gamma] \in \text{Hom}_{\pi_1(X)}(x, y)$ we have

$$[\bar{\gamma}] \cdot [\gamma] = [\text{const}_x] \quad [\gamma] \cdot [\bar{\gamma}] = [\text{const}_y] .$$

Definition 1.16. (fundamental groupoid and fundamental groups)

Let X be a topological space. Then set of points of X together with the sets $\text{Hom}_{\pi_1(X)}(x, y)$ of homotopy relative boundary-classes of paths (def. 1.13) for all points of points and equipped with the concatenation operation from prop. 1.15 is called the fundamental groupoid of X , denoted


$$\Pi_1(X) .$$

Given a choice of point $x \in X$, then one writes

$$\pi_1(X, x) := \text{Hom}_{\pi_1(X)}(x, x) .$$

Prop. 1.15 says that under concatenation of paths, this set is a group. As such it is called the fundamental group of X at x .

The following picture indicates the four non-equivalent non-trivial generators of the fundamental group of the oriented surface of genus 2:



graphics grabbed from [Lawson 03](#)

Example 1.17. ([fundamental group of Euclidean space](#))

For $n \in \mathbb{N}$ and $x \in \mathbb{R}^n$ any point in the n -dimensional [Euclidean space](#) (regarded with its [metric topology](#)) we have that the [fundamental group](#) (def. 1.16) at that point is trivial:

$$\pi_1(\mathbb{R}^n, x) = *.$$

Remark 1.18. (basepoints)

Definition 1.16 intentionally offers two variants of the definition.

The first, the [fundamental groupoid](#) is canonically given, without choosing a basepoint. As a result, it is a structure that is not quite a [group](#) but, slightly more generally, a “[groupoid](#)” (a “group with many objects”). We discuss the concept of [groupoids](#) below.

The second, the [fundamental group](#), is a genuine group, but its definition requires picking a base point $x \in X$.

In this context it is useful to say that

1. a [pointed topological space](#) (X, x) is
 1. a [topological space](#) X ;
 2. a $x \in X$ in the underlying set.
2. a [homomorphism](#) of pointed topological spaces $f : (X, x) \rightarrow (Y, y)$ is a base-point preserving continuous function, namely
 1. a [continuous function](#) $f : X \rightarrow Y$
 2. such that $f(x) = y$.

Hence there is a [category](#), to be denoted, $\text{Top}^{*/}$, whose [objects](#) are the [pointed topological spaces](#), and whose [morphisms](#) are the base-point preserving continuous functions.

Similarly, a [homotopy](#) between morphisms $f, f' : (X, x) \rightarrow (Y, y)$ in $\text{Top}^{*/}$ is a

homotopy $\eta: f \Rightarrow f'$ of underlying continuous functions, as in def. 1.2, such that the corresponding function

$$\eta : X \times [0, 1] \rightarrow Y$$

preserves the basepoints in that

$$\forall_{t \in [0, 1]} \eta(x, t) = y .$$

These pointed homotopies still form an equivalence relation as in prop. 1.3 and hence quotienting these out yields the pointed analogue of the homotopy category from def. 1.4, now denoted

$$\kappa : \mathrm{Top}^{*/} \rightarrow \mathrm{Ho}(\mathrm{Top}^{*/}) .$$

In general it is hard to explicitly compute the fundamental group of a topological space. But often it is already useful to know if two spaces have the same fundamental group or not:

Definition 1.19. (pushforward of elements of fundamental groups)

Let (X, x) and (Y, y) be pointed topological space (remark 1.18) and let

$$f : X \rightarrow Y$$

be a continuous function which respects the chosen points, in that $f(x) = y$.

Then there is an induced homomorphism of fundamental groups (def. 1.16)

$$\begin{array}{ccc} \pi_1(X, x) & \xrightarrow{f_*} & \pi_1(Y, y) \\ [\gamma] & \mapsto & [f \circ \gamma] \end{array}$$

given by sending a closed path $\gamma: [0, 1] \rightarrow X$ to the composite

$$f \circ \gamma : [0, 1] \xrightarrow{\gamma} X \xrightarrow{f} Y .$$

Remark 1.20. (fundamental group is functor on pointed topological spaces)

The pushforward operation in def. 1.19 is functorial, now on the category $\mathrm{Top}^{*/}$ of pointed topological spaces (remark 1.18)

$$\pi_1 : \mathrm{Top}^{*/} \rightarrow \mathrm{Grp} .$$

Proposition 1.21. (fundamental group depends only on homotopy classes)

Let $X, Y \in \mathrm{Top}^{*/}$ be pointed topological space and let $f_1, f_2 : X \rightarrow Y$ be two base-point preserving continuous functions. If there is a pointed homotopy (def. 1.2, remark 1.18)

$$\eta : f_1 \Rightarrow f_2$$

then the induced [homomorphisms](#) on fundamental groups (def. 1.19) agree

$$(f_1)_* = (f_2)_* : \pi_1(X, x) \rightarrow \pi_1(Y, y) .$$

In particular if $f : X \rightarrow Y$ is a [homotopy equivalence](#) (def. 1.5) then $f_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ is an [isomorphism](#).

Proof. This follows by the fact that homotopy respects composition (prop. 1.3):

If $\gamma : [0, 1] \rightarrow X$ is a closed path representing a given element of $\pi_1(X, x)$, then the homotopy $f_1 \Rightarrow f_2$ induces a homotopy

$$f_1 \circ \gamma \Rightarrow f_2 \circ \gamma$$

and therefore these represent the same elements in $\pi_1(Y, y)$.

It follows that if f is a homotopy equivalence with homotopy inverse g , then $g_* : \pi_1(Y, y) \rightarrow \pi_1(X, x)$ is an [inverse morphism](#) to $f_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ and hence f_* is an [isomorphism](#). ■

Remark 1.22. Prop. 1.21 says that the fundamental group functor from def. 1.19 and remark 1.20 factors through the [classical pointed homotopy category](#) from remark 1.18:

$$\begin{array}{ccc} \mathbf{Top}^{*/} & \xrightarrow{\pi_1} & \mathbf{Grp} \\ \kappa \downarrow & \nearrow & \\ \mathbf{Ho}(\mathbf{Top}^{*/}) & & \end{array} .$$

Definition 1.23. ([simply connected topological space](#))

A topological space X for which

1. $\pi_0(X) \simeq *$ ([path connected](#))
2. $\pi_1(X, x) \simeq 1$ (the [fundamental group](#) is [trivial](#), def. 1.16),

is called [simply connected](#).

We will need also the following local version:

Definition 1.24. ([semi-locally simply connected topological space](#))

A [topological space](#) X is called [semi-locally simply connected](#) if every point $x \in X$ has a [neighbourhood](#) $U_x \subset X$ such that every loop in X is contractible as a loop in X , hence such that the induced morphism of [fundamental groups](#) (def. 1.19)

$$\pi_1(U, x) \rightarrow \pi_1(X, x)$$

is trivial (i.e. sends everything to the [neutral element](#)).

If every x has a neighbourhood U_x which is itself simply connected, then X is called a [locally simply connected topological space](#). This implies semi-local simply-connectedness.

Example 1.25. ([Euclidean space](#) is [simply connected](#))

For $n \in \mathbb{N}$, then the [Euclidean space](#) \mathbb{R}^n is a [simply connected topological space](#) (def. 1.23).

Groupoids


In def. 1.16 we extracted the [fundamental group](#) at some point $x \in X$ from a larger algebraic structure, that incorporates all the basepoints, to be called the [fundamental groupoid](#). This larger algebraic structure of [groupoids](#) is usefully made explicit for the formulation and proof of the [fundamental theorem of covering spaces](#) (theorem 2.25 below) and the development of [homotopy theory](#) in general.

Where a [group](#) may be thought of as a *group of symmetry transformations* that [isomorphically](#) relates one [object](#) to itself (the [symmetries](#) of one object, such as the [isometries](#) of a [polyhedron](#)) a *groupoid* is a collection of symmetry transformations acting between possibly more than one object.

Hence a groupoid consists of a [set](#) of objects x, y, z, \dots and for each [pair](#) of objects (x, y) there is a set of transformations, usually denoted by arrows

$$x \xrightarrow{f} y$$

which may be composed if they are composable (i.e. if the first ends where the second starts)



$$\begin{array}{ccc} & y & \\ f \nearrow & & \searrow g \\ x & \xrightarrow{g \circ f} & z \end{array}$$

such that this composition is [associative](#) and such that for each object x there is identity transformation $x \xrightarrow{\text{id}_x} x$ in that this is a [neutral element](#) for the composition of transformations, whenever defined.

So far this structure is what is called a [small category](#). What makes this a ([small](#)) *groupoid* is that all these transformations are to be “symmetries” in that they are [invertible morphisms](#) meaning that for each transformation $x \xrightarrow{f} y$ there is a

transformation the other way around $y \xrightarrow{f^{-1}} x$ such that

$$f^{-1} \circ f = \text{id}_x \quad f \circ f^{-1} = \text{id}_y .$$

If there is only a single object x , then this definition reduces to that of a [group](#), and in this sense groupoids are “groups with many objects”. Conversely, given any groupoid \mathcal{G} and a choice of one of its objects x , then the subcollection of transformations from and to x is a group, sometimes called the [automorphism group](#) $\text{Aut}_{\mathcal{G}}(x)$ of x in \mathcal{G} .

Just as for groups, the “transformations” above need not necessarily be given by concrete transformations (say by [bijections](#) between [objects](#) which are [sets](#)). Just as for groups, such a concrete realization is always possible, but is an extra choice (called a [representation](#) of the groupoid). Generally one calls these “transformations” [morphisms](#): $x \xrightarrow{f} y$ is a morphism with “[source](#)” x and “[domain](#)” y .

An archetypical example of a groupoid is the [fundamental groupoid](#) $\Pi_1(X)$ of a [topological space](#) (def. 1.27 below, for introduction see [here](#)): For X a topological space, this is the groupoid whose

- [objects](#) are the points $x \in X$;
- [morphisms](#) $x \xrightarrow{[\gamma]} y$ are the [homotopy relative boundary-equivalence classes](#) $[\gamma]$ of [paths](#) $\gamma: [0, 1] \rightarrow X$ in X , with $\gamma(0) = x$ and $\gamma(1) = y$;

and [composition](#) is given, on representatives, by [concatenation](#) of paths. Here the class of the [reverse path](#) $\bar{\gamma} : t \mapsto \gamma(1 - t)$ constitutes the inverse morphism, making this a groupoid.

If one *chooses* a point $x \in X$, then the corresponding group at that point is the [fundamental group](#) $\pi_1(X, x) := \text{Aut}_{\Pi_1(X)}(x)$ of X at that point.

This highlights one of the reasons for being interested in groupoids over groups: Sometimes this allows to avoid unnatural ad-hoc choices and it serves to streamline and simplify the theory.

A [homomorphism](#) between groupoids is the obvious: a [function](#) between their underlying [objects](#) together with a function between their morphisms which respects [source](#) and [target](#) objects as well as [composition](#) and [identity morphisms](#). If one thinks of the groupoid as a special case of a [category](#), then this is a [functor](#). Between groupoids with only a single object this is the same as a [group homomorphism](#).

For example if $f : X \rightarrow Y$ is a [continuous function](#) between topological spaces, then postcomposition of [paths](#) with this function induces a groupoid homomorphism $f_* : \Pi_1(X) \rightarrow \Pi_1(Y)$ between the [fundamental groupoids](#) from above.

Groupoids with groupoid homomorphisms ([functors](#)) between them form a

[category Grp](#) (def. 1.33 below) which includes the category [Grp](#) of [groups](#) as the [full subcategory](#) of the groupoids with a single object. This makes precise how groupoid theory is a generalization of [group theory](#).

However, for groupoids more than for groups one is typically interested in “[conjugation actions](#)” on homomorphisms. These are richer for groupoids than for groups, because one may conjugate with a different morphism at each object. If we think of groupoids as special cases of [categories](#), then these “conjugation actions on homomorphisms” are [natural transformations](#) between [functors](#).

For examples if $f, g : X \rightarrow Y$ are two [continuous functions](#) between [topological spaces](#), and if $\eta : f \Rightarrow g$ is a [homotopy](#) from f to g , then the [homotopy relative boundary](#) classes of the [paths](#) $\eta(x, -) : [0, 1] \rightarrow Y$ constitute a natural

transformation between $f_*, g_* : \Pi_1(X) \rightarrow \Pi_1(Y)$ in that for all paths $x_1 \xrightarrow{[\gamma]} x_2$ in X we have the “conjugation relation”

$$\begin{array}{ccc} f(x_1) & \xrightarrow{[\eta(x_1, -)]} & g(x_1) \\ [\eta(x_1, -)] \cdot [f \circ \gamma] = [g \circ \gamma] \cdot [\eta(x_2, -)] & \text{i.e.} & \begin{array}{ccc} [f \circ \gamma] \downarrow & & \downarrow [f \circ \gamma] \\ f(x_1) & \xrightarrow{[\eta(x_2, -)]} & g(x_2) \end{array} \end{array}$$

Definition 1.26. ([groupoid](#))

A [small groupoid](#) \mathcal{G} is

1. a [set](#) X , to be called the *set of [objects](#)*;
2. for all [pairs](#) of objects $(x, y) \in X \times X$ a [set](#) $\text{Hom}(x, y)$, to be called the *set of [morphisms](#) with [domain](#) or [source](#) x and [codomain](#) or [target](#) y* ;
3. for all [triples](#) of objects $(x, y, z) \in X \times X \times X$ a [function](#)

$$\circ_{x,y,z} : \text{Hom}(y, z) \times \text{Hom}(x, y) \rightarrow \text{Hom}(x, z)$$

to be called [composition](#)

4. for all objects $x \in X$ an element

$$\text{id}_x \in \text{Hom}(x, x)$$

to be called the [identity morphism](#) on x ;

5. for all pairs $x, y \in \text{Hom}(x, y)$ of objects a function

$$(-)^{-1} : \text{Hom}(x, y) \rightarrow \text{Hom}(y, x)$$

to be called the [inverse-assigning function](#)

such that

1. ([associativity](#)) for all [quadruples](#) of objects $x_1, x_2, x_3, x_4 \in X$ and all [triples](#) of

morphisms $f \in \text{Hom}(x_1, x_2)$, $g \in \text{Hom}(x_2, x_3)$ and $h \in \text{Hom}(x_3, x_4)$ an equality

$$h \circ (g \circ f) = (h \circ g) \circ f$$

2. (unitality) for all pairs of objects $x, y \in X$ and all morphisms $f \in \text{Hom}(x, y)$ equalities

$$\text{id}_y \circ f = f \quad f \circ \text{id}_x = f$$

3. (invertibility) for all pairs of objects $x, y \in X$ and every morphism $f \in \text{Hom}(x, y)$ equalities

$$f^{-1} \circ f = \text{id}_x \quad f \circ f^{-1} = \text{id}_y .$$

If $\mathcal{G}_1, \mathcal{G}_2$ are two groupoids, then a homomorphism or functor between them, denoted

$$F : \mathcal{G}_1 \rightarrow \mathcal{G}_2$$

is

1. a function $F_0 : X_1 \rightarrow X_2$ between the respective sets of objects;
2. for each pair $x, y \in X_1$ of objects a function

$$F_{x,y} : \text{Hom}_{\mathcal{G}_1}(x, y) \rightarrow \text{Hom}_{\mathcal{G}_2}(F_0(x), F_0(y))$$

between sets of morphisms

such that

1. (respect for composition) for all triples $x, y, z \in X_1$ and all $f \in \text{Hom}(x, y)$ and $g \in \text{Hom}(y, z)$ an equality

$$F_{y,z}(g) \circ_2 F_{x,y}(f) = F_{x,z}(g \circ_1 f)$$

2. (respect for identities) for all $x \in X$ an equality

$$F_{x,x}(\text{id}_x) = \text{id}_{F_0(x)} .$$

For $\mathcal{G}_1, \mathcal{G}_2$ two groupoids, and for $F, G : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ two groupoid homomorphisms/functors, then a *conjugation* or homotopy or natural transformation (necessarily a natural isomorphism)

$$\eta : F \Rightarrow G$$

is

- for each object $x \in X_1$ of \mathcal{G}_1 a morphism $\eta_x \in \text{Hom}_{\mathcal{G}_2}(F(x), G(x))$

such that

- for all $x, y \in X_1$ and $f \in \text{Hom}_{\mathcal{G}_1}(x, y)$ an [equality](#)

$$\eta_y \circ_2 F(f) = G(f) \circ \eta_x \quad \begin{array}{ccc} F(x) & \xrightarrow{\eta_x} & G(x) \\ F(f) \downarrow & & \downarrow G(f) \\ F(y) & \xrightarrow{\eta_y} & G(y) \end{array}$$

For $\mathcal{G}_1, \mathcal{G}_2$ two groupoids and $F, G, H: \mathcal{G}_1 \rightarrow \mathcal{G}_2$ three functors between them and $\eta_1 : F \Rightarrow G$ and $\eta_2 : G \Rightarrow H$ conjugation actions/natural isomorphisms between these, there is the composite

$$\eta_2 \circ \eta_1 : F \Rightarrow H$$

with components the composite of the components

$$(\eta_2 \circ \eta_1)(x) := \eta_2(x) \circ \eta_1(x) .$$

This yields for any two groupoid a [hom-groupoid](#)

$$\text{Hom}_{\text{Grpd}}(\mathcal{G}_1, \mathcal{G}_2)$$

whose objects are the groupoid homomorphisms / functors, and whose morphisms are the conjugation actions / natural transformations.

The archetypical example of a groupoid we already encountered above:

Example 1.27. ([fundamental groupoid](#))

For X a [topological space](#), then its [fundamental groupoid](#) (as in def. 1.16) has as set of objects the underlying set of X , and for $x, y \in X$ two points, the set of homomorphisms is the set of [paths](#) from x to y modulo [homotopy relative boundary](#):

$$\text{Hom}_{\Pi_1(X)}(x, y) = (P_{x, y}) / \sim_h$$

and [composition](#) is given by [concatenation of paths](#).

Remark 1.28. ([groupoids](#) are special cases of [categories](#))

A [small groupoid](#) (def. 1.26) is equivalently a [small category](#) in which all [morphisms](#) are [isomorphisms](#).

While therefore groupoid theory may be regarded as a special case of [category theory](#), it is noteworthy that the two theories are quite different in character. For example [higher groupoid theory](#) is [homotopy theory](#) which is rich but quite tractable, for instance via tools such as [simplicial homotopy theory](#) or [homotopy type theory](#), while [higher category theory](#) is intricate and becomes tractable mostly by making recourse to higher groupoid theory in the guise of [\(infinity,1\)-category theory](#) and [\(infinity,n\)-categories](#).

Example 1.29. (groupoid [core](#) of a [category](#))

For \mathcal{C} any ([small](#)) [category](#), then there is a maximal groupoid inside

$$\mathrm{Core}(\mathcal{C}) \hookrightarrow \mathcal{C}$$

sometimes called the [core](#) of \mathcal{C} . This is obtained from \mathcal{C} simply by discarding all those [morphisms](#) that are not [isomorphisms](#).

For instance

- For $\mathcal{C} = \mathbf{Set}$ then $\mathrm{Core}(\mathbf{Set})$ is the groupoid of [sets](#) and [bijections](#) between them.

For $\mathcal{C} = \mathbf{FinSet}$ then the [skeleton](#) of this groupoid (prop. 1.47) is the disjoint union of deloopings (example 1.41) of all the [symmetric groups](#):

$$\mathrm{Core}(\mathbf{FinSet}) \simeq \bigsqcup_{n \in \mathbb{N}} \Sigma(n)$$

- For $\mathcal{C} = \mathbf{Vect}$ then $\mathrm{Core}(\mathbf{Vect})$ is the groupoid of [vector spaces](#) and [linear bijections](#) between them.

For $\mathcal{C} = \mathbf{FinVect}$ then the [skeleton](#) of this groupoid is the disjoint union of deloopings of all the [general linear groups](#)

$$\mathrm{Core}(\mathbf{FinVect}) \simeq \bigsqcup_{n \in \mathbb{N}} \mathrm{GL}(n) .$$

Example 1.30. ([discrete groupoid](#))

For X any set, there is the [discrete groupoid](#) $\mathrm{Disc}(X)$, whose set of objects is X and whose only morphisms are [identity morphisms](#).

This is also the [fundamental groupoid](#) (example 1.27) of the [discrete topological space](#) on the set X .

Example 1.31. ([disjoint union/coproduct of groupoids](#))

Let $\{\mathcal{G}_i\}_{i \in I}$ be a [set](#) of [groupoids](#). Then their [disjoint union](#) ([coproduct](#)) is the groupoid

$$\bigsqcup_{i \in I} \mathcal{G}_i$$

whose set of objects is the disjoint union of the sets of objects of the summand groupoids, and whose sets of morphisms between two objects is that of \mathcal{G}_i if both objects are from this groupoid, and is [empty](#) otherwise.

Definitio 1.32. ([product of groupoids](#))

Let $\{\mathcal{G}_i\}_{i \in I}$ be a [set](#) of [groupoids](#). Their [product groupoid](#) is the [groupoid]

$$\prod_{i \in I} \mathcal{G}_i$$

whose set of objects is the [Cartesian product](#) of the sets of objects of the factor groupoids

$$\left(\prod_{i \in I} \mathcal{G}_i\right)_0 := \prod_{i \in I} (\mathcal{G}_i)_0$$

and whose set of [morphisms](#) between [tuples](#) $(x_i)_{i \in I}$ and $(y_i)_{i \in I}$ is the corresponding Cartesian product of morphisms, with elements denoted

$$(x_i)_{i \in I} \xrightarrow{(f_i)_{i \in I}} (y_i)_{i \in I} .$$

For instance if each of the groupoids is the [delooping](#) $\mathcal{G}_i = BG_i$ of a [group](#) G_i (example 1.40) then the product groupoid is the delooping groupoid of the [direct product group](#):

$$\prod_{i \in I} BG_i \simeq B \prod_{i \in I} G_i .$$

As another example, if $\bigsqcup_{i \in I} \mathcal{G}_i$ is the [coproduct](#) groupoid from example 1.31, and if \mathcal{G} is any groupoid, then a groupoid homomorphism of the form

$$\bigsqcup_{i \in I} \mathcal{G}_i \rightarrow \mathcal{G}$$

is equivalently a [tuple](#) $(f_i)_{i \in I}$ of groupoid homomorphisms

$$\mathcal{G}_i \xrightarrow{f_i} \mathcal{G} .$$

The analogous statement holds for homotopies between groupoid homomorphisms, and so one find that the [hom-groupoid](#) out of a coproduct of groupoids is the product groupoid of the separate hom-groupoids:

$$\mathrm{Hom}_{\mathrm{Grpd}}\left(\bigsqcup_{i \in I} \mathcal{G}_i, \mathcal{G}\right) \simeq \prod_{i \in I} \mathrm{Hom}_{\mathrm{Grpd}}(\mathcal{G}_i, \mathcal{G}) .$$

Remark 1.33. ([1-category of groupoids](#))

From def. 1.26 we see that there is a [category](#) whose

- [objects](#) are the small groupoids;
- [morphisms](#) are the groupoid homomorphisms ([functors](#)).

But since this [1-category](#) does not reflect the existence of [homotopies/natural isomorphisms](#) between homomorphisms/[functors](#) of groupoids (def. 1.26) this [1-category](#) is not what one is interested in when considering [homotopy theory/higher category theory](#).

In order to obtain the right notion of category of groupoids that does reflect homotopies, we first consider now the *horizontal* composition of

homotopies/natural transformations.

Lemma 1.34. (horizontal composition of homotopies with morphisms)

Let $\mathcal{G}_1, \mathcal{G}_2, \mathcal{G}_3, \mathcal{G}_4$ be groupoid and let

$$\mathcal{G}_1 \xrightarrow{F_1} \mathcal{G}_2 \quad \begin{array}{c} \xrightarrow{F'_2} \\ \Downarrow \eta \\ \xrightarrow{F_2} \end{array} \mathcal{G}_3 \xrightarrow{F_3} \mathcal{G}_3$$

be morphisms and a homotopy η . Then there is a homotopy

$$\mathcal{G}_1 \xrightarrow{F_3 \circ F'_2 \circ F_1} \mathcal{G}_2 \quad \begin{array}{c} \xrightarrow{F_3 \circ F'_2 \circ F_1} \\ \Downarrow F_2 \cdot \eta \cdot F_1 \\ \xrightarrow{F_3 \circ F'_2 \circ F_1} \end{array} \mathcal{G}_2$$

between the respective composites, with components given by

$$(F_2 \cdot \eta \cdot F_1)(x) := F_2(\eta(F_1(x))) .$$

This operation constitutes a groupoid homomorphism/functor

$$F_3 \cdot (-) \cdot F_1 : \text{Hom}_{\text{Grpd}}(\mathcal{G}_2, \mathcal{G}_3) \rightarrow \text{Hom}_{\text{Grpd}}(\mathcal{G}_1, \mathcal{G}_4) .$$

Proof. The respect for identities is clear. To see the respect for composition, let

$$\begin{array}{c} \xrightarrow{F} \\ \Downarrow \eta_1 \\ \mathcal{G}_2 \xrightarrow{G} \mathcal{G}_3 \\ \Downarrow \eta_2 \\ \xrightarrow{H} \end{array}$$

be two composable homotopies. We need to show that

$$F_3 \cdot (\eta_2 \circ \eta_1 \cdot F_1) = (F_3 \cdot \eta_2 \cdot F_1) \circ (F_3 \cdot \eta_1 \cdot F_1) .$$

Now for x any object of \mathcal{G}_1 we find

$$\begin{aligned} (F_3 \cdot (\eta_2 \circ \eta_1 \cdot F_1))(x) &:= F_2((\eta_2 \circ \eta_1)(F_1(x))) \\ &:= F_3(\eta_2(F_1(x)) \circ \eta_1(F_1(x))) \\ &= F_2(\eta_2(F_1(x))) \circ F_2(\eta_1(F_1(x))) \\ &= ((F_3 \cdot \eta_2 \cdot F_1) \circ (F_3 \cdot \eta_1 \cdot F_1))(x) \end{aligned}$$

Here all steps are unwinding of the definition of horizontal and of ordinary (vertical) composition of homotopies, except the third equality, which is the functoriality of F_2 . ■

Lemma 1.35. (horizontal composition of homotopies)

Consider a [diagram](#) of [groupoids](#), groupoid homomorphisms (functors) and homotopies (natural transformations) as follows:

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{F_1} & \mathcal{G}_2 & \xrightarrow{F_2} & \mathcal{G}_3 \\ \Downarrow \eta_1 & & \Downarrow \eta_2 & & \\ \mathcal{G}_1 & \xrightarrow{F'_1} & \mathcal{G}_2 & \xrightarrow{F'_2} & \mathcal{G}_3 \end{array}$$

The [horizontal composition](#) of the homotopies to a single homotopy of the form

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{F_2 \circ F_1} & \mathcal{G}_3 \\ \Downarrow \eta_2 \cdot \eta_1 & & \\ \mathcal{G}_1 & \xrightarrow{F'_2 \circ F'_1} & \mathcal{G}_3 \end{array}$$

may be defined in terms of the horizontal composition of homotopies with morphisms (lemma 1.34) and the ("vertical") composition of homotopies with themselves, in two different ways, namely by decomposing the above diagram as

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{F_1} & \mathcal{G}_2 & \xrightarrow{F_2} & \mathcal{G}_3 \\ \Downarrow \eta_1 & & & & \\ \mathcal{G}_1 & \xrightarrow{F'_1} & \mathcal{G}_2 & & \mathcal{G}_3 \end{array}$$

$$\begin{array}{ccccc} \mathcal{G}_1 & & \mathcal{G}_2 & \xrightarrow{F_2} & \mathcal{G}_3 \\ & \xrightarrow{F'_1} & \mathcal{G}_2 & \Downarrow \eta_2 & \\ & & \mathcal{G}_2 & \xrightarrow{F'_2} & \mathcal{G}_3 \end{array}$$

or as

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{F_1} & \mathcal{G}_2 & \xrightarrow{F_2} & \mathcal{G}_3 \\ & & \Downarrow \eta_2 & & \\ & & \mathcal{G}_2 & \xrightarrow{F'_2} & \mathcal{G}_3 \end{array}$$

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{F_1} & \mathcal{G}_2 & & \mathcal{G}_3 \\ \Downarrow \eta_1 & & & \xrightarrow{F'_2} & \\ \mathcal{G}_1 & \xrightarrow{F'_1} & \mathcal{G}_2 & & \mathcal{G}_3 \end{array}$$

In the first case we get

$$\eta_2 \cdot \eta_1 := (\eta_2 \cdot F'_1) \circ (F_2 \cdot \eta_1)$$

while in the second case we get

$$\eta_2 \cdot \eta_1 := (F'_2 \cdot \eta_1) \circ (\eta_2 \cdot F_1) .$$

These two definitions coincide.

Proof. For x an object of \mathcal{G}_1 , then we need that the following square [diagram commutes](#) in \mathcal{G}_3

$$\begin{array}{ccc}
 F_2(F_1(x)) & \xrightarrow{(F_2 \cdot \eta_1)(x)} & F_2(F'_1(x)) \\
 (\eta_2 \cdot F_1)(x) \downarrow & & \downarrow (\eta_2 \cdot F'_1)(x) \\
 F'_2(F_1(x)) & \xrightarrow{(F'_2 \cdot \eta_1)(x)} & F'_2(F'_1(x))
 \end{array}
 =
 \begin{array}{ccc}
 F_2(F_1(x)) & \xrightarrow{F_2(\eta_1(x))} & F_2(F'_1(x)) \\
 \eta_2(F_1(x)) \downarrow & & \downarrow \eta_2(F'_1(x)) \\
 F'_2(F_1(x)) & \xrightarrow{F'_2(\eta_1(x))} & F'_2(F'_1(x))
 \end{array}$$

But the commutativity of the square on the right is the defining compatibility condition on the components of η_2 applied to the morphism $\eta_1(x)$ in \mathcal{G}_2 . ■

Proposition 1.36. (horizontal composition with homotopy is natural transformation)

Consider groupoids, homomorphisms and homotopies of the form

$$\begin{array}{ccc}
 \mathcal{G}_1 & \xrightarrow{F_1} & \mathcal{G}_2 \\
 \Downarrow \eta_1 & & \\
 \mathcal{G}_1 & \xrightarrow{F'_1} & \mathcal{G}_2
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{G}_3 & \xrightarrow{F_3} & \mathcal{G}_4 \\
 \Downarrow \eta_3 & & \\
 \mathcal{G}_3 & \xrightarrow{F'_3} & \mathcal{G}_4
 \end{array}
 .$$

Then horizontal composition with the homotopies (lemma 1.35) constitutes a natural transformation between the functors of horizontal composition with morphisms (lemma 1.34)

$$(\eta_3 \cdot (-) \cdot \eta_1) : (F_3 \cdot (-) \cdot F_1) \Rightarrow (F'_3 \cdot (-) \cdot F'_1) : \text{Hom}_{\text{Grpd}}(\mathcal{G}_2, \mathcal{G}_2) \rightarrow \text{Hom}_{\text{Grpd}}(\mathcal{G}_1, \mathcal{G}_4) .$$

Proof. By lemma 1.35. ■

It first of all follows that the following makes sense

Definition 1.37. (homotopy category of groupoids)

There is also the homotopy category $\text{Ho}(\text{Grpd})$ whose

- objects are small groupoids;
- morphisms are equivalence classes of groupoid homomorphisms modulo homotopy (i.e. functors modulo natural transformations).

This is usually denoted $\text{Ho}(\text{Grpd})$.

Of course what the above really means is that, without quotienting out homotopies, groupoids form a 2-category, in fact a (2,1)-category, in fact an enriched category which is enriched over the naive 1-category of groupoids from remark 1.33, hence a strict 2-category with hom-groupoids.

Definition 1.38. (equivalence of groupoids)

Given two groupoids \mathcal{G}_1 and \mathcal{G}_2 , then a homomorphism

$$F : \mathcal{G}_1 \rightarrow \mathcal{G}_2$$

is an equivalence if it is an isomorphism in the homotopy category $\text{Ho}(\text{Grpd})$ (def. 1.37), hence if there exists a homomorphism the other way around

$$G : \mathcal{G}_2 \rightarrow \mathcal{G}_1$$

and a homotopy/natural transformations of the form

$$G \circ F \simeq \text{id}_{\mathcal{G}_1} \quad F \circ G \simeq \text{id}_{\mathcal{G}_2} .$$

Example 1.39. ((2,1)-functoriality of fundamental groupoid)

If X and Y are [topological spaces](#) and $f : X \rightarrow Y$ is a [continuous function](#) between them, then this induces a groupoid homomorphism (functor) between the respective [fundamental groupoids](#) (def. 1.27)

$$F_f : \Pi_1(X) \rightarrow \Pi_1(Y)$$

given on objects by the underlying function of f

$$(F_f)_0 := f$$

and given on the class of a [path](#) by the evident postcomposition with f

$$(F_f)_{x,y} : (x \xrightarrow{[\gamma]} y) \mapsto (f(x) \xrightarrow{[f \circ \gamma]} f(y)) .$$

This construction clearly respects [identity morphisms](#) and [composition](#) and hence is itself a [functor](#) of the form

$$\Pi_1 : \mathbf{Top} \rightarrow \mathbf{Grpd}_1$$

from the [category Top](#) of [topological space](#) to the [1-category Grpd](#) of groupoids.

But more is true: If $f, g : X \rightarrow Y$ are two [continuous function](#) and

$$\eta : f \Rightarrow g$$

is a [left homotopy](#) between them, hence a continuous function

$$\eta : X \times [0, 1] \rightarrow Y$$

such that $\eta(-, 0) = f$ and $\eta(-, 1) = g$, then this induces a homotopy between the above groupoid homomorphisms (a natural transformation of functors).

This shows that the fundamental groupoid functor in fact descends to [homotopy categories](#)

$$\Pi_1 : \mathbf{Ho}(\mathbf{Top}) \rightarrow \mathbf{Ho}(\mathbf{Grpd}) .$$

(In fact this means it even extends to a [\(2,1\)-functor](#) from the [\(2,1\)-category](#) of topological spaces, continuous functions, and [higher homotopy](#)-classes of left homotopies, to that of groupoids.)

As a direct consequence it follows that if there is a [homotopy equivalence](#)

$$X \simeq_h Y$$

between [topological spaces](#), then there is an induced [equivalence of groupoids](#) between their [fundamental groupoids](#)

$$\Pi_1(X) \simeq \Pi_1(Y) .$$

Hence the [fundamental groupoid](#) is a [homotopy invariant](#) of topological spaces. Of course by prop. 1.46 the fundamental groupoid is equivalent, as a groupoid, to the disjoint union of the [deloopings](#) of all the [fundamental groups](#) of the given topological spaces, one for each [connected component](#), and hence this is equivalently the statement that the set of connected components and the fundamental groups of a topological space are homotopy invariants.

Example 1.40. ([delooping of a group](#))

Let G be a [group](#). Then there is a groupoid, denoted BG , with a single object p , with morphisms

$$\mathrm{Hom}_{BG}(p, p) := G$$

the elements of G , with composition the multiplication in G , with identity morphism the [neutral element](#) in G and with inverse morphisms the inverse elements in G .

This is also called the [delooping](#) of G (because the [loop space object](#) of BG at the unique point is the given group: $\Omega BG \simeq G$).

For G_1, G_2 two groups, then there is a [natural bijection](#) between [group homomorphisms](#) $\phi: G_1 \rightarrow G_2$ and groupoid homomorphisms $BG_1 \rightarrow BG_2$: the latter are all of the form $B\phi$, with $(B\phi)_0$ uniquely fixed and $(B\phi)_{p,p} = \phi$.

This means that the construction $B(-)$ is a [fully faithful functor](#)

$$B(-) : \mathbf{Grp} \hookrightarrow \mathbf{Grpd}_1$$

into from the category [Grp](#) of groups to the [1-category](#) of [groupoids](#).

But beware that this functor is not fully faithful when homotopies of groupoids are taken into account, because there are in general non-trivial homotopies between morphisms of the form

$$B\phi_1, B\phi_2 : BG \rightarrow BH$$

By definition, such a homotopy (natural transformation) $\eta : B\phi_1 \Rightarrow B\phi_2$ is a choice of a single element $\eta_p \in H$ such that for all $g \in G$ we have

$$\begin{array}{ccc} p & \xrightarrow{h} & p \\ \phi_1(g) \downarrow & & \downarrow \phi_2(g) \\ p & \xrightarrow[h]{} & p \end{array} \quad \phi_2(g) = h \cdot \phi_1(g) \cdot h^{-1}$$

hence such that

$$\phi_2 = \text{Ad}_h \circ \phi_1 .$$

Therefore notably the induced functor

$$B(-) : \text{Grp} \rightarrow \text{Ho}(\text{Grp})$$

to the [homotopy category](#) of groupoids is not fully faithful.

But since BG is canonically a [pointed object](#) in groupoids, we may also regard [delooping](#) as a functor

$$B(-) : \text{Grp} \rightarrow \text{Grpd}^{*/}$$

to the [category of pointed objects](#) of [Grpd](#). Since groupoid homomorphisms $BG_1 \rightarrow BG_2$ necessarily preserve the basepoint, this makes no difference at this point. But as we now pass to the [homotopy category](#)

$$B(-) : \text{Grp} \hookrightarrow \text{Ho}(\text{Grpd}^{*/})$$

then also the homotopies are required to preserve the absepoint, and for homotopies between homomorphisms between delooped groups this means, since there only is a single point, that these homotopies are all trivial. Hence regarded this way the functor is a [fully faithful functor](#) again, hence an [equivalence of categories](#) onto its [essential image](#). By prop. 1.47 below this essential image consists precisely of the (pointed) [connected](#) groupoids:

Groups are equivalently pointed connected groupoids.

Example 1.41. ([disjoint union](#) of [delooping](#) groupoids)

Let $\{G_i\}_{i \in I}$ be a [set](#) of [groups](#). Then there is a groupoid $\bigsqcup_{i \in I} BG_i$ which is the disjoint union groupoid (example 1.31) of the [delooping](#) groupoids BG_i (example 1.40).

Its set of objects is the index set I , and

$$\text{Hom}(i, j) = \begin{cases} G_i & | \quad i = j \\ \emptyset & | \quad \text{otherwise} \end{cases}$$

Definition 1.42. ([connected components](#) of a groupoid)

Given a [groupoid](#) \mathcal{G} with set of objects X , then the [relation](#) “there exists a morphism from x to y ”, i.e.

$$(x \sim y) := (\text{Hom}(x, y) \neq \emptyset)$$

is clearly an [equivalence relation](#) on X . The corresponding set of [equivalence classes](#) is denoted

$$\pi_0(\mathcal{G})$$

and called the set of connected components of \mathcal{G} .

Definition 1.43. (automorphism groups)

Given a groupoid \mathcal{G} and an object x , then under composition the set $\mathrm{Hom}_{\mathcal{G}}(x, x)$ forms a group. This is called the automorphism group $\mathrm{Aut}_{\mathcal{G}}(x)$ or *vertex group* or *isotropy group* of x in \mathcal{G} .

For each object x in a groupoid \mathcal{G} , there is a canonical groupoid homomorphism

$$B \mathrm{Aut}_{\mathcal{G}}(x) \hookrightarrow \mathcal{G}$$

from the delooping groupoid (def. 1.40) of the automorphism group. This takes the unique object of $B \mathrm{Aut}_{\mathcal{G}}(x)$ to x and takes every automorphism of x “to itself”, regarded now again as a morphism in \mathcal{G} .

Definition 1.44. (weak homotopy equivalence of groupoids)

Let \mathcal{G}_1 and \mathcal{G}_2 be groupoids. Then a morphism (functor)

$$F : \mathcal{G}_1 \rightarrow \mathcal{G}_2$$

is called a weak homotopy equivalence if

1. it induces a bijection on connected components (def. 1.42):

$$\pi_0(F) : \pi_0(\mathcal{G}_1) \xrightarrow{\cong} \pi_0(\mathcal{G}_2)$$

2. for each object x of \mathcal{G}_1 the morphism

$$F_{x,x} : \mathrm{Aut}_{\mathcal{G}_1}(x) \xrightarrow{\cong} \mathrm{Aut}_{\mathcal{G}_2}(F_0(X))$$

is an isomorphism of automorphism groups (def. 1.43)

Lemma 1.45. (automorphism group depends on basepoint only up to conjugation)

For \mathcal{G} a groupoid, let x and y be two objects in the same connected component (def. 1.42). Then there is a group isomorphism

$$\mathrm{Aut}_{\mathcal{G}}(x) \simeq \mathrm{Aut}_{\mathcal{G}}(y)$$

between their automorphism groups (def. 1.43).

Proof. By assumption, there exists some morphism from x to y

$$x \xrightarrow{f} y.$$

The operation of conjugation with this morphism

$$\begin{aligned} \mathrm{Aut}_{\mathcal{G}}(x) &\xrightarrow{\mathrm{Ad}_f} \mathrm{Aut}_{\mathcal{G}}(y) \\ g &\mapsto f^{-1} \circ g \circ f \end{aligned}$$

is clearly a group isomorphism as required. ■

Lemma 1.46. (equivalences between disjoint unions of delooping groupoids)

Let $\{G_i\}_{i \in I}$ and $\{H_j\}_{j \in J}$ be sets of groups and consider a homomorphism (functor)

$$F : \sqcup_{i \in I} G_i \longrightarrow \sqcup_{j \in J} H_j$$

between the corresponding disjoint unions of delooping groupoids (example 1.40).

Then the following are equivalent:

1. F is an equivalence of groupoids (def. 1.38);
2. F is a weak homotopy equivalence (def. 1.44).

Proof. The implication $2) \Rightarrow 1)$ is immediate.

In the other direction, assume that F is an equivalence of groupoids, and let G be an inverse up to natural isomorphism. It is clear that both induces bijections on connected components. To see that both are isomorphisms of automorphisms groups, observe that the conditions for the natural isomorphisms

$$\alpha : G \circ F \Rightarrow \mathrm{id} \qquad \beta : F \circ G \Rightarrow \mathrm{id}$$

are in each separate delooping groupoid BH_j of the form

$$\begin{array}{ccc} * & \xrightarrow{\alpha} & * \\ F_{F_0(i), F_0(i)}(F_{i,i}(f)) \downarrow & & \downarrow \mathrm{id} \\ * & \xrightarrow{\alpha} & * \end{array} \qquad \begin{array}{ccc} * & \xrightarrow{\beta} & * \\ F_{G_0(j), G_0(j)}(G_{j,j}(f)) \downarrow & & \downarrow \mathrm{id} \\ * & \xrightarrow{\beta} & * \end{array}$$

since there is only a single object. But this means $F_{i,i}$ and $F_{j,j}$ are group isomorphisms. ■

Proposition 1.47. (every groupoid is equivalent to a disjoint union of group deloopings)

Assuming the axiom of choice, then:

For \mathcal{G} any groupoid, then there exists a set $\{G_i\}_{i \in I}$ of groups and an equivalence of groupoids (def. 1.38)

$$\mathcal{G} \simeq \sqcup_{i \in I} BG_i$$

between \mathcal{G} and a [disjoint union](#) of [delooping](#) groupoids (example 1.41). This is called a [skeleton](#) of \mathcal{G} .

Concretely, this exists for $I = \pi_0(\mathcal{G})$ the set of [connected components](#) of \mathcal{G} (def. 1.42) and for $G_i := \text{Aut}_{\mathcal{G}}(x_i)$ the [automorphism group](#) (def. 1.43) of any object x_i in the given connected component.

Proof. Using the [axiom of choice](#) we may find a set $\{x_i\}_{i \in \pi_0(\mathcal{G})}$ of objects of \mathcal{G} , with x_i being in the [connected component](#) $i \in \pi_0(\mathcal{G})$.

This choice induces a functor

$$\text{inc} : \bigsqcup_{i \in \pi_0(\mathcal{G})} \text{Aut}_{\mathcal{G}}(x_i) \rightarrow \mathcal{G}$$

which takes each object and morphism “to itself”.

Now using the [axiom of choice](#) once more, we choose in each connected component $i \in \pi_0(\mathcal{G})$ and for each object y in that connected component a morphism

$$x_i \xrightarrow{f_{x_i, y}} y .$$

Using this we obtain a functor the other way around

$$p : \mathcal{G} \rightarrow \bigsqcup_{i \in \pi_0(\mathcal{G})} \text{Aut}_{\mathcal{G}}(x_i)$$

which sends each object to its connected component, and which for pairs of objects y, z of \mathcal{G} is given by conjugation with the morphisms chosen above:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{G}}(y, z) & \xrightarrow{p_{y, z}} & \text{Aut}_{\mathcal{G}}(x_i) \\ y & & y \xleftarrow{f_{x_i, y}} x_i \\ f \downarrow & \mapsto & f \downarrow \\ z & & z \xrightarrow[f_{x_i, z}^{-1}]{} x_i \end{array} .$$

It is now sufficient to show that there are conjugations/natural isomorphisms

$$p \circ \text{inc} \simeq \text{id} \quad \text{inc} \circ p \simeq \text{id} .$$

For the first this is immediate, since we even have equality

$$p \circ \text{inc} = \text{id} .$$

For the second we observe that choosing

$$\eta(y) := f_{x_i, y}$$

yields a naturality square by the above construction:

$$\begin{array}{ccc}
 x_i & \xrightarrow{f_{x_i,y}} & y \\
 f_{x_i,z} \circ f \circ f_{x_i,y}^{-1} \downarrow & & \downarrow f \\
 x_i & \xrightarrow{f_{x_i,z}} & z
 \end{array}$$

■

Proposition 1.48. (**weak homotopy equivalence is equivalence of groupoids**)

Let $F : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ be a homomorphism of groupoids.

Assuming the axiom of choice then the following are equivalent:

1. F is an equivalence of groupoids (def. 1.38);
2. F is a weak homotopy equivalence in that
 1. it induces an bijection of sets of connected components (def. 1.42);

$$\pi_0(F) : \pi_0(\mathcal{G}_1) \xrightarrow{\cong} \pi_0(\mathcal{G}_2),$$

3. for each object $x \in \mathcal{G}_1$ it induces an isomorphism of automorphism groups (def. 1.43):

$$F_{x,x} : \text{Aut}_{\mathcal{G}_1}(x) \xrightarrow{\cong} \text{Aut}_{\mathcal{G}_2}(F_0(x)) .$$

Proof. In one direction, if F has an inverse up to natural isomorphism, then this induces by definition a bijection on connected components, and it induces isomorphism on homotopy groups by lemma 1.45.

In the other direction, choose equivalences to skeleta as in prop. 1.47 to get a commuting diagram in the 1-category of groupoids as follows:

$$\begin{array}{ccc}
 \mathcal{G}_1 & \xleftarrow[\cong]{\text{inc}_1} & \bigsqcup_{i \in \pi_0(\mathcal{G}_1)} \text{Aut}_{\mathcal{G}_1}(x_i) \\
 F \downarrow & & \downarrow \tilde{F} \\
 \mathcal{G}_2 & \xleftarrow[\text{inc}_2]{\cong} & \bigsqcup_{i \in \pi_0(\mathcal{G}_1)} \text{Aut}_{\mathcal{G}_2}(F_0(x_i))
 \end{array} .$$

Here inc_1 and inc_2 are equivalences of groupoids by prop. 1.47. Moreover, by assumption that F is a weak homotopy equivalence \tilde{F} is the union of deloopings of isomorphisms of groups, and hence has a strict inverse, in particular a homotopy inverse, hence is in particular an equivalence of groupoids.

In conclusion, when regarded as a diagram in the homotopy category $\text{Ho}(\text{Grpd})$ (def. 1.37), the top, bottom and right morphism of the above diagram are isomorphisms. It follows that also f is an isomorphism in $\text{Ho}(\text{Grpd})$. But this means exactly that it is a homotopy equivalence of groupoids, by def. 1.38. ■

2. Covering spaces

A *covering space* (def. 2.1 below) is a “continuous [fiber bundle](#) of sets” over a topological space, in just the same way as a [topological vector bundle](#) is a “continuous [fiber bundle](#) of vector spaces”.

Definition 2.1. ([covering space](#))

Let X be a [topological space](#). A [covering space](#) of X is a [continuous function](#)

$$p:E \rightarrow X$$

such that there exists an [open cover](#) $\sqcup_i U_i \rightarrow X$, such that restricted to each U_i then $E \rightarrow X$ is [homeomorphic](#) over U_i to the [product topological space](#) ([this example](#)) of U_i with the [discrete topological space](#) ([this example](#)) on a [set](#) F_i ,

In summary this says that $p:E \rightarrow X$ is a covering space if there exists a [pullback diagram](#) in [Top](#) of the form

$$\begin{array}{ccc} \sqcup_i U_i \times \text{Disc}(F_i) & \longrightarrow & E \\ \downarrow & \text{(pb)} & \downarrow p \\ \sqcup_{i \in I} U_i & \longrightarrow & X \end{array}$$

For $x \in U_i \subset X$ a point, then the elements in $F_x = F_i$ are called the [leaves](#) of the covering at x .

Given two covering spaces $p_i:E_i \rightarrow X$, then a [homomorphism](#) between them is a [continuous function](#) $f:E_1 \rightarrow E_2$ between the total covering spaces, which respects the [fibers](#) in that the following [diagram commutes](#)

$$\begin{array}{ccc} E_1 & \xrightarrow{f} & E_2 \\ p_1 \searrow & & \swarrow p_2 \\ & X & \end{array}$$

This defines a [category](#) $\text{Cov}(X)$, the [category of covering spaces](#) over X , whose

- [objects](#) are the covering spaces over X ;
- [morphisms](#) are the homomorphisms between these.

Example 2.2. ([trivial covering space](#))

For X a [topological space](#) and S a [set](#) with $\text{Disc}(S)$ the [discrete topological space](#) on that set, then the [projection](#) out of the [product topological space](#)

$$\text{pr}_1 : X \times \text{Disc}(S) \rightarrow X$$

is a [covering space](#), called the *trivial covering space* over X with [fiber](#) $\text{Disc}(S)$.


If $E \xrightarrow{p} X$ is any covering space, then an [isomorphism](#) of covering spaces of the form

$$\begin{array}{ccc} E & \xrightarrow{\cong} & X \times \text{Disc}(S) \\ p \searrow & & \swarrow \text{pr}_2 \\ & X & \end{array}$$

is called a *trivialization* of $E \xrightarrow{p} X$.

It is in this sense that every covering space E is, by definition, locally trivializable.

Example 2.3. (covering of circle by circle)



Regard the [circle](#) $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ as the [topological subspace](#) of elements of unit [absolute value](#) in the [complex plane](#). For $k \in \mathbb{N}$, consider the continuous function

$$p := (-)^k : S^1 \rightarrow S^1$$

given by taking a complex number to its k th power. This may be thought of as the result of “winding the circle k times around itself”. Precisely, for $k \geq 1$ this is a [covering space](#) (def. 2.1) with k leaves at each point.

graphics grabbed from [Hatcher](#)

Example 2.4. (covering of circle by real line)

Consider the [continuous function](#)

$$\exp(2\pi i(-)) : \mathbb{R}^1 \rightarrow S^1$$

from the [real line](#) to the [circle](#), which,

1. with the circle regarded as the unit circle in the [complex plane](#) \mathbb{C} , is given by

$$t \mapsto \exp(2\pi i t)$$

2. with the circle regarded as the unit circle in \mathbb{R}^2 , is given by

$$t \mapsto (\cos(2\pi t), \sin(2\pi t)) .$$



We may think of this as the result of “winding the line around the circle ad infinitum”. Precisely, this is a [covering space](#) (def. 2.1) with the [leaves](#) at each point forming the set \mathbb{Z} of [natural numbers](#).

Here are some basic properties of covering spaces:

Proposition 2.5. ([covering projections are open maps](#))

If $p:E \rightarrow X$ is a covering space projection, then p is an [open map](#).

Proof. By definition of covering space there exists an [open cover](#) $\{U_i \subset X\}_{i \in I}$ and [homeomorphisms](#) $p^{-1}(U_i) \simeq U_i \times \text{Disc}(F_i)$ for all $i \in I$. Since the [projections](#) out of a [product topological space](#) are [open maps](#) (this prop.), it follows that p is an open map when restricted to any of the $p^{-1}(U_i)$. But a general open subset $W \subset E$ is the union of its restrictions to these subspaces:

$$W = \bigcup_{i \in I} (W \cap p^{-1}(U_i)) .$$

Since images preserve unions (this prop.) it follows that

$$p(W) = \bigcup_{i \in I} p(W \cap p^{-1}(U_i))$$

is a union of open sets, and hence itself open. ■

Lemma 2.6. ([fiber-wise diagonal of covering space is open and closed](#))

Let $E \xrightarrow{p} X$ be a covering space. Consider the [fiber product](#)

$$E \times_X E := \{(e_1, e_2) \in E \times E \mid p(e_1) = p(e_2)\}$$

hence (by the discussion at [Top - Universal constructions](#)) the [topological subspace](#) of the [product space](#) $E \times E$, as shown on the right. By the [universal property](#) of the [fiber product](#), there is the [diagonal continuous function](#)

$$\begin{array}{ccc} E & \longrightarrow & E \times_X E \\ e & \mapsto & (e, e) \end{array} .$$

Then the [image](#) of E under this function is an [open subset](#) and a [closed subset](#):

$$\Delta(E) \subset E \times_X E \text{ is open and closed .}$$

Proof. First to see that it is an open subset. It is sufficient to show that for any $e \in E$ there exists an open neighbourhood of $(e, e) \in E \times_X E$.

Now by definition of covering spaces, there exists an open neighbourhood $U_{p(e)} \subset X$ of $p(e) \in X$ such that

$$\begin{array}{ccc}
 U_{p(e)} \times \text{Disc}(p^{-1}(p(e))) & \xrightarrow{\cong} & E|_{U_{p(e)}} \\
 \text{pr}_1 \searrow & & \swarrow p \\
 & U_{p(e)} &
 \end{array}
 .$$

It follows that $U_{p(e)} \times \{e\} \subset E$ is an open neighbourhood. Hence by the nature of the [product topology](#), $U_{p(e)} \times U_{p(e)} \subset E \times E$ is an open neighbourhood of (e, e) in $E \times E$ and hence by the nature of the [subspace topology](#) the restriction

$$(E \times_X E) \cap (U_{p(e)} \times U_{p(e)}) \subset E \times_X E$$

is an open neighbourhood of (e, e) in $E \times_X E$.

Now to see that the diagonal is closed, hence that the complement $(E \times_X E) \setminus \Delta(E)$ is an open subset, it is sufficient to show that every point (e_1, e_2) with $e_1 \neq e_2$ but $p(e_1) = p(e_2)$ has an open neighbourhood in this complement.

As before, there is an open neighbourhood $U \subset X$ of $p(e_1) = p(e_2)$ over which the cover trivializes, and hence $U \times \{e_1\}, U \times \{e_2\} \subset E$ are open neighbourhoods of e_1 and e_2 , respectively. These are disjoint by the assumption that $e_1 \neq e_2$. As above, this means that the intersection

$$(E \times_X E) \cap ((U \times \{e_1\}) \times (U \times \{e_2\})) \subset (E \times_X E) \setminus \Delta(E)$$

is an open subset of the complement of the diagonal in the fiber product. ■

Lifting properties

If $E \xrightarrow{p} X$ is any [continuous function](#) (possibly a [covering space](#) or a [topological vector bundle](#)) then a [section](#) is a continuous function $\sigma: X \rightarrow E$ which sends each point in the base to a point in the [fiber](#) above it, hence which makes this [diagram commute](#):

$$\begin{array}{ccc}
 & E & \\
 \sigma \nearrow & & \downarrow p \\
 X & = & X
 \end{array}$$

We may think of this as “lifting” each point in the base to point in the fibers “through” the projection map p . More generally if $Y \hookrightarrow X$ is a subspace, we may consider such lifts only over Y

$$\begin{array}{ccc}
 & E & \\
 \sigma \nearrow & & \downarrow p \\
 Y & \hookrightarrow & X
 \end{array}$$

sometimes called a “local section”. But this suggests that for $Y \xrightarrow{f} X$ any continuous function, we consider “lifting its image through p ”

$$\begin{array}{ccc} & E & \\ \sigma \nearrow & & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

For example if $Y = [0, 1]$ is the [topological interval](#), then $f: [0, 1] \rightarrow X$ is a [path](#) in the base space X , and a lift through p of this is a path in the total space which “runs above” the given path. Such lifts of paths through covering projections is the topic of [monodromy](#) below.

Here it is of interest to consider the lifting problem subject to some constraint. For instance we will want to consider lifts of paths $\gamma: [0, 1] \rightarrow X$ through a covering projection, subject to the condition that the starting point $\gamma(0)$ is lifted to a prescribed point $p \in E$.

Since such a point is equivalently a continuous function $\text{const}_p: * \rightarrow X$ out of the [point space](#), this is the same as asking for a continuous function σ that makes both triangles in the following [diagram commute](#):

$$\begin{array}{ccccc} * & & \xrightarrow{\text{const}_p} & E & \\ \text{const}_0 \downarrow & & \sigma \nearrow & & \downarrow p \\ [0, 1] & & \xrightarrow{\gamma} & X & \end{array}$$

This is an example of a general situation which plays a central role in [homotopy theory](#): We say that a square [commuting diagram](#)

$$\begin{array}{ccc} A & \longrightarrow & E \\ i \downarrow & & \downarrow p \\ B & \longrightarrow & X \end{array}$$

is a [lifting problem](#) and that a diagonal morphism

$$\begin{array}{ccc} A & \longrightarrow & E \\ i \downarrow & \nearrow & \downarrow p \\ B & \longrightarrow & X \end{array}$$

such that both resulting triangles commute is a [lift](#). If such a lift exists for for the given p and for each i taken from some [class](#) of morphisms, then one says that p has the [right lifting property](#) against this class.

We now discuss some [right lifting properties](#) satisfied by covering spaces:

1. [homotopy-lifting property](#),
2. the [lifting theorem](#).

These lifting properties will be used in [below](#) for the computation of [fundamental groups](#) of some topological spaces.

Lemma 2.7. ([lifts](#) out of [connected space](#) into [covering spaces](#) are unique relative to any point)

Let

1. $E \xrightarrow{p} X$ be a covering space,
2. Y a [connected topological space](#)
3. $f : Y \rightarrow X$ a [continuous function](#).
4. $\hat{f}_1, \hat{f}_2 : Y \rightarrow E$ two [lifts](#) of f , in that the following [diagram commutes](#):

$$\begin{array}{ccc} & E & \\ \hat{f}_i \nearrow & & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

for $i \in \{1, 2\}$.

If there exists $y \in Y$ such that $\hat{f}_1(y) = \hat{f}_2(y)$ then the two lifts already agree everywhere: $\hat{f}_1 = \hat{f}_2$.

Proof. By the [universal property](#) of the [fiber product](#)

$$E \times_X E := \{(e_1, e_2) \in E \times E \mid p(e_1) = p(e_2)\} \subset E \times E$$

the two lifts determine a single continuous function of the form

$$(\hat{f}_1, \hat{f}_2) : Y \rightarrow E \times_X E.$$

Write

$$\Delta(E) := \{(e, e) \in E \times_X E \mid e \in E\}$$

for the [diagonal](#) on E in the fiber product. By lemma [2.6](#) this is an open subset and a closed subset of the fiber product space. Hence by continuity of (\hat{f}_1, \hat{f}_2) also its pre-image

$$(\hat{f}_1, \hat{f}_2)^{-1}(\Delta(E)) \subset Y$$

is both closed and open, hence also its [complement](#) is open in Y .

Moreover, the assumption that the functions \hat{f}_1 and \hat{f}_2 agree in at least one point

means that the above pre-image is [non-empty](#). Therefore the assumption that Y is [connected](#) implies that this pre-image coincides with all of Y . This is the statement to be proven. ■

Lemma 2.8. (*path lifting property*)

Let $p:E \rightarrow X$ be any [covering space](#). Given

1. $\gamma:[0,1] \rightarrow X$ a [path](#) in X ,
2. $\hat{x}_0 \in E$ be a lift of its starting point, hence such that $p(\hat{x}_0) = \gamma(0)$

then there exists a unique path $\hat{\gamma}:[0,1] \rightarrow E$ such that

1. it is a lift of the original path: $p \circ \hat{\gamma} = \gamma$;
2. it starts at the given lifted point: $\hat{\gamma}(0) = \hat{x}_0$.

In other words, every [commuting diagram](#) in [Top](#) of the form

$$\begin{array}{ccc} \{0\} & \xrightarrow{\hat{x}_0} & E \\ \downarrow & & \downarrow p \\ [0,1] & \xrightarrow{\gamma} & X \end{array}$$

has a unique [lift](#):

$$\begin{array}{ccccc} \{0\} & & \xrightarrow{\hat{x}_0} & & E \\ \downarrow & \hat{\gamma} \nearrow & & \downarrow p. & \\ [0,1] & & \xrightarrow{\gamma} & & X \end{array}$$

Proof. First consider the case that the covering space is trivial, hence of the [Cartesian product](#) form

$$\text{pr}_1 : X \times \text{Disc}(S) \rightarrow X .$$

By the [universal property](#) of the [product topological spaces](#) in this case a lift $\hat{\gamma}:[0,1] \rightarrow X \times \text{Disc}(S)$ is equivalently a [pair](#) of continuous functions

$$\text{pr}_1(\hat{\gamma}):[0,1] \rightarrow X \quad \text{pr}_2(\hat{\gamma}):[0,1] \rightarrow \text{Disc}(S) ,$$

Now the lifting condition explicitly fixes $\text{pr}_1(\hat{\gamma}) = \gamma$. Moreover, a continuous function into a [discrete topological space](#) $\text{Disc}(S)$ is [locally constant](#), and since $[0,1]$ is a [connected topological space](#) this means that $\text{pr}_2(\hat{\gamma})$ is in fact a [constant function](#) ([this example](#)), hence uniquely fixed to be $\text{pr}_2(\hat{\gamma}) = \hat{x}_0$.

This shows the statement for the case of trivial covering spaces.

Now consider any covering space $p:E \rightarrow X$. By definition of covering spaces, there exists for every point $x \in X$ a [open neighbourhood](#) $U_x \subset X$ such that the restriction of E to U_x becomes a trivial covering space:

$$p^{-1}(U_x) \simeq U_x \times \text{Disc}(p^{-1}(x)) .$$

Consider such a choice

$$\{U_x \subset X\}_{x \in X} .$$

This is an [open cover](#) of X . Accordingly, the [pre-images](#)

$$\{\gamma^{-1}(U_x) \subset [0, 1]\}_{x \in X}$$

constitute an open cover of the [topological interval](#) $[0, 1]$.

Now the [closed interval](#) is a [compact topological space](#), so that this cover has a finite open subcover. By the [Euclidean metric topology](#), each element in this finite subcover is a disjoint union of open intervals. The collection of all these open intervals is an open refinement of the original cover, and by compactness it once more has a finite subcover, now such that each element of the subcover is guaranteed to be a single open interval.

This means that we find a [finite number](#) of points

$$t_0 < t_1 < \cdots < t_{n+1} \in [0, 1]$$

with $t_0 = 0$ and $t_{n+1} = 1$ such that for all $0 < j \leq n$ there is $x_j \in X$ such that the corresponding path segment

$$\gamma([t_j, t_{j+1}]) \subset X$$

is contained in U_{x_j} from above.

Now assume that $\hat{\gamma}|_{[0, t_j]}$ has been found. Then by the triviality of the covering space over U_{x_j} and the first argument above, there is a unique lift of $\gamma|_{[t_j, t_{j+1}]}$ to a continuous function $\hat{\gamma}|_{[t_j, t_{j+1}]}$ with starting point $\hat{\gamma}(t_j)$. Since $[0, t_{j+1}]$ is the [pushout](#) $[0, t_j] \sqcup_{\{t_j\}} [t_j, t_{j+1}]$ ([this example](#)), it follows that $\hat{\gamma}|_{[0, t_j]}$ and $\hat{\gamma}|_{[t_j, t_{j+1}]}$ uniquely glue to a continuous function $\hat{\gamma}|_{[0, t_{j+1}]}$ which lifts $\gamma|_{[0, t_{j+1}]}$.

By [induction](#) over j , this yields the required lift $\hat{\gamma}$.

Conversely, given any lift, $\hat{\gamma}$, then its restrictions $\hat{\gamma}|_{[t_j, t_{j+1}]}$ are uniquely fixed by the above inductive argument. Therefore also the total lift is unique. Alternatively, uniqueness of the lifts is a special case of lemma [2.7](#). ■

Proposition 2.9. ([homotopy lifting property of covering spaces](#))

Let

1. $E \xrightarrow{p} X$ be a covering space;

2. Y a [topological space](#).

Then every [lifting problem](#) of the form

$$\begin{array}{ccc} Y & \xrightarrow{\hat{f}} & E \\ (\text{id}_Y, \text{const}_0) \downarrow & & \downarrow p \\ Y \times [0, 1] & \xrightarrow{\eta} & X \end{array}$$

has a unique [lift](#)

$$\begin{array}{ccc} Y & \xrightarrow{\hat{f}} & E \\ (\text{id}_Y, \text{const}_0) \downarrow & \hat{\eta} \nearrow & \downarrow p \\ Y \times [0, 1] & \xrightarrow{\eta} & X \end{array}$$

Proof. It is clear what the lift must be: For every point $y \in Y$ the situation restricts to that of path lifting

$$\begin{array}{ccccc} * & \xrightarrow{\text{const}_y} & Y & \xrightarrow{f} & E \\ \text{const}_0 \downarrow & & (\text{id}_Y, \text{const}_0) \downarrow & \hat{\eta} \nearrow & \downarrow p \\ [0, 1] & \xrightarrow{(\text{const}_y, \text{id})} & Y \times [0, 1] & \xrightarrow{\eta} & X \end{array}$$

And so at each point $y \in Y$ the lift of $\eta(x, -)$ must be the unique path that lifts this with starting point $\hat{f}(y)$. We just need to see that this lift is a continuous function.

To that end we generalize the proof of the path lifting to connected open neighbourhoods of points in Y :

Let $\{U_i \subset X\}_{i \in I}$ be an open cover over which the covering space trivializes. Then the pre-images $\{\eta^{-1}(U_i) \subset Y \times [0, 1]\}_{i \in I}$ is an open cover of the product space. By nature of the [product space topology](#) and the [Euclidean topology](#) on $[0, 1]$, each of the $\eta^{-1}(U_i)$ is a union of Cartesian products $V_j \times I_j$ with $V_i \subset Y$ an open subset of Y and $I_i \subset [0, 1]$ an interval. Hence there is an open cover of the form

$$\{V_j \times I_j \subset Y \times [0, 1]\}_{j \in J}$$

with the property that for each j there exists $i \in I$ with $\eta(V_j \times I_j) \subset U_i$.

Now by the fact that $[0, 1]$ is a [compact topological space](#), for each $y \in Y$ there exists a finite set $K_y \subset J$ such that

$$\{V_k \times I_k\}_{k \in K_y} \subset K$$

still restricts to a cover of $\{y\} \times [I]$. Since K is finite, the intersection

$$V_y := \bigcap_{k \in K_y}$$

is still open, and so also

$$\{V_y \times I_k\}_{k \in K_y}$$

still restricts to a cover of $\{y\} \times [0, 1]$.

This means that the same argument as for the path lifting in lemma 2.8 provides a unique lift $\widehat{\eta}|_{V_y \times [0, 1]}$ for each $y \in Y$.

Moreover, for $y_1, y_2 \in Y$ two points, these lifts clearly have to agree on $V_{y_1} \cap V_{y_2}$.

Since $\{V_y \times [0, 1] \subset Y \times [0, 1]\}_{y \in Y}$ is an open cover, means that there is a unique function $\hat{\eta}$ that restricts to all these local lifts (this prop). This is the required lift. ■

Remark 2.10. (**covering spaces are Hurewicz fibrations**)

Continuous functions that satisfy the [homotopy lifting property](#), hence that have the [right lifting property](#) against continuous functions of the form $Y \xrightarrow{(\text{id}, \text{const}_0)} Y \times [0, 1]$ are called [Hurewicz fibrations](#). Hence prop. 2.9 says that covering projections are in particular Hurewicz fibrations.

Example 2.11. (**homotopy lifting property for given lifts of paths relative starting point**)

Let $p: E \rightarrow X$ be a [covering space](#). Then given a [homotopy](#) relative the starting point between two [paths](#) in X ,

$$\eta : \gamma_1 \Rightarrow \gamma_2$$

there is for every lift $\hat{\gamma}_1, \hat{\gamma}_2$ of these two paths to paths in E with the same starting point a unique homotopy

$$\hat{\eta} : \hat{\gamma}_1 \Rightarrow \hat{\gamma}_2$$

between the lifted paths that lifts the given homotopy:

For [commuting squares](#) of the form

$$\begin{array}{ccc} ([0, 1] \times \{0\}) \cup (\{0, 1\} \times [0, 1]) & \xrightarrow{(\gamma_1, \gamma_2)} & E \\ \downarrow & \hat{\eta} \nearrow & \downarrow p \\ [0, 1] \times [0, 1] & \xrightarrow{\eta} & X \end{array}$$

there is a unique diagonal [lift](#) in the lower diagram, as shown.

Moreover if the homotopy η also fixes the endpoint, then so does the lifted homotopy $\hat{\eta}$.

Proof. There are horizontal [homeomorphisms](#) such that the following diagram commutes

$$\begin{array}{ccc} [0, 1] & \xrightarrow{\cong} & ([0, 1] \times \{0\}) \cap (\{0, 1\} \times [0, 1]) \\ \downarrow & & \downarrow \\ [0, 1] \times [0, 1] & \xrightarrow{\cong} & [0, 1] \times [0, 1] \end{array} .$$

With this the statement follows from [2.9](#). ■

Example 2.12. Let $(E, e) \xrightarrow{p} (X, x)$ be a [pointed covering space](#) and let $f: (Y, y) \rightarrow (X, x)$ be a point-preserving [continuous function](#) such that the image of the [fundamental group](#) of (Y, y) is contained within the image of the fundamental group of (E, e) in that of (X, x) :

$$f_*(\pi_1(Y, y)) \subset p_*(\pi_1(E, e)) \subset \pi_1(X, x) .$$

Then for ℓ_Y a [path](#) in (Y, y) that happens to be a [loop](#), every lift of its image path $f \circ \ell$ in (X, x) to a path $\widehat{f \circ \ell_Y}$ in (E, e) is also a loop there.

Proof. By assumption, there is a loop ℓ_E in (E, e) and a homotopy fixing the endpoints of the form

$$\eta_X : p \circ \ell_E \Rightarrow f \circ \ell_Y .$$

Then by the homotopy lifting property as in [example 2.11](#), there is a homotopy in (E, e) relative to the basepoint

$$\eta_E : \ell_E \Rightarrow \widehat{f \circ \ell_Y}$$

and lifting the homotopy η_X . Therefore η_E is in fact a homotopy between loops, and so $\widehat{f \circ \ell_Y}$ is indeed a loop. ■

Proposition 2.13. (lifting theorem)

Let

1. $p: E \rightarrow X$ be a [covering space](#);
2. $e \in E$ a point, with $x := p(e)$ denoting its image,
3. Y be a [connected](#) and [locally path-connected topological space](#);
4. $y \in Y$ a point
5. $f: (Y, y) \rightarrow (X, x)$ a [continuous function](#) such that $f(y) = x$.

Then the following are equivalent:

1. There exists a unique lift \hat{f} in the diagram

$$\begin{array}{ccc} & (E, e) & \\ \hat{f} \nearrow & & \downarrow p \\ (Y, y) & \xrightarrow{f} & (X, x) \end{array}$$

of pointed topological spaces.

2. The image of the fundamental group of Y under f in that of X is contained in the image of the fundamental group of E under p :

$$f_*(\pi_1(Y, y)) \subset p_*(\pi_1(E, e))$$

Moreover, if Y is path-connected, then the lift in the first item is unique.

Proof. The implication $1) \Rightarrow 2)$ is immediate. We need to show that the second statement already implies the first.

Since Y is connected and locally path-connected, it is also a path-connected topological space (this prop.). Hence for every point $y' \in Y$ there exists a path γ connecting y with y' and hence a path $f \circ \gamma$ connecting x with $f(y')$. By the path-lifting property (lemma 2.8) this has a unique lift

$$\begin{array}{ccc} \{0\} & \xrightarrow{e} & E \\ \downarrow \widehat{f \circ \gamma} \nearrow & & \downarrow p. \\ [0, 1] & \xrightarrow{f \circ \gamma} & X \end{array}$$

Therefore

$$\hat{f}(y') := \widehat{f \circ \gamma}(1)$$

is a lift of $f(y')$.

We claim now that this pointwise construction is independent of the choice γ , and that as a function of y' it is indeed continuous. This will prove the claim.

Now by the path lifting lemma 2.8 the lift $\widehat{f \circ \gamma}$ is unique given $f \circ \gamma$, and hence $\hat{f}(y')$ depends at most on the choice of γ .

Hence let $\gamma' : [0, 1] \rightarrow Y$ be another path in Y that connects y with y' . We need to show that then $\widehat{f \circ \gamma'} = \widehat{f \circ \gamma}$.

First observe that if γ' is related to γ by a homotopy, so that then also $f \circ \gamma'$ is related to $f \circ \gamma$ by a homotopy, then this is the statement of the homotopy lifting property as in example 2.11.

Next write $\bar{\gamma}' \cdot \gamma$ for the [path concatenation](#) of the path γ with the [reverse path](#) of the path γ' , hence a loop in Y , so that $f \circ (\bar{\gamma}' \cdot \gamma)$ is a loop in X . The assumption that $f_*(\pi_1(Y, y)) \subset p_*(\pi_1(E, e))$ implies (example [2.12](#)) that the path $\widehat{f \circ (\bar{\gamma}' \cdot \gamma)}$ which lifts this loop to E is itself a loop in E .

By uniqueness of path lifting, this means that the lift of $f \circ (\gamma' \cdot (\bar{\gamma}' \cdot \gamma))$ coincides with that of $f \circ \gamma'$. But $\bar{\gamma}' \cdot (\gamma' \cdot \gamma)$ is homotopic (via reparameterization) to just γ . Hence it follows now with the first statement that the lift of $f \circ \gamma'$ indeed coincides with that of $f \circ \gamma$.

This shows that the above prescription for \hat{f} is well defined.

It only remains to show that the function \hat{f} obtained this way is continuous.

Let $y' \in Y$ be a point and $W_{\hat{f}(y')} \subset E$ an open neighbourhood of its image in E . It is sufficient to see that there is an open neighbourhood $V_{y'} \subset Y$ such that $\hat{f}(V_{y'}) \subset W_{\hat{f}(y')}$.

Let $U_{f(y')} \subset X$ be an open neighbourhood over which p trivializes. Then the restriction

$$p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')} \subset U_{f(y')} \times \text{Disc}(p^{-1}(f(y')))$$

is an open subset of the product space. Consider its further restriction

$$(U_{f(y')} \times \{\hat{f}(y')\}) \cap (p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')})$$

to the [leaf](#)

$$U_{f(y')} \times \{\hat{f}(y')\} \subset U_{f(y')} \times p^{-1}(f(y'))$$

which is itself an open subset. Since p is an [open map](#) ([this prop.](#)), the subset

$$p\left((U_{f(y')} \times \{\hat{f}(y')\}) \cap (p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')})\right) \subset X$$

is open, hence so is its pre-image

$$f^{-1}\left(p\left((U_{f(y')} \times \{\hat{f}(y')\}) \cap (p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')})\right)\right) \subset Y.$$

Since Y is assumed to be [locally path-connected](#), there exists a path-connected open neighbourhood

$$V_{y'} \subset f^{-1}\left(p\left((U_{f(y')} \times \{\hat{f}(y')\}) \cap (p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')})\right)\right).$$

By the uniqueness of path lifting, the image of that under \hat{f} is

$$\begin{aligned}
\hat{f}(V_{y'}) &= f(V_{y'}) \times \{\hat{f}(y')\} \\
&\subset p\left(\left(U_{f(y')} \times \{\hat{f}(y')\}\right) \cap \left(p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')}\right)\right) \times \{\hat{f}(y')\} \\
&\simeq \left(U_{f(y')} \times \{\hat{f}(y')\}\right) \cap \left(p^{-1}(U_{f(y')}) \cap W_{\hat{f}(y')}\right) \\
&\subset W_{\hat{f}(y')}
\end{aligned}$$

This shows that the lifted function is continuous. Finally that this continuous lift is unique is the statement of lemma 2.7. ■

The lifting theorem implies that there are “universal” covering spaces:

Definition 2.14. ([universal covering space](#))

A [covering space](#) $E \xrightarrow{p} X$ is called a [universal covering space](#) if the total space E is a [simply connected topological space](#) (def. 1.23)

Proposition 2.15. For X a [locally path-connected topological space](#), then any two [universal covering spaces](#) over X (def. 2.14) are isomorphic.

Proof. Since both E_1 and E_2 are simply connected, the assumption of the lifting theorem for covering spaces is satisfied (prop. 2.13). This says that there are horizontal continuous function making the following diagrams commute:

$$\begin{array}{ccccc}
E_1 & \xrightarrow{f} & E_2 & & E_2 & \xrightarrow{g} & E_1 \\
p_1 \searrow & & \swarrow p_2 & & p_1 \searrow & & \swarrow p_2 \\
& & X & & & & X \\
& & E_i & \xrightarrow{\text{id}} & E_i & & \\
& & p_i \searrow & & \swarrow p_i & & \\
& & & & X & &
\end{array}$$

and that these are *unique* once we specify the image of a single point, which we may freely do (in the given fiber).

So if we pick any point $x \in X$ and $\hat{x}_1 \in E_1$ with $p(\hat{x}_1) = x$ and $\hat{x}_2 \in E_2$ with $p(\hat{x}_2) = x$ and specify that $f(\hat{x}_1) = \hat{x}_2$ and $g(\hat{x}_2) = \hat{x}_1$ then uniqueness applied to the composites implies $f \circ g = \text{id}_{E_2}$ and $g \circ f = \text{id}_{E_1}$. ■

Monodromy

Since the lift of a path through a [covering space projection](#) is unique once the lift of the starting point is chosen (lemma 2.8) every path in the base space determines a [function](#) between the [fiber sets](#) over its endpoints. By the [homotopy lifting property](#) of covering spaces as in example 2.11 this function only depends on the [equivalence class](#) of the path under [homotopy relative boundary](#). Therefore this fiber-assignment is in fact an [action](#) of the [fundamental groupoid](#)

of the base space on sets, called a [groupoid representation](#) (def. 2.16 below). In particular, associated with any homotopy-class of a [loop](#), hence of an element in the [fundamental group](#), there is associated a [bijection](#) of the fiber over the loop's basepoint with itself, hence a [permutation representation](#) of the [fundamental group](#). This is called the [monodromy](#) of the covering space. It is a measure for how the coverign space fails to be globally trivial.

In fact the [fundamental theorem of covering spaces](#) (prop. 2.25) below says that the [monodromy](#) representation characterizes the covering spaces completely and faithfully. This means that covering spaces may be dealt with completely with tools from [group theory](#) and [representation theory](#), a fact that we make use of in the computation of examples [below](#).

Definition 2.16. ([groupoid representation](#))

Let \mathcal{G} be a [groupoid](#). Then:

A [linear representation](#) of \mathcal{G} is a groupoid homomorphism ([functor](#))

$$\rho : \mathcal{G} \rightarrow \text{Core}(\text{Vect})$$

to the groupoid [core](#) of the category [Vect](#) of [vector spaces](#) (example 1.29). Hence this is

1. For each object x of \mathcal{G} a [vector space](#) V_x ;
2. for each morphism $x \xrightarrow{f} y$ of \mathcal{G} a [linear map](#) $\rho(f) : V_x \rightarrow V_y$

such that

1. (respect for composition) for all composable morphisms $x \xrightarrow{f} y \xrightarrow{g} z$ in the groupoid we have an [equality](#)

$$\rho(g) \circ \rho(f) = \rho(g \circ f)$$

2. (respect for identities) for each object x of the groupoid we have an equality

$$\rho(\text{id}_x) = \text{id}_{V_x} .$$

Similarly a [permutation representation](#) of \mathcal{G} is a groupoid homomorphism ([functor](#))

$$\rho : \mathcal{G} \rightarrow \text{Core}(\text{Set})$$

to the groupoid core of [Set](#). Hence this is

1. For each object x of \mathcal{G} a [set](#) S_x ;

2. for each morphism $x \xrightarrow{f} y$ of \mathcal{G} a [function](#) $\rho(f) : S_x \rightarrow S_y$

such that composition and identities are respected, as above.

For ρ_1 and ρ_2 two such representations, then a homomorphism of representations

$$\phi : \rho_1 \longrightarrow \rho_2$$

is a [natural transformation](#) between these functors, hence is

- for each object x of the groupoid a (linear) function

$$(V_1)_x \xrightarrow{\phi(x)} (V_2)_x$$

- such that for all morphisms $x \xrightarrow{f} y$ we have

$$\begin{array}{ccc} (V_1)_x & \xrightarrow{\phi(x)} & (V_2)_x \\ \rho_1(f) \downarrow & & \downarrow \phi_2(f) \\ (V_1)_y & \xrightarrow{\phi(y)} & (V_2)_y \end{array} \quad \phi(y) \circ \rho_1(f) = \rho_2(x) \circ \phi(x)$$

By def. [1.26](#) the representations of \mathcal{G} in $\text{Core}(\mathcal{C})$ and homomorphisms between them constitute a [groupoid](#) called the [representation groupoid](#)

$$\text{Rep}(\mathcal{G}) := \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{C})) .$$

Example/Definition 2.17. ([group representations](#) are [groupoid representations](#) of [delooping groupoids](#))

If here $\mathcal{B}G$ is the [delooping](#) groupoid of a [group](#) G (example [1.40](#)), then a [groupoid representation](#) of BG is a [group representation](#) of G (def. [2.16](#)), and one writes

$$\text{Rep}(G) := \text{Rep}(BG)$$

for the [representation groupoid](#).

For each object $x \in X$ the canonical inclusion of the delooping groupoid of the [automorphism group](#) (from def. [1.43](#))

$$\text{inc}_x : B \text{Aut}_{\text{math}} \hookrightarrow \mathcal{G}$$

induces by precomposition a homomorphism of representation groupoids:

$$\text{Hom}(\text{inc}_x, \text{Core}(\mathcal{C})) : \text{Rep}(\mathcal{G}, \mathcal{C}) \longrightarrow \text{Rep}(\text{Aut}_{\mathcal{G}}(x), \mathcal{C}) .$$

We say that a groupoid representation is *faithful* or *free* if for all objects x its restriction to a group representation of $\text{Aut}_{\mathcal{G}}(x)$ this way is transitive or free, respectively.

Here the representation ρ of a group G on some set S

1. transitive if for all pairs of elements $s_1, s_2 \in S$ there is a $g \in G$ such that $\rho(g)(s_1) = s_2$;
2. free if whenever $g(s) = s$ holds for all $s \in S$ then g is the neutral elements.

Proposition 2.18. (groupoid representations are products of group representations)

Assuming the axiom of choice then the following holds:

Let \mathcal{G} be a groupoid. Then its groupoid of groupoid representations $\text{Rep}(\mathcal{G})$ (def. 2.16) is equivalent (def. 1.38) to the product groupoid (example 1.32) indexed by the set of connected components $\pi_0(\mathcal{G})$ (def. 1.42) of group representations (example 2.17) of the automorphism group $G_i := \text{Aut}_{\mathcal{G}}(x_i)$ (def. 1.43) for x_i any object in the i th connected component:

$$\text{Rep}(\mathcal{G}) \simeq \prod_{i \in \pi_0(\mathcal{G})} \text{Rep}(G_i) .$$

Proof. Let \mathcal{C} be the category that the representation is on (e.g. $\mathcal{C} = \text{Set}$ for permutation representations). Then by definition

$$\text{Rep}(\mathcal{G}) = \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{C})) .$$

Consider the injection functor of the skeleton from lemma 1.46

$$\text{inc} : \bigsqcup_{i \in \pi_0(\mathcal{G})} BG_i \longrightarrow \mathcal{G} .$$

By lemma 1.34 the pre-composition with this constitutes a functor

$$\text{inc}^* : \text{Hom}(\mathcal{G}, \mathcal{C}) \longrightarrow \text{Hom}\left(\bigsqcup_{i \in \pi_0(\mathcal{G})} BG_i, \mathcal{C}\right)$$

and by combining lemma 1.46 with lemma 1.36 this is an equivalence of groupoids. Finally, by example 1.32 the groupoid on the right is the product groupoid as claimed. ■

Definition 2.19. (monodromy of a covering space)

Let X be a topological space and $E \xrightarrow{p} X$ a covering space (def. 2.1). Write $\Pi_1(X)$ for the fundamental groupoid of X (example 1.27).

Define a groupoid homomorphism

$$\text{Fib}_E : \Pi_1(X) \longrightarrow \text{Core}(\text{Set})$$

to the groupoid core of the category Set of sets (example 2.16), hence a permutation groupoid representation (example 2.16), as follows:

1. to a point $x \in X$ assign the **fiber** $p^{-1}(\{x\}) \in \mathbf{Set}$;
2. to the **homotopy class** of a **path** γ connecting $x := \gamma(0)$ with $y := \gamma(1)$ in X assign the function $p^{-1}(\{x\}) \rightarrow p^{-1}(\{y\})$ which takes $\hat{x} \in p^{-1}(\{x\})$ to the endpoint of a path $\hat{\gamma}$ in E which lifts γ through p with starting point $\hat{\gamma}(0) = \hat{x}$

$$\begin{aligned} p^{-1}(x) &\longrightarrow p^{-1}(y) \\ (\hat{x} = \hat{\gamma}(0)) &\mapsto \hat{\gamma}(1) \end{aligned} .$$

This construction is well defined for a given representative γ due to the unique path-lifting property of covering spaces (lemma 2.8) and it is independent of the choice of γ in the given homotopy class of paths due to the **homotopy lifting property** (example 2.11). Similarly, these two lifting properties give that this construction respects composition in $\Pi_1(X)$ and hence is indeed a homomorphism of groupoids (a functor).

Proposition 2.20. (extracting **monodromy is **functorial**)**

Given a **isomorphism** between two **covering spaces** $E_i \xrightarrow{p_i} X$, hence a **homeomorphism** $f: E_1 \rightarrow E_2$ which respects **fibers** in that the **diagram**

$$\begin{array}{ccc} E_1 & \xrightarrow{f} & E_2 \\ p_1 \searrow & & \swarrow p_2 \\ & X & \end{array}$$

commutes, then the component functions

$$f|_{\{x\}} : p_1^{-1}(\{x\}) \rightarrow p_2^{-1}(\{x\})$$

are compatible with the monodromy \mathbf{Fib}_E (def. 2.19) along any **path** γ between points x and y from def. 2.19 in that the following **diagrams of sets commute**

$$\begin{array}{ccc} p_1^{-1}(x) & \xrightarrow{f|_{\{x\}}} & p_2^{-1}(x) \\ \mathbf{Fib}_{E_1}([\gamma]) \downarrow & & \downarrow \mathbf{Fib}_{E_2}([\gamma]) \\ p_1^{-1}(y) & \xrightarrow{f|_{\{y\}}} & p_2^{-1}(y) \end{array}$$

This means that f induces a homotopy (**natural transformation**) between the monodromy homomorphisms (functors)

$$\begin{array}{ccc} & \xrightarrow{\mathbf{Fib}_{E_1}} & \\ \Pi_1(X) & \Downarrow \text{Core}(\mathbf{Set}) & \\ & \xrightarrow{\mathbf{Fib}_{E_2}} & \end{array}$$

of E_1 and E_2 , respectively, and hence that constructing monodromy is itself a

functor from the category of covering spaces of X to that of permutation representations of the fundamental groupoid of X :

$$\text{Fib} : \text{Cov}(X) \rightarrow \text{Rep}(\Pi_1(X), \text{Set}) .$$

Proof. Let $e \in p_1^{-1}(x)$ be an element, and $\hat{\gamma} : [0, 1] \rightarrow E_1$ a lift of γ to E_1 with $\hat{\gamma}(0) = e$. This means by definition of monodromy that

$$\text{Fib}_{E_1}([\gamma]) : e \mapsto \hat{\gamma}(1)$$

Now $f \circ \hat{\gamma}$ is a lift of γ to E_2 , with $(f \circ \gamma)(0) = f(\hat{\gamma}(0)) = f(e)$. Therefore

$$\text{Fib}_{E_2}([\gamma]) : f(e) \mapsto f(\hat{\gamma}(1)) .$$

This means that the square commutes, as claimed. ■

Example 2.21. (fundamental groupoid of covering space)

Let $E \xrightarrow{p} X$ be a covering space.

Then the fundamental groupoid $\Pi_1(E)$ of the total space E is the groupoid whose

- objects are pairs (x, \hat{x}) consisting of a point $x \in X$ and an element $\hat{x} \in \text{Fib}_E(x)$;
- morphisms $[\hat{\gamma}] : (x, \hat{x}) \rightarrow (x', \hat{x}')$ are morphisms $[\gamma] : x \rightarrow x'$ in $\Pi_1(X)$ such that $\text{Fib}_E([\gamma])(\hat{x}) = \hat{x}'$.

This is also called the Grothendieck construction of the monodromy functor $\text{Fib}_E : \Pi_1(X) \rightarrow \text{Core}(\text{Set})$, and denoted

$$\Pi_1(E) \simeq \int_{\Pi_1(X)} \text{Fib}_E .$$

Proof. By the uniqueness of the path-lifting, lemma 2.8 and the very definition of the monodromy functor. ■

Example 2.22. (covering space is simply connected if monodromy is free and transitive)

Let X be a path-connected topological space and let $E \xrightarrow{p} X$ be a covering space. Then the total space E is

1. path-connected precisely if the monodromy Fib_E is a transitive action;
2. simply connected (def. 1.23) precisely if the monodromy Fib_E is a transitive and free action.

Proof. By example 2.21. ■

Definition 2.23. (reconstruction of covering spaces from monodromy)

Let

1. (X, τ) be a locally path-connected semi-locally simply connected topological space,
2. $\rho \in \text{Rep}(\Pi_1(X), \text{Set})$ a permutation representation of its fundamental groupoid.

Consider the disjoint union set of all the sets appearing in this representation

$$E(\rho) := \bigsqcup_{x \in X} \rho(x)$$

For

1. $U \subset X$ an open subset
 1. which is path-connected
 2. for which every element of the fundamental group $\pi_1(U, x)$ becomes trivial under $\pi_1(U, x) \rightarrow \pi_1(X, x)$,
2. for $\hat{x} \in \rho(x)$ with $x \in U$

consider the subset

$$V_{U, \hat{x}} := \{\rho(\gamma)(\hat{x}) \mid x' \in U, \gamma \text{ path from } x \text{ to } x'\} \subset E(\rho).$$

The collection of these defines a base for a topology (prop. 2.24 below). Write τ_ρ for the corresponding topology. Then

$$(E(\rho), \tau_\rho)$$

is a topological space. It canonically comes with the function

$$\begin{array}{ccc} E(\rho) & \xrightarrow{p} & X \\ \hat{x} \in \rho(x) & \mapsto & x \end{array}.$$

Finally, for

$$f : \rho_1 \rightarrow \rho_2$$

a homomorphism of permutation representations, there is the evident induced function

$$\begin{array}{ccc} E(\rho_1) & \xrightarrow{\text{Rec}(f)} & E(\rho_2) \\ (\hat{x} \in \rho_1(x)) & \mapsto & (f_x(\hat{x}) \in \rho_2(x)) \end{array}.$$

Proposition 2.24. *The construction $\rho \mapsto E(\rho)$ in def. 2.23 is well defined and yields a [covering space](#) of X .*

Moreover, the construction $f \mapsto \text{Rec}(f)$ yields a homomorphism of covering spaces.

Proof. First to see that we indeed have a [topology](#), we need to check (by [this prop.](#)) that every point is contained in some base element, and that every point in the intersection of two base elements has a base neighbourhood that is still contained in that intersection.

So let $x \in X$ be a point. By the assumption that X is [semi-locally simply connected](#) there exists an [open neighbourhood](#) $U_x \subset X$ such that every loop in U_x on x is contractible in X . By the assumption that X is a [locally path-connected topological space](#), this contains an open neighbourhood $U'_x \subset U_x$ which is [path connected](#) and, as every subset of U_x , it still has the property that every loop in U'_x based on x is contractible as a loop in X . Now let $\hat{x} \in E$ be any point over x , then it is contained in the base open $V_{U'_x, \hat{x}}$.

The argument for the base open neighbourhoods contained in intersections is similar.

Then we need to see that $p: E(\rho) \rightarrow X$ is a [continuous function](#). Since taking pre-images preserves unions ([this prop.](#)), and since by semi-local simply connectedness and local path connectedness every neighbourhood contains an open neighbourhood $U \subset X$ that labels a base open, it is sufficient to see that $p^{-1}(U)$ is a base open. But by the very assumption on U , there is a unique morphism in $\Pi_1(X)$ from any point $x \in U$ to any other point in U , so that ρ applied to these paths establishes a bijection of sets

$$p^{-1}(U) \simeq \bigsqcup_{\hat{x} \in \rho(x)} V_{U, \hat{x}} \simeq U \times \rho(x),$$

thus exhibiting $p^{-1}(U)$ as a union of base opens.

Finally we need to see that this continuous function p is a covering projection, hence that every point $x \in X$ has a neighbourhood U such that $p^{-1}(U) \simeq U \times \rho(x)$. But this is again the case for those U all whose loops are contractible in X , by the above identification via ρ , and these exist around every point by semi-local simply-connectedness of X .

This shows that $p: E(\rho) \rightarrow X$ is a covering space. It remains to see that $\text{Rec}(f): E(\rho_1) \rightarrow E(\rho_2)$ is a homomorphism of covering spaces. Now by construction it is immediate that this is a function over X , in that this [diagram commutes](#):

$$\begin{array}{ccc}
 E(\rho_1) & \xrightarrow{\text{Rec}(f)} & E(\rho_2) \\
 \searrow & & \swarrow \\
 & X &
 \end{array}$$

So it only remains to see that $\text{Rec}(f)$ is a [continuous function](#). So consider $V_{U, y_2 \in \rho_2(x)}$ a base open of $E(\rho_2)$. By [naturality](#) of f

$$\begin{array}{ccc}
 \rho_1(x') & \xrightarrow{f_{x'}} & \rho_2(x') \\
 \rho_1(\gamma) \uparrow \simeq & & \simeq \uparrow \rho_2(\gamma) \\
 \rho_1(x) & \xrightarrow{f_x} & \rho_2(x)
 \end{array}$$

its pre-image under $\text{Rec}(f)$ is

$$\text{Rec}(f)^{-1}(V_{U, y_2 \in \rho_2(x)}) = \bigsqcup_{y_1 \in f^{-1}(y_2)} V_{U, y_1 \in \rho_1(x)}$$

and hence a union of base opens. ■

Proposition 2.25. ([fundamental theorem of covering spaces](#))

Let X be a [locally path-connected](#) and [semi-locally simply-connected topological space](#) (def. 1.24). Then the operations on

1. extracting the [monodromy](#) Fib_E of a [covering space](#) E over X (def. 2.19, prop. 2.20)
2. [reconstructing a covering space from monodromy](#) $\text{Rec}(\rho)$ (def. 2.23, prop. 2.24)

constitute an [equivalence of groupoids](#) (def. 1.38

$$\text{Core}(\text{Cov}(X)) \begin{array}{c} \xleftarrow{\text{Rec}} \\ \xrightarrow{\text{Fib}} \end{array} \text{Rep}(\Pi_1(X), \text{Set})$$

between the groupoid $\text{Core}(\text{Cov}(X))$ (example 1.29, def. 2.1) whose [objects](#) are covering spaces over X , and whose [morphisms](#) are [isomorphisms](#) between these (def. 2.1) and the groupoid $\text{Rep}(\Pi_1(X), \text{Set})$ of [permutation groupoid representations](#) (def. 2.16) of the [fundamental groupoid](#) $\Pi_1(X)$ of X (example 1.27).

Proof. First we demonstrate a homotopy ([natural isomorphism](#)) of the form

$$\text{id}_{\text{Rep}(\Pi_1(X), \text{Set})} \xrightarrow{\simeq} \text{Fib} \circ \text{Rec} .$$

To this end, given $\rho \in \text{Rep}(\Pi_1(X), \text{Set})$ a [permutation groupoid representation](#), we need to exhibit in turn a homotopy ([natural isomorphism](#)) of permutation representations.

$$\eta_\rho : \rho \rightarrow \text{Fib}(\text{Rec}(\rho))$$

First consider what the right hand side is like: By def. [2.23](#) of [Rec](#) and def. [2.19](#) of [Fib](#) we have for every $x \in X$ an actual equality

$$\text{Fib}(\text{Rec}(\rho))(x) = \rho(x) .$$

To similarly understand the value of $\text{Fib}(\text{Rec}(\rho))$ on morphisms $[\gamma] \in \Pi_1(X)$, let $\gamma : [0, 1] \rightarrow X$ be a representing [path](#) in X . As in the proof of the path lifting lemma [2.8](#) we find a [finite number](#) of paths $\{\gamma_i\}_{i \in \{1, n\}}$ such that

1. regarded as morphisms $[\gamma_i]$ in $\Pi_1(X)$ they [compose](#) to $[\gamma]$:

$$[\gamma] = [\gamma_n] \circ \cdots \circ [\gamma_2] \circ [\gamma_1]$$

2. each γ_i factors through an open subset $U_i \subset X$ over which $\text{Rec}(\rho)$ trivializes.

Hence by [functoriality](#) of $\text{Fib}(\text{Rec}(\rho))$ it is sufficient to understand its value on these paths γ_i . But on these we have again by direct unwinding of the definitions that

$$\text{Fib}(\text{Rec}(\rho))([\gamma_i]) = \rho([\gamma_i]) .$$

This means that if we take

$$\eta_\rho(x) : \rho(x) \xrightarrow{=} \text{Fib}(\text{Rec}(\rho))$$

to be the above identification, then this is a homotopy/[natural isomorphism](#) as required.

It remains to see that these morphism η_ρ are themselves natural in ρ , hence that for each morphism $\phi : \rho \rightarrow \rho'$ the diagram

$$\begin{array}{ccc} \rho & \xrightarrow{\phi} & \rho' \\ \text{eta}_\rho \downarrow & & \downarrow \eta_{\rho'} \\ \text{Fib}(\text{Rec}(\rho)) & \xrightarrow[\text{Fib}(\text{Rec}(\phi))]{} & \text{Fib}(\text{Rec}(\rho')) \end{array}$$

commutes as a diagram in $\text{Rep}(\Pi_1(X), \text{Set})$. Since these morphisms are themselves groupoid homotopies (natural isomorphisms) this is the case precisely if for all $x \in X$ the corresponding component diagram commutes. But by the above this is

$$\begin{array}{ccc} \rho(x) & \xrightarrow{\phi(x)} & \rho'(x) \\ = \downarrow & & \downarrow = \\ \text{Fib}(\text{Rec}(\rho))(x) & \xrightarrow[\text{Fib}(\text{Rec}(\phi))(x)]{} & \text{Fib}(\text{Rec}(\rho'))(x) \end{array}$$

and hence this means that the top and bottom horizontal morphism are in fact equal. Directz unwiinding of the definitions shows that this is indeed the case.

Now we demonstrate a homotopy ([natural isomorphism](#)) of the form

$$\mathrm{Rec} \circ \mathrm{Fib} \xrightarrow{\simeq} \mathrm{id}_{\mathrm{Core}(\mathrm{Cov}(X))} .$$

For $E \in \mathrm{Cov}(X)$ a covering space, we need to exhibit a natural isomorphism of covering spaces of the form

$$\epsilon_E : \mathrm{Rec}(\mathrm{Fib}(E)) \rightarrow E .$$

Again by def. 2.23 of Rec and def. 2.19 of Fib the underlying set of $\mathrm{Rec}(\mathrm{Fib}(E))$ is actually equal to that of E , hence it is sufficient to check that this [identity function](#) on underlying sets is a [homeomorphism](#) of [topological spaces](#).

By the assumption that X is [locally path-connected](#) and [semi-locally simply connected](#), it is sufficient to check for $U \subset X$ an open path-connected subset and $x \in X$ a point with the property that $\pi_1(U, x) \rightarrow \pi_1(X, x)$ lands is constant on the trivial element, that the open subsets of E of the form $U \times \{\hat{x}\} \subset p^{-1}(U)$ form a basis for the topology of $\mathrm{Rec}(\mathrm{Fib}(E))$. But this is the case by definition of Rec .

This proves the equivalence. ■

Example 2.26. ([universal covering space](#) reconstructed from [free](#) and [transitive fundamental group representation](#))

Let X be a [topological space](#) which is

- [path-connected](#),
- [locally path-connected](#)
- and [semi-locally simply-connected](#)

Then a [universal covering space](#) of X (def. \ref{CoveringUniversal}) exists.

Proof. By example 2.22 the covering space is connected and simply connected precisely if its [monodromy](#) representation is free and transitive. By the [fundamental theorem of covering spaces](#) (prop. 2.25) every [permutation representation](#) of the [fundamental group](#) $\pi_1(X)$ arises as the monodromy of some covering space. Hence it remains to see that a free and transitive representation of $\pi_1(X)$ exists: The action of any group G on itself, by left multiplication, is free and transitive. ■

3. Examples

We now use the theorems established above to compute the [fundamental groups](#) of topological spaces in some basic examples. In particular we prove the archetypical example saying that the [fundamental group of the circle is the integers](#) (prop. 3.1 below).

Fundamental groups

Proposition 3.1. (fundamental group of the circle is the integers)

The fundamental group π_1 of the circle S^1 is the additive group of integers:

$$\pi_1(S^1) \xrightarrow{\cong} \mathbb{Z}$$

and the isomorphism is given by assigning winding number.

Proof. The universal covering space $\widehat{S^1}$ of S^1 is the real line (by this example):

$$p := (\cos(2\pi(-)), \sin(2\pi(-))) : \mathbb{R}^1 \rightarrow S^1 .$$

Since the circle is locally path-connected (this example) and semi-locally simply connected (this example) the fundamental theorem of covering spaces applies and gives that the automorphism group of \mathbb{R}^1 over S^1 equals the automorphism group of its monodromy permutation representation:

$$\mathrm{Aut}_{\mathrm{Cov}(S^1)}(\mathbb{R}^1) \simeq \mathrm{Aut}_{\pi_1(S^1)\mathrm{Set}}(\mathrm{Fib}_{S^1}) .$$

Moreover, as a corollary of the fundamental theorem of covering spaces we have that the monodromy representation of a universal covering space is given by the action of the fundamental group $\pi_1(S)$ on itself (this prop.).

But the automorphism group of any group regarded as an action on itself by left multiplication is canonically isomorphic to that group itself (by this example), hence we have

$$\mathrm{Aut}_{\pi_1(S^1)\mathrm{Set}}(\mathrm{Fib}_{S^1}) \simeq \mathrm{Aut}_{\pi_1(S^1)\mathrm{Set}}(\pi_1(S^1)) \simeq \pi_1(S^1) .$$


Therefore to conclude the proof it is now sufficient to show that

$$\mathrm{Aut}_{\mathrm{Cov}(S^1)}(\mathbb{R}^1) \simeq \mathbb{Z} .$$

To that end, consider a homeomorphism of the form

$$\begin{array}{ccc} \mathbb{R}^1 & \xrightarrow[\simeq]{f} & \mathbb{R}^1 \\ p \searrow & & \swarrow p \\ & S^1 & \end{array} .$$

Let $s \in S^1$ be any point, and consider the restriction of f to the fibers over the complement:



$$\begin{array}{ccc}
 p^{-1}(S^1 \setminus \{s\}) & \xrightarrow[\simeq]{f} & p^{-1}(S^1 \setminus \{s\}) \\
 p \searrow & & \swarrow p \\
 & S^1 \setminus \{s\} &
 \end{array} .$$

By the [covering space](#) property we have (via [this example](#)) a [homeomorphism](#)

$$p^{-1}(S^1 \setminus \{s\}) \simeq (0, 1) \times \text{Disc}(\mathbb{Z}) .$$

Therefore, up to homeomorphism, the restricted function is of the form

$$\begin{array}{ccc}
 (0, 1) \times \text{Disc}(\mathbb{Z}) & \xrightarrow[\simeq]{f} & (0, 1) \times \text{Disc}(\mathbb{Z}) \\
 \text{pr}_1 \searrow & & \swarrow \text{pr}_1 \\
 & (0, 1) &
 \end{array} .$$

By the [universal property](#) of the [product topological space](#) this means that f is equivalently given by its two components

$$(0, 1) \times \text{Disc}(\mathbb{Z}) \xrightarrow{\text{pr}_1 \circ f} (0, 1) \quad (0, 1) \times \text{Disc}(\mathbb{Z}) \xrightarrow{\text{pr}_2 \circ f} \text{Disc}(\mathbb{Z}) .$$

By the [commutativity](#) of the above [diagram](#), the first component is fixed to be pr_1 . Moreover, by the fact that $\text{Disc}(\mathbb{Z})$ is a [discrete space](#) it follows that the second component is a [locally constant function](#) (by [this example](#)). Therefore, since the [product space](#) with a [discrete space](#) is a [disjoint union space](#) (via [this example](#))

$$(0, 1) \times \text{Disc}(\mathbb{Z}) \simeq \bigsqcup_{n \in \mathbb{Z}} (0, 1)$$

and since the disjoint summands $(0, 1)$ are [connected topological spaces](#) ([this example](#)), it follows that the second component is a [constant function](#) on each of these summands (by [this example](#)).

Finally, since every function out of a [discrete topological space](#) is continuous, it follows in conclusion that the restriction of f to the fibers over $S^1 \setminus \{s\}$ is entirely encoded in an [endofunction](#) of the set of [integers](#)

$$\phi : \mathbb{Z} \rightarrow \mathbb{Z}$$

by

$$\begin{array}{ccc}
 S^1 \setminus \{s\} \times \text{Disc}(\mathbb{Z}) & \xrightarrow{f} & S^1 \setminus \{s\} \times \text{Disc}(\mathbb{Z}) \\
 (t, k) & \mapsto & (t, \phi(k))
 \end{array} .$$

Now let $s' \in S^1$ be another point, distinct from s . The same analysis as above applies now to the restriction of f to $S^1 \setminus \{s'\}$ and yields a function

$$\phi' : \mathbb{Z} \rightarrow \mathbb{Z} .$$

Since

$$\{p^{-1}(S^1 \setminus \{s\}) \subset \mathbb{R}^1, p^{-1}(S^1 \setminus \{s'\}) \subset \mathbb{R}^1\}$$

is an [open cover](#) of \mathbb{R}^1 , it follows that f is uniquely fixed by its restrictions to these two subsets.

Now unwinding the definition of p shows that the condition that the two restrictions coincide on the intersection $S^1 \setminus \{s, s'\}$ implies that there is $n \in \mathbb{Z}$ such that $\phi(k) = k + n$ and $\phi'(k) = k + n$.

This shows that $\text{Aut}_{\text{Cov}(S^1)}(\mathbb{R}^1) \simeq \mathbb{Z}$. ■

Example 3.2. ([isomorphism classes](#) of [coverings](#) of the circle are [conjugacy classes](#) in the [symmetric group](#))

The [monodromy](#) construction assigns to an [isomorphism class](#) of covering spaces over the [circle](#) S^1 with [fibers](#) consisting of n elements [conjugacy classes](#) of elements the [symmetric group](#) $\Sigma(n)$:

$$\left\{ \begin{array}{l} \text{isomorphism classes of} \\ \text{finite covering spaces} \\ \text{over the circle} \end{array} \right\} \simeq \left\{ \begin{array}{l} \text{conjugacy classes of} \\ \text{elements of a symmetric group} \end{array} \right\}$$

To see this we may without restriction choose a basepoint $x \in S^1$ so that a monodromy representation is equivalently a groupoid morphism of the form (prop. 2.18)

$$\rho : B\mathbb{Z} \xrightarrow{\simeq} B\pi_1(S^1, x) \xrightarrow{\rho} \text{Core}(\text{Set}) .$$

Since \mathbb{Z} is the [free abelian group](#) on a single generator, such as morphism is uniquely determined by the image of $1 \in \mathbb{Z}$. This is taken to some isomorphism of the set $p^{-1}(x)$. If we choose any identification $\phi : p^{-1}(x) \xrightarrow{\simeq} \{1, \dots, n\}$, then this defines an element $\sigma \in \Sigma(n)$ in the [symmetric group](#):

$$\begin{array}{ccccc} x & \mapsto & p^{-1}(x) & \xrightarrow[\simeq]{\phi} & \{1, \dots, n\} \\ 1 \downarrow & & \rho(1) \downarrow & & \downarrow \sigma \\ x & \mapsto & p^{-1}(x) & \xrightarrow[\phi]{\simeq} & \{1, \dots, n\} \end{array} .$$

Now if

$$f : E_1 \xrightarrow{\simeq} E_2$$

is an isomorphism of covering spaces, then by the [fundamental theorem of covering spaces](#) (prop. 2.25) this corresponds bijectively to a homomorphism of representations

$$\text{Fib}(f) : \text{Fib}_{E_1} \xrightarrow{\simeq} \text{Fib}_{E_2}$$

which in turn is by definition a homotopy (natural isomorphism) between the monodromy functors $\text{Fib}_{E_i} : B\mathbb{Z} \rightarrow \text{Core}(\text{Set})$.

The combination of the naturality square of this natural isomorphism with the above identification yields the following diagram

$$\begin{array}{ccccccc}
 \{1, \dots, n\} & \xrightarrow{\phi_1^{-1}} & p_1^{-1}(x) & \xrightarrow{f|_{\{x\}}} & p_2^{-1}(x) & \xrightarrow{\phi_2} & \{1, \dots, n\} \\
 \sigma_1 \downarrow & & \text{Fib}_{E_1}(1) \downarrow & & \downarrow \text{Fib}_{E_2}(1) & & \downarrow \sigma_2 \\
 \{1, \dots, n\} & \xrightarrow{\phi_1^{-1}} & p_1^{-1}(x) & \xrightarrow{f|_{\{x\}}} & p_2^{-1}(x) & \xrightarrow{\phi_2} & \{1, \dots, n\}
 \end{array}$$

The commutativity of the total rectangle says that the permutations σ_1 and σ_2 are related by conjugation with the element $\phi_2 \circ f|_{\{x\}} \circ \phi_1^{-1}$.

Example 3.3. (three-sheeted covers of the circle)

Consider the three-sheeted [covering spaces](#) of the [circle](#).

By [example 3.2](#) these are, up to isomorphism, given by the [conjugacy classes](#) of the elements of the [symmetric group](#) $\Sigma(3)$ on three elements. These in turn are labeled by the [cycle](#) structure of the elements ([this prop.](#)).

For the symmetric group on three elements there are three such classes

$$\begin{aligned}
 (1\ 2\ 3) \\
 (1\ 2)(3) \\
 (1)(2)(3)
 \end{aligned}$$

The corresponding covering spaces of the circle are shown in the graphics.

graphics grabbed from [Hatcher](#)

Higher homotopy groups

(...)



This concludes the introduction to basic homotopy theory.

For introduction to more general and abstract homotopy theory see at [Introduction to Homotopy Theory](#).

An incarnation of [homotopy theory](#) in [linear algebra](#) is [homological algebra](#). For introduction to that see at [Introduction to Homological Algebra](#).

4. References

A textbook account is in

- [Tammo tom Dieck](#), sections 2 and 3 of *Algebraic Topology*, EMS 2006 ([pdf](#))

Lecture notes include

- [Jesper Møller](#), *The fundamental group and covering spaces* (2011) ([pdf](#))

Revised on July 19, 2017 16:25:30 by [Urs Schreiber](#)