

Homotopy axiomatic cohesion

Some basic ideas

Urs Schreiber

December 6, 2011

Special thanks to

- ▶ Mike Shulman
- ▶ Dave Carchedi
- ▶ Richard Williamson

Details and references at

<http://ncatlab.org/schreiber/show/differential+cohomology+in+a+cohesive+topos>

Outline

Geometry

Internal language

Homotopy theory

Cohesive ∞ -toposes

Goal:

Pair

geometry + homotopy
theory

axiomatically.

Motivation:

Gauge theory in physics:

- ▶ fields are smooth functions on spacetime;
- ▶ gauge transformations are “smooth homotopies”;
- ▶ gauge-of-gauge transformation are “higher smooth homotopies”.

Axiomatization unifies varying details,
such as replacing smooth geometry
by supergeometry.

I

Geometry: Cohesive toposes

In geometry, toposes play
two different roles

1. as generalized topological spaces;
2. as *collections* of geometric spaces.

Axiomatic cohesion characterizes
toposes \mathbf{H} of the second kind.
(Lawvere)

First axiom – \mathbf{H} is local:

There exist adjoint functors

$$\mathbf{H} \begin{array}{c} \xleftarrow{\text{Disc}} \\ \xrightarrow{\Gamma} \\ \xleftarrow{\text{coDisc}} \end{array} \text{Set} ,$$

Interpretation:

$$\mathbf{H} \begin{array}{c} \xleftarrow{\text{discrete cohesive structure}} \\ \xrightarrow{\text{underlying point set}} \\ \xleftarrow{\text{codiscrete cohesive structure}} \end{array} \text{Set}$$

Second axiom – \mathbf{H} is strongly connected:

$$\begin{array}{ccc} & \xrightarrow{\exists \Pi} & \\ \mathbf{H} & \begin{array}{c} \xleftarrow{\text{Disc}} \\ \xrightarrow{\Gamma} \\ \xleftarrow{\text{coDisc}} \end{array} & \text{Set} \end{array},$$

Π preserves finite products.

Interpretation:

$$\mathbf{H}(X, \text{Disc}S) \simeq \text{Set}(\Pi X, S)$$

says: Π sends any object to its set of *connected components*.

Consequence

The space $* \in \mathbf{H}$ underlying \mathbf{H} looks like a fat point:

- ▶ strong connectedness: $\Pi_* = *$;
- ▶ locality: if $X \rightarrow *$ is epi, then it has a section.

So $* \in \mathbf{H}$

- ▶ is connected
- ▶ “looks contractible”;

Consequence

Since the generalized space
underlying $X \in \mathbf{H}$
is the slice topos

$$\mathbf{H}/X \xrightarrow{\text{étale}} \mathbf{H}/_*$$

every $X \in \mathbf{H}$ is a space *locally modeled* on the fat point $\mathbf{H}/_*$:
the *abstract cohesive blob*.

Examples

- ▶ $\mathbf{H} = \text{Sh}(\text{TopologicalManifolds})$;
- ▶ $\mathbf{H} = \text{Sh}(\text{SmoothManifolds})$,
contains generalized smooth
spaces such as *diffeological spaces*.
- ▶ $\mathbf{H} = \text{CahierTopos}$,
a model for synthetic differential
geometry;

II

Internal language

We may reflect cohesion structure back into \mathbf{H} .

$$(\Pi \dashv \flat \dashv \sharp) : \mathbf{H} \begin{array}{c} \xrightarrow{\Pi} \\ \xleftarrow{\text{Disc}} \\ \xrightarrow{\Gamma} \end{array} \text{Set} \begin{array}{c} \xrightarrow{\text{Disc}} \\ \xleftarrow{\Gamma} \\ \xrightarrow{\text{coDisc}} \end{array} \mathbf{H}$$

and express these endofunctors in the *internal language* of \mathbf{H} .

Internal Type Theory language of a topos \mathbf{H}

Dictionary

type	$\vdash X : \text{Type}$	object $X \in \mathbf{H}$
term	$\vdash x : X$	morphism $x : * \rightarrow X$
dependent type	$x : X \vdash A(x) : \text{Type}$	morphism $A \rightarrow X$
type of propositions	$\vdash \text{Prop} : \text{Type}$	subobject classifier Ω
proposition	$x : X \vdash \phi(x) : \text{Prop}$	monomorphism $A \hookrightarrow X$ \simeq morphism $\phi : X \rightarrow \Omega$
dependent sum	$\vdash \exists x, A(x)$	$A \in \mathbf{H}$
dependent product	$\vdash \forall x, A(x)$	object of sections $\Gamma_X(A)$

In this internal logic,
 cohesion incarnates as *modalities* (qualified truth).
 Notably \sharp incarnates as a *geometric modality*:

$$\sharp : \text{Prop} \rightarrow \text{Prop}$$

defined by

$$\sharp : \Omega \xrightarrow{\text{unit}} \sharp\Omega \xrightarrow{\chi_{\sharp\text{true}}} \Omega .$$

Sends subobjects $\phi : A \hookrightarrow X$ to the pullback

$$\begin{array}{ccc} & \longrightarrow & \sharp A \\ \downarrow \sharp\phi & & \downarrow \\ X & \longrightarrow & \sharp X \end{array} .$$

Say $\phi(x : X) : \text{Prop}$ is *discretely true*
(true over a discrete space)
if $\sharp\phi$ is true.

For instance in $\mathbf{H} = \text{Sh}(\text{SmthMfd})$

$\text{isClosed}(\omega : \text{Differential}n\text{Form})$

is discretely true.

$$\begin{array}{ccc} \Omega_{\text{cl}}^n(-) & & \Omega^n(-) \\ \downarrow & \xrightarrow{\sharp} & \downarrow \text{id} \\ \Omega^n(-) & & \Omega^n(-) \end{array}$$

But:

Internal characterization of cohesion
requires to lift modalities from
propositions to all types

$$\sharp : \text{Type} \rightarrow \text{Type}$$

Needed to characterize reflection:

$$\forall X Y : \text{Type}, \text{isCodiscrete}(Y) :$$

$$[\sharp X, Y] \xrightarrow{\sim} [X \rightarrow Y].$$

III

Homotopy (type) theory

Identity types. Let $\text{Id}_X(x, y)$ be the type of *proofs* that terms x and y are equal.

Now drop the assumption that there are unique such terms:

there may be different proofs of the same fact, different *paths* from x to y .

Consider paths of paths of paths...

$$\gamma, \rho : \text{Id}_{\text{Id}_X(x, y)}(\alpha, \beta : \text{Id}_X(x, y))$$

“ ∞ -groupoid” : $X \simeq \left\{ \begin{array}{c} \alpha \\ \begin{array}{ccc} x & \begin{array}{c} \curvearrowright \gamma \\ \curvearrowleft \rho \end{array} & y \\ \beta \end{array} \end{array} \right\}$

This now has an interpretation (Awodey, Warren) in **H** a *model topos* (Rezk, Lurie):

- ▶ class of “weak” equivalences $\xrightarrow{\simeq}$;
- ▶ class of “bundle morphisms” $\longrightarrow =$ fibrations;
- ▶ compatible sSet-enrichment.

Refined dictionary:

type	$\vdash X : \text{Type}$	fibrant object $A \longrightarrow *$
dependent type	$x : X \vdash A(x) : \text{Type}$	fibration $A \longrightarrow X$
identity type	$x \ y : X \vdash \text{Id}_X(x, y) : \text{Type}$	path object $X^{\Delta[1]} \xrightarrow{(s,t)} X \times X$
type of small types	Type	<i>small object</i> classifier

IV

Synthesis:

Cohesive ∞ -toposes

Using this, the relevant structures on a topos do
have internal axiomatization:

- ▶ locality;
- ▶ connectivity;
- ▶ cohesion.

Mike Shulman, *Internalizing the external* (2011)

The resulting structure is automatically
homotopy cohesion.

Example:

model topos

$$\mathbf{H} := \text{PSh}(\text{SmothMfd}, \text{sSet})$$

with

- ▶ weak equivalences are the *stalkwise weak homotopy equivalences*;

Context for higher differential geometry:

$$\mathbf{H} = \{ \text{smooth } \infty\text{-groupoids} \}$$

The extra left adjoint

$$\Pi : \mathbf{H} \rightarrow \mathbf{sSet}$$

now sends a manifold X to its *path* ∞ -groupoid

$$\Pi X \simeq \left\{ \begin{array}{c} \begin{array}{ccc} & \gamma & \\ \swarrow & \text{---} & \searrow \\ x & & y \\ \nwarrow & \text{---} & \nearrow \\ & \tilde{\gamma} & \end{array} \\ \Sigma \quad \tilde{\Sigma} \end{array} \right\}$$

that consists of actual *geometric paths*, *geometric surfaces*, etc. in X .

Outlook:

Intrinsic notion of
geometric paths
induces
intrinsic notion of “dynamics”:
differential cohomology.

End.

Details and further material at

<http://ncatlab.org/schreiber/show/differential+cohomology+in+a+cohesive+topos>