



Faculty of Science

Simplicial Complexes

A short Introduction to Algebraic Topology and Discrete
Geometry

Kenny Erleben
kenny@diku.dk

Department of Computer Science
University of Copenhagen

2010



The Definition of a Simplex

A simplex is defined as the point set consisting of the convex hull of a set of linear independent points.

- Let $\{v_i\}^{n+1}$ denote a linear independent point set containing $n + 1$ points. Henceforth named the vertex set and its elements the vertices. The simplex, σ_n , is defined as the point set,

$$\sigma_n \equiv \left\{ v \mid v = \sum_{i=1}^{n+1} \lambda_i v_i, \quad \sum_{i=1}^{n+1} \lambda_i = 1, \quad 0 \leq \lambda_i \leq 1 \quad \forall i \right\}$$

- In three dimensional Euclidean space we can have up to four linear independent vertices. This implies that $n = 0, 1, 2, 3$ are the only possible choices in a three dimensional space.



The Vertex Set Operation

Let $\{v_i\}^{n+1}$ denote a linear independent point set containing $n + 1$ points and defining the point set of the simplex, σ_n .

$$\mathbf{vert}(\sigma_n) \equiv \{v_i\}^{n+1} = \{v_1, v_2, \dots, v_{n+1}\}$$

As short-hand notation we use the labeling $\sigma_n \equiv \{v_1, v_2, \dots, v_{n+1}\}$ as the notation that defines the simplex.



Simplex Dimension

The number of linear independent vertex basis vectors of the point-set of the simplex will be denoted as the dimension of the simplex. Thus we have

$$\mathbf{dim}(\sigma_n) \equiv n$$

for $n = 0, 1, 2, 3, 4$ and so on.



The Orientation of a Simplex

The orientation of a simplex is given by the ordering of the vertex set up to an even permutation (even number of two element swaps)

- Thus, there exist only two classes of orientations

Example: given $\sigma_2 = \{v_1, v_2, v_3\}$ then

- $\{v_1, v_2, v_3\}$, $\{v_2, v_3, v_1\}$, and $\{v_3, v_1, v_2\}$ are of same orientation
- $\{v_2, v_1, v_3\}$, $\{v_1, v_3, v_2\}$, and $\{v_3, v_2, v_1\}$ are of same orientation

but $\{v_1, v_2, v_3\}$ and $\{v_2, v_1, v_3\}$ are of different orientations.



More on Orientations

- A zero-simplex (a single vertex) has no orientation
- The two different orientations are often designated by a sign, $+1$ or -1 .
- One convention for picking an orientation is to use the determinant of the vertex basis,

$$\mathbf{sgn}(\sigma_n) \equiv \mathbf{sgn}(\det([(v_2 - v_1) \quad (v_3 - v_1) \quad \cdots \quad (v_{n+1} - v_1)]))$$

- If we are given the orientation $\mathbf{sgn}(\sigma_1) = \mathbf{sgn}(\{v_1, v_2\})$ then the opposite orientation is written as $\mathbf{sgn}(\{v_2, v_1\})$
- Or even more shorthand we use $\{v_1, v_2\} = -\{v_2, v_1\}$



Examples of Notation

So we have



$$\mathbf{sgn}(\{v_1, v_2, v_3\}) = \mathbf{sgn}(\mathbf{det}([v_2 - v_1 \quad v_3 - v_1]))$$



$$\mathbf{sgn}(\{v_1, v_2, v_3\}) = \mathbf{sgn}(\{v_2, v_3, v_1\}) = \mathbf{sgn}(\{v_3, v_1, v_2\})$$



$$\mathbf{sgn}(\{v_2, v_1, v_3\}) = \mathbf{sgn}(\{v_1, v_3, v_2\}) = \mathbf{sgn}(\{v_3, v_2, v_1\})$$



$$\mathbf{sgn}(\{v_1, v_2, v_3\}) = -\mathbf{sgn}(\{v_2, v_1, v_3\})$$



A Simplex Face (Sub-simplex)

A face σ_m of a simplex σ_n is a simplex spanned by the subset of vertices of $\{v_i\}^{n+1}$

$$\mathbf{vert}(\sigma_m) \subseteq \mathbf{vert}(\sigma_n)$$

Observe

- Any face is itself a simplex
- By definition of a face any simplex is a face of itself.

If $\mathbf{dim}(\sigma_m) < \mathbf{dim}(\sigma_n)$ we call σ_m a proper face of σ_n .



The Boundary of a Simplex

We will define the boundary, Γ , of a simplex, σ to denote the set of faces having fewer elements in the vertex set than σ .

$$\Gamma(\sigma_n) \equiv \{\sigma_m \mid m < n \wedge \mathbf{vert}(\sigma_m) \subseteq \mathbf{vert}(\sigma_n)\}$$

Thus for $\sigma_2 = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ (a triangle), the boundary is by our definition

$$\Gamma(\sigma_2) = \{\{\mathbf{v}_1, \mathbf{v}_2\}, \{\mathbf{v}_1, \mathbf{v}_3\}, \{\mathbf{v}_2, \mathbf{v}_3\}, \{\mathbf{v}_1\}, \{\mathbf{v}_2\}, \{\mathbf{v}_3\}\}$$

Thus, it is merely the edges and the vertices of the triangle.



The Boundary Operator

A slightly different definition defines the boundary operator of σ_n to be all faces having exactly $n - 1$ elements in their vertex sets.

$$\partial(\sigma_n) \equiv \{\sigma_m \mid m = n - 1 \quad \wedge \quad \mathbf{vert}(\sigma_m) \subseteq \mathbf{vert}(\sigma_n)\}$$

Usually the orientations of the faces must be handled carefully.

- The boundary operator yields a set of $n + 1$ simplexes

$$\partial(\sigma_n) = \{\sigma_m^j\}_{j=1}^{n+1} \text{ where}$$


$$\sigma_m^j = (-1)^{j+1} \{v_1, \dots, \hat{v}_j, \dots, v_{n+1}\}$$

and \hat{v}_j means that v_j is dropped.

Observe that from a “geometric point set” viewpoint $\partial(\sigma_n) \equiv \Gamma(\sigma_n)$, only “topological set-wise” $\partial(\sigma_n) \neq \Gamma(\sigma_n)$.



Self Training 1



$$\text{span}(\{1, 2, 3\}) = 2$$

$$\text{span}(\{1, 2, 5\}) = 2$$

$$\rho(\{1, 2, 3\}) = 2$$

$$\rho(\{1, 2, 5\}) = 2$$

$$\rho(\{1, 2\}) = ?$$

$$\rho(\{4\}) = ?$$



The Closure of a Simplex

The closure operation is the union of the simplex and its boundary.
Here is a combinatorial notion of closure

$$\mathbf{cl}(\sigma_n) \equiv \sigma_n \cup \Gamma(\sigma_n)$$

Thus for $\sigma_2 = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ we have

$$\mathbf{cl}(\sigma_2) = \{\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}, \{\mathbf{v}_1, \mathbf{v}_2\}, \{\mathbf{v}_1, \mathbf{v}_3\}, \{\mathbf{v}_2, \mathbf{v}_3\}, \{\mathbf{v}_1\}, \{\mathbf{v}_2\}, \{\mathbf{v}_3\}\}$$

From a point set viewpoint one could just as easily have used the boundary operator $\partial(\sigma_n)$ in place of $\Gamma(\sigma_n)$ in the above definition.




The Closure of a Set of Simplexes

Given a set of simplexes $\mathcal{K} = \{\sigma^1, \dots, \sigma^N\}$ the closure of \mathcal{K} is defined as

$$\mathbf{cl}(\mathcal{K}) \equiv \bigcup_{\sigma^k \in \mathcal{K}} \mathbf{cl}(\sigma^k)$$



Self Training 2



$$K = \{ \{1, 2, 3\}, \{1, 5, 2\} \}$$

$$cl(K) = ?$$



The Interior of a Simplex

We define the interior of a simplex as the point set of the simplex minus the points on the boundary.

$$\mathbf{int}(\sigma_n) \equiv \mathbf{cl}(\sigma_n) \setminus \Gamma(\sigma_n)$$

- Observe that all point sets are closed. Thus the vertices of a triangle are contained in the edges of the triangle and both the vertices and the edges of the triangle are contained in the triangle.

Observe from a point-set viewpoint we have

$$\mathbf{int}(\sigma_n) = \mathbf{cl}(\sigma_n) \setminus \partial(\sigma_n).$$



Adjacent Simplexes

Two simplexes σ^i and σ^j are said to be adjacent if and only if

- $\mathbf{dim}(\sigma^i) = \mathbf{dim}(\sigma^j)$
- and they share a common face

$$\sigma^k = \sigma^i \cap \sigma^j \neq \emptyset$$

- and the dimension of the common face is exactly one lower than the dimension of the simplexes


$$\mathbf{dim}(\sigma^k) = n - 1$$

where $n = \mathbf{dim}(\sigma^i) = \mathbf{dim}(\sigma^j)$

We define the boolean binary relation $\mathbf{adj}(\sigma^i, \sigma^k)$ to be true if and only if σ^i and σ^j are adjacent simplexes and false otherwise.



Self Training 3



$$\text{adj}(\{1, 2, 3\}, \{1, 5, 2\}) = ?$$

$$\text{adj}(\{1, 2\}, \{2, 3\}) = ?$$

$$\text{adj}(\{1, 2, 3\}, \{3, 4\}) = ?$$

$$\text{adj}(\{3, 4\}, \{1, 2\}) = ?$$

$$\text{adj}(\{1\}, \{2\}) = ?$$



The Simplicial Complex

A simplicial complex is a finite collection \mathcal{K} of simplexes and the following two properties are always true


- Every face $\sigma^k \subset \sigma^j$ of each simplex $\sigma^j \in \mathcal{K}$ is also a simplex in \mathcal{K}
- Any intersection of two simplexes σ^i and σ^j from \mathcal{K} is

$$\sigma^i \cap \sigma^j = \begin{cases} \emptyset \\ \sigma^k \in \mathcal{K} \end{cases}$$



Self Training 4

For each of simplex collections below determine which are simplicial complexes



$$K_1 = \{ \{1,2,3\}, \{1,5,2\}, \{1,2\}, \{2,3\}, \{3,1\}, \{2,5\}, \{5,1\}, \\ \{1\}, \{2\}, \{3\}, \{5\} \}$$

$$K_2 = K_1 \cup \{3,4\}$$

$$K_3 = K_1 \cup \{6\}$$

$$K_4 = \{ \{7,8,9\}, \{10,11,12\} \} \cup \{ \{7,8,9\} \} \cup \{ \{10,11,12\} \}$$

$$K_5 = \text{cl}(K_2)$$



The Star (one-ring) of a Simplex

Given $\sigma \in \mathcal{K}$ then the star operator is given by

$$\mathbf{star}(\sigma) \equiv \{\sigma_n \mid \sigma_n \in \mathcal{K} \wedge \mathbf{vert}(\sigma) \subset \mathbf{vert}(\sigma_n)\}$$


That is the set of all simplexes that σ is a face of.

- A top-simplex is defined as having $\mathbf{star}(\sigma) = \sigma$
- The dimension of a simplicial complex is equal to the highest dimension top simplex in the simplicial complex

The star operator is sometimes called the co-boundary operator.



Self Training 5



$$K = \{ \{1,2,3\}, \{1,4,2\}, \{2,5\} \}$$

$$\text{star}(\{2\}) = ?$$

$$\text{star}(\{1,2\}) = ?$$



The Discrete Manifold

An n -dimensional discrete manifold is an n -dimensional simplicial complex that satisfies


- For each simplex the union of all n -dimensional incident n -simplexes forms an n -dimensional ball
- or a half-ball if the simplex is on the boundary

Thus, each $n - 1$ -dimensional simplex has exactly two adjacent n -dimensional simplexes if not on the boundary and exactly one n -dimensional simplex otherwise.




Self Training 6


Determine which examples are discrete manifolds and which are not



$$K_1 = \text{cl}(\{\{1,2\}, \{2,3\}, \{2,4\}\})$$



$$K_2 = \text{cl}(\{\{5,6,7\}, \{5,7,8\}, \{7,9\}\})$$



$$K_3 = \text{cl}(\{\{10,11,12\}, \{11,13,12\}, \{12,13,14\}\})$$



That is It!

Questions?



Further Reading

- Siggraph Asia 2008 course notes: Discrete Differential Geometry: An applied Introduction. (Read Chapters 7 and 8)
- Marek Krzysztof Misztal, Deformable Simplicial Complexes, PhD Thesis, IMM, DTU, 2010




Study Group

- Do the training exercises and discuss the definitions of each operation
- If you have time do the “Extras For Self-Study ” listed on a later slide



Answers for Self Training 1



$$\text{sgn}(\{1, 2, 3\}) = +1$$

$$\text{sgn}(\{1, 2, 5\}) = -1$$

$$d(\{1, 2, 3\}) = \{ \{2, 3\}, \{3, 1\}, \{1, 2\} \}$$


$$d(\{1, 2, 5\}) = \{ \{2, 5\}, \{5, 1\}, \{1, 2\} \}$$

$$d(\{1, 2\}) = \{ \{1\}, \{2\} \}$$

$$d(\{4\}) = \emptyset$$



Answers for Self Training 2




$$K = \{ \{1, 2, 3\}, \{1, 5, 2\} \}$$

$$cl(K) = \{ \{1, 2, 3\}, \{1, 5, 2\}, \{1, 2\}, \{2, 3\}, \{3, 1\}, \{5, 2\}, \{1, 5\}, \{1, 3\}, \{2, 3\}, \{3\}, \{5\} \}$$



Answers for Self Training 3



$$\text{adj}(\{1, 2, 3\}, \{1, 5, 2\}) = \text{TRUE}$$

$$\text{adj}(\{1, 2\}, \{2, 3\}) = \text{TRUE}$$


$$\text{adj}(\{1, 2, 3\}, \{3, 4\}) = \text{FALSE}$$

$$\text{adj}(\{3, 4\}, \{1, 2\}) = \text{FALSE}$$

$$\text{adj}(\{1\}, \{2\}) = \text{FALSE}$$



Answers for Self Training 4



K_1 is a simplicial complex

K_2 is not a simplicial complex


K_3 is a simplicial complex

K_4 is not a simplicial complex

K_5 is a simplicial complex



Answers for Self Training 5



$$K_1 = \text{cl}(\{ \{1, 2, 3\}, \{1, 4, 2\}, \{2, 5\} \})$$

$$\text{star}(\{2\}) = \{ \{2, 5\}, \{2, 3\}, \{2, 1\}, \{2, 4\}, \\ \{1, 2, 3\}, \{1, 4, 2\} \}$$

$$\text{star}(\{1, 2\}) = \{ \{1, 2, 3\}, \{1, 4, 2\} \}$$



Answers for Self Training 6



K_1 is not a discrete manifold



K_2 is not a discrete manifold



K_3 is 2D discrete manifold



Extras For Self-Study

- A Discrete Manifold is said to have consistent orientation if all top-simplexes has the same orientation
- The link of a simplex $\sigma \in \mathcal{K}$ from a simplicial complex \mathcal{K} is defined as

$$\mathbf{link}(\sigma) \equiv \mathbf{cl}(\mathbf{star}(\sigma)) \setminus \mathbf{star}(\mathbf{cl}(\sigma))$$

- Chains, Co-chains and Skeletons and much more...



What have We Learned?

- Geometry (=point-sets) and topology (= combinatorics) are two different things
- What we consider a nice mesh – the discrete manifold
- Star and link operators – are nice for making local changes
- Boundary and co-boundary operators are really useful for finite volume methods etc..

