

M-theory and cohomotopy

Hisham Sati

New York University Abu Dhabi (NYUAD)

M-Theory and Mathematics

NYUAD Workshop

27–30 January 2020

Outline

I. From 11d sugra to M-theory

II. Where do fields live?

III. (Twisted) Cohomotopy vs. (twisted) cohomology description of the M-theory fields

1. *Rationally.*

2. *Integrally.*

3. *Differentially.*

IV. Further applications: branes and gauge theory

Joint with: Urs Schreiber, Domenico Fiorenza, Dan Grady, Vincent Braunack-Mayer

[BMSS]= Braunack-Mayer-S.-Schreiber


[FSS]= Fiorenza-S.-Schreiber

[GS]= Grady-S.

[S]= S.

[SS]= S.-Schreiber

Richness of M-theory



I. From 11d supergravity to M-theory

Bosonic 11D supergravity

- **Bosonic Lagrangian:** given by the eleven-form [[Cremmer-Julia-Scherk](#)]

$$\mathcal{L}_{11}^{\text{bos}} = R * \mathbf{1} - \frac{1}{2} G_4 \wedge * G_4 - \frac{1}{6} G_4 \wedge G_4 \wedge C_3$$

- **Equations of motion:** The variation $\frac{\delta L_{(11),\text{bos}}}{\delta C_3} = 0$ for C_3 gives the corresponding equation of motion

$$d * G_4 + \frac{1}{2} G_4 \wedge G_4 = 0 . \quad (1)$$

- **Bianchi identity:**

$$dG_4 = 0 . \quad (2)$$

- The second order equation (1) can be written in a first order form, by first writing $d(*G_4 + \frac{1}{2} C_3 \wedge G_4) = 0$ so that

$$*G_4 = G_7 := dC_6 - \frac{1}{2} C_3 \wedge G_4 , \quad (3)$$

where C_6 is the potential of G_7 , the Hodge dual field strength to G_4 in 11 dimensions.

The effect of the fermions

- The fermionic field $\psi \in \Gamma(S \otimes TM)$ (the gravitino) satisfies the generalized Dirac equation, the Rarita-Schwinger equation

$$D_{RS}\psi = 0, \quad \psi \in \Gamma(S \otimes T^*M).$$

(involves mixing of terms).

- The fields themselves are in fact combinations of bosonic and fermionic fields. Physics literature usually writes:

$$G_4^{\text{super}} = \underbrace{G_4}_{\sim \text{topology/geometry}} + \underbrace{\bar{\psi}\Gamma_2\psi}_{\sim \text{topology/geometry}}$$

- Similarly for the connections

$$\omega^{\text{super}} = \omega + \text{fermion-bilinears}$$


[See [Duff-Nilsson-Pope](#)]

Strategy: Extract topology/higher geometry from bosons and fermions separately.

II. Where do fields live?

Generalities on what physics wants


Nontrivial physical entities, such as fields, charges, etc. generically take values in cohomology.




- i. **Generalized:** Capture essential topological and bundles aspects.
- ii. **Twisting:** Account for symmetries via automorphisms.
- iii. **Differentially refined:** Include geometric data, such as connections, Chern character form, smooth structure, smooth representatives of maps ...

Differential refinement

- Introduce geometric data via differential forms (connections, Chern forms, ...), i.e., retain differential form representatives of cohomology classes.



- Amalgam of an underlying (topological) cohomology theory and the data of differential forms:



- That is, we have a fiber product or twisted product

$$\text{“Differential cohomology} = \text{Cohomology} \times_{\text{de Rham}} \text{Forms”}$$

Differential generalized cohomology

- Start with a generalized cohomology theory h
- $\Omega(X, h_*) := \Omega(X) \otimes_{\mathbb{Z}} h_*$ Smooth differential forms with coefficients in $h_* := h(*)$
- $\Omega_{\text{cl}}(X, h_*) \subseteq \Omega(X, h_*)$ closed forms
- $H_{\text{dR}}(X, h_*)$ cohomology of the complex $(\Omega(X, h_*), d)$


Definition

A **smooth extension** of h is a contravariant functor

$\widehat{h} : \text{Compact Smooth Manifolds} \rightarrow \text{Graded Abelian Grps}$

$$\begin{array}{ccc} & & \Omega_{\text{cl}}(X, h_*) \\ & \nearrow R & \downarrow \\ \widehat{h}(X) & & H_{\text{dR}}(X, h_*) \\ & \searrow I & \uparrow \\ & & h(X) \end{array}$$

Twisted \cap Differential \cap Generalized



Examples ([GS])

- 1 Type I (II) RR fields live in twisted differential KO-theory $\widehat{KO}_{\hat{\tau}}$ (K-theory $\widehat{K}_{\hat{\tau}}$).
 - 2 Differential refinements of various twisted cohomology theories.
- Fields in M -theory are proposed to live in a theory of this type [S06]. Which one?

III. (Twisted) Cohomotopy vs. (twisted) cohomology description of the M-theory fields

Cohomotopy versus cohomology

- **Cohomology** of Y with R -coefficients: $[Y, K(R, n)] \cong H^n(Y; R)$. **old**
- **Cohomotopy** of Y with R -coefficients: $[Y, S_R^n] \cong \pi_R^n(Y)$. **new**

Compare cohomotopy to cohomology of various flavors:

- 1 Rational: $S_{\mathbb{Q}}^4$ vs. $H^4(-; \mathbb{Q})$.
- 2 Integral: $S_{\mathbb{Z}}^4$ vs. $H^4(-; \mathbb{Z})$.
- 3 Differential: \widehat{S}^4 vs. $\widehat{H}^4(-)$.

1. Rationally

Definition

The field equations of (a limit) of M-theory on an 11-dimensional manifold Y^{11} are

$$\begin{aligned}d * G_4 &= \frac{1}{2} G_4 \wedge G_4 \\dG_4 &= 0\end{aligned}$$

- **Q. What topological & geometric information can the above system provide us?**
 - Rational structures: Differential forms, rational cohomology, rational homotopy theory ...
 - More refined structures: (twisted) 2-gerbes, (twisted) String structures, orientations ...
- A priori, G_4 should be described by a map $f : Y \rightarrow K(\mathbb{Z}, 4) \rightsquigarrow H^4(Y; \mathbb{Z})$
- Differential refinement \widehat{G}_4 corresponds to $Y \rightarrow B^3U(1)_{\nabla} \rightsquigarrow \widehat{H}^4(Y)$
- Product structure on Eilenberg-MacLane spaces is *cup product*, with **no** a priori information about *trivialization*.
- Need (G_4, G_7) satisfying above $\Leftrightarrow Y \rightarrow ?$.
- Need $(\widehat{G}_4, \widehat{G}_7)$ satisfying above $\Leftrightarrow Y \rightarrow \widehat{?}$.

Rational degree four twists [S]

- Consider a 3-form C_3 with $G_4 = dC_3$. We can build a differential with G_4 as $d_{G_4} = d + v_3^{-1} G_4 \wedge$

Observation

The de Rham complex can be twisted by a differential of the form $d + v_{2i-1}^{-1} G_{2i} \wedge$ provided that G_{2i} is closed and v_{2i-1} is Grassmann algebra-valued.

- Form a duality-symmetric graded uniform degree form $G = v_3^{-1} G_4 + v_6^{-1} G_7$. This expression can now be used to twist the de Rham differential, leading to

$$d_G = d + G \wedge = d + v_3^{-1} G_4 \wedge + v_6^{-1} G_7 \wedge .$$

Observation

The de Rham complex can be twisted by the differential d_G provided

- $\{v_3, v_3\} = v_6$
- $dG_7 = \frac{1}{2} G_4 \wedge G_4$.

The first condition is the M-theory gauge algebra and the second is the equation of motion.

Observation (The Sullivan model as the equations of motion [S])

The above equations correspond to the Sullivan DGCA model of the 4-sphere S^4

$$\mathcal{M}(S^4) = (\wedge(y_4, y_7); dy_7 = y_4^2, dy_4 = 0)$$

What about the factor of $\frac{1}{2}$?

- Whitehead bracket $[\iota_4, \iota_4]_W : S^7 \rightarrow S^4$ generates \mathbb{Z} (\mathbb{Q})-summand in $\pi_7(S^4)$.
- There is an extra symmetry as we are in the dimension of a Hopf fibration, i.e. σ the \mathbb{H} -Hopf map and so the generator is $\sigma = \frac{1}{2}[\iota_4, \iota_4]_W$.

Observation (Quillen model as the M-theory gauge algebra [FSS])

The Sullivan model for S^{2n} is given by the DGCA

$$\mathcal{M}(S^{2n}) = (\wedge(x_{2n}, x_{4n-1}); dx_{2n} = 0, dx_{4n-1} = x_{2n}^2).$$

Imposing the Maurer-Cartan equation on the degree 1 element

$x_{2n}\xi_{1-2n} + x_{4n-1}\xi_{2-4n}$ we find the Lie bracket dual to the differential is given by

$$[\xi_{1-2n}, \xi_{1-2n}] = 2\xi_{2-4n}$$

with all the other brackets zero.

Example ($n = 2$)

The graded Lie algebra $\mathbb{R}\xi_{-3} \oplus \mathbb{R}\xi_{-6}$ with bracket $[\xi_{-3}, \xi_{-3}] = 2\xi_{-6}$ (Quillen model) can be identified with the M-theory gauge Lie algebra.


Proposal ([S])

Higher gauge fields in M-theory are cocycles in (rational) **cohomotopy**.

Developed via *Rational Homotopy Theory (RHT)* in [FSS]: $X \xrightarrow{(G_4, G_7)} S_{\mathbb{R}}^4$.

- $[Y, S_{\mathbb{Q}}^4] = \pi_{\mathbb{Q}}^4(Y)$ rational cohomotopy.
- Ultimately interested in full $\text{Map}(Y, S^4) \ni f$.
- Geometry + physics \Rightarrow **differential cohomotopy** [FSS]
- Formulate in stacks/chain complexes.

RHT. Generalized Chern character maps are examples of *rationalization*



RHT amenable to computations due to *Sullivan models*: differential graded-commutative algebras (dgc-algebras) on a finite number of generating elements (spanning the rational homotopy groups) subject to differential relations (enforcing the intended rational cohomology groups). In Suga: “FDA”s.

Examples

	Rational super space	Loop super L_∞ -algebra	Chevalley-Eilenberg super dgc-algebras ("Sullivan models", "FDA's")
General	X	$\mathfrak{l}X$	$CE(\mathfrak{l}X)$
Super spacetime	$\mathbb{T}^d, \mathbf{1} \mathbb{N}$	$\mathbb{R}^d, \mathbf{1} \mathbb{N}$	$\mathbb{R}[\{\psi^\alpha\}_{\alpha=1}^N, \{e^a\}_{a=0}^d] / \left(\begin{array}{l} d\psi^\alpha = \mathbf{0} \\ de^a = \frac{1}{\psi} \Gamma^a \psi \end{array} \right)$
Eilenberg-MacLane space	$K(\mathbb{R}, p+2)$ $\simeq_{\mathbb{R}} B^{p+1} S^1$	$\mathbb{R}[p+1]$	$\mathbb{R}[c_{p+2}] / (dc_{p+2} = \mathbf{0})$
Odd-dimensional sphere	$S_{\mathbb{R}}^{2k+1}$	$\mathfrak{l}(S^{2k+1})$	$\mathbb{R}[\omega_{2k+1}] / (d\omega_{2k+1} = \mathbf{0})$
Even-dimensional sphere	$S_{\mathbb{R}}^{2k}$	$\mathfrak{l}(S^{2k})$	$\mathbb{R}[\omega_{2k}, \omega_{4k-1}] / \left(\begin{array}{l} d\omega_{2k} = \mathbf{0} \\ d\omega_{4k-1} = -\omega_{2k} \wedge \omega_{2k} \end{array} \right)$
M2-extended super spacetime	$\widehat{\mathbb{T}^{10,1} \mathbf{32}}$	$\mathfrak{m}2\text{brane}$	$\mathbb{R}[\{\psi^\alpha\}_{\alpha=1}^{32}, \{e^a\}_{a=0}^{10}, h_3] / \left(\begin{array}{l} d\psi^\alpha = \mathbf{0} \\ de^a = \frac{1}{\psi} \Gamma^a \psi \\ dh_3 = \frac{i}{2} (\overline{\psi} \Gamma_{ab} \psi) \wedge e^a \wedge e^b \end{array} \right)$

- 1 Reduction via a circle bundle \Rightarrow new functors formalizing dimensional reduction via loop (and mapping) spaces with rich structure retained (topological, geometric, gauge).
- 2 The rational data of S^4 on the total space Y^{11} of a circle bundle $S^1 \rightarrow Y^{11} \rightarrow X^{10}$ leads exactly to rational data of twisted K-theory on base X^{10} . \rightarrow [see Vincent's talk]
- 3 Even if we take *flat + rational* we can still see a lot of structure: Study of cocycles in Super-Minkowski space recovers cocycles in rational twisted K-theory.
- 4 Furthermore, T-duality can be derived at the level of supercocycles.

Branes from supercocycles

- **Superspace formulation of 11d supergravity** [D'Auria-Fre]: fully controlled by an iterated pair of invariant super-cocycles μ_{M2} and μ_{M5} on $D = 11, N = 1$ super Minkowski spacetime.
- In the super homotopy-theoretic formulation [FSS]:

$$\begin{array}{ccc}
 K(\mathbb{R}, 3) & & K(\mathbb{R}, 3) \\
 \downarrow & & \downarrow \\
 \boxed{\widehat{\mathbb{T}^{10,1|32}}} & \xrightarrow{\mu_{M5}} & K(\mathbb{R}, 7) \\
 \text{fib}(\mu_{M2}) \downarrow & & \downarrow \\
 \mathbb{T}^{10,1|32} & \xrightarrow{\mu_{M2}} & K(\mathbb{R}, 4)
 \end{array}$$

\mathbb{R} -quaternionic Hopf fibration

$$\mu_{M5} = \frac{1}{5!} (\overline{\psi} \Gamma_{a_1 \dots a_5} \psi) e^{a_1} \wedge \dots \wedge e^{a_5} + h_3 \wedge \mu_{M2}$$

$$\mu_{M2} = \frac{i}{2} (\overline{\psi} \Gamma_{a_1 a_2} \psi) e^{a_1} \wedge e^{a_2}$$

which are the super-flux forms to which the M2-brane and M5-brane couple, in⁽⁴⁾ their incarnation as Green-Schwarz-type sigma models [FSS].

- $\widehat{\mathbb{T}^{10,1|32}}$ = m2brane arises as the homotopy fiber of μ_{M2} and is the extended super Minkowski spacetime or the M2-brane super Lie 3-algebra.
- μ_{M2} = super-form component of *magnetic flux* sourced by charged M5-branes.
- μ_{M5} = super-form component of *electric flux* source by charged M2-branes.

So these cocycles are avatars of M-brane charge/flux at the level of super RHT.

Twisted K-theory in type II from M-theory

- ① **Type IIA.** [BMSS] The double dimensional reduction of rational M-brane supercocycles (μ_{M2}, μ_{M5}) is indeed the tuple of F1/Dp-brane supercocycles $(\mu_{F1}, \mu_{D0}, \mu_{D2}, \mu_{D4}, \mu_{D6}, \mu_{D8})$ in rational twisted K-theory, which the literature demands to be the rational image of a cocycle in actual twisted K-theory.

Objects	Cohomology theory
M-branes	twisted Cohomotopy
D-branes	twisted K-theory



double dimensional
reduction/oxidation

[see talk by Vincent]

- ② **Type IIB.** Characterization of T-duality for circle and sphere bundles using RHT [FSS].
- **Novel effect:** T-duality in super-exceptional spacetimes in 11d M-theory [FSS][SS].

2. Integrally

① Rationally and **stably** $S_{\mathbb{Q}}^4$ is just the Eilenberg-MacLane space $K(\mathbb{Q}, 4)$, and

$$H^4(Y^{11}; \mathbb{Q}) \cong \pi_s^4(Y^{11}) \otimes \mathbb{Q}.$$

② In the **unstable** case, schematically, we have

Rational cohomotopy = Rational cohomology + trivialization of the cup square

Integrally and stably we do see new effects.

- In between full non-abelian cohomotopy and abelian ordinary cohomology sits **stable cohomotopy**, represented not by actual spheres, but by their stabilization to the sphere spectrum.
- There is a description of the C-field in each one of these flavors [FSS][BMSS].

Cohomology theory	Rational cohomology	Integral cohomology	Stable cohomotopy	Non-abelian cohomotopy
Cocycle	G_4	\tilde{G}_4	$\Sigma^\infty c$	c

Hypothesis H. *The C-field is charge-quantized in cohomotopy theory, even non-rationally.*

Cancellation of main anomalies of M-theory follows naturally from cohomotopy:

- ① C-field charge quantization in twisted cohomotopy implies various fundamental anomaly cancellation and quantization conditions [FSS].
- ② Similar effects for D-branes and orientifolds [SS].

Lifting rational S^4 to integral S^4

- If we start with the rational 4-sphere $S^4_{\mathbb{Q}}$, then how can we lift it to an “integral” space?
- The actual 4-sphere S^4 stands out as not only the most natural but the finite-dimensional one.

$$\begin{array}{ccc}
 & \xrightarrow{\text{Integral, torsion}} & S^4 \\
 Y & \xrightarrow{\text{Rational, non-torsion}} & S^4_{\mathbb{Q}} \\
 & & \downarrow
 \end{array} \tag{5}$$

- Start with integral cohomology as describing the (shifted/twisted) C-field and then transition to a description in terms of cohomotopy. By representability, this amounts to lifting


$$\begin{array}{ccc}
 & \xrightarrow{\text{Nonlinear prequantum}} & S^4 \\
 Y & \xrightarrow{\text{Linear quantum}} & K(\mathbb{Z}, 4) \\
 & & \downarrow \iota
 \end{array} \tag{6}$$

The map ι assembles, upon taking homotopy classes, into the integral cohomology $H^4(S^4; \mathbb{Z})$ generated by a fundamental class.

- Description:

$$\text{C-field in } \pi^4(Y^{11}) \iff \text{C-field in } H^4(Y^{11}; \mathbb{Z}) + \text{nontrivial conditions.}$$

Proposition (Integral Postnikov tower for S^4 [GS])



Note that at the top level the three conditions vanish necessarily on Y^{11} , for dimension reasons.

Cohomotopy in deg 4 \sim Integral 4-cohomology + four sets of obstructions.

Pulling back to spacetime Y , where the fundamental class ι_4 pulls back to the field

$$G_4 - \frac{1}{2}\lambda =: \tilde{G}_4 = f^* \iota_4$$

where $\lambda = \frac{1}{2}p_1$ is the first Spin characteristic class of TY .

(i) First obstruction.

$$\boxed{\text{Sq}^2 \tilde{G}_4 \stackrel{!}{=} 0 \in H^6(Y; \mathbb{Z}_2)}$$

This follows from anomaly cancellation in M-theory [FSS].

(ii) Second obstruction.

$$\boxed{f^*(\alpha_7) \stackrel{!}{=} 0 \in H^7(Y; \mathbb{Z}_2)}$$

where α_7 is a secondary operation, restricting fiberwise to $\text{Sq}^2 \iota_5$.
No candidate degree 5 classes.

(iii) Third obstructions.

$$\boxed{f^*(\text{"Sq}^4 \iota_4\text{"}) \stackrel{!}{=} 0 \in H^8(Y; \mathbb{Z}_8)}$$

- Note that by construction, this implies also that (upon mod 2 reduction)

$$f^*(\text{Sq}^4 \iota_4) = \text{Sq}^4 f^*(\iota_4) = \text{Sq}^4 \tilde{G}_4 = \tilde{G}_4 \cup \tilde{G}_4 = 0 \in H^8(Y; \mathbb{Z}_2).$$

- Recall that rationally we have the EOM $d * G_4^{\text{form}} = \frac{1}{2} G_4^{\text{form}} \wedge G_4^{\text{form}}$..
- Coefficients being \mathbb{Z}_8 rather than \mathbb{Z}_2 : Fields reduced modulo 4:
 $\frac{1}{2}\lambda \rightsquigarrow$ modding out p_1 by 4. (Pontrjagin square operation).

We also have $\mathcal{P}_3^1 \iota_4 = 0$.

- Mod 3 reductions are shown to play a prominent role in topological considerations in M-theory [S], where similar conditions, including $\mathcal{P}_3^1 \rho_3 G_4 = 0$, have been highlighted in the context of *Spin K-theory*.

(iv) Fourth obstruction.

$$f^*(P_{11}) \stackrel{!}{=} 0$$

where P_{11} is a class which fiberwise restricts to $\text{Sq}^4 \iota_7$.

- Reminiscent of $G_4 \wedge G_7$.
- The universal coefficient theorem gives detectable effect for M-theory on orientable spacetimes.

(v) Fifth obstructions.

$$\text{“Sq}^8 \iota_4 \stackrel{!}{=} 0, \quad \iota_4^3 \stackrel{!}{=} 0, \quad \mathcal{P}_5^1 \iota_4 \stackrel{!}{=} 0$$

These obstructions necessarily vanish on Y^{11} . However on a 12-manifold Z^{12} , for analyzing the congruences of the Chern-Simons term in the M-theory action, the three conditions are nontrivial (but natural to have).

Proposition (Cohomotopy vs. cohomology for the C-field)

Consider the M-theory (shifted) C-field \tilde{G}_4 as an integral cohomology class in degree four. Then if \tilde{G}_4 lifts to a cohomotopy class $\mathcal{G}_4 \in \pi^4(Y^{11})$ the following obstructions necessarily vanish

- (i) $\text{Sq}^2 \tilde{G}_4 = 0 \in H^6(Y^{11}; \mathbb{Z}_2)$.
- (ii) $\mathcal{P}_3^1(\tilde{G}_4) = 0 \in H^8(Y^{11}; \mathbb{Z}_3)$.
- (iii) $\text{Sq}^4 \tilde{G}_4 = \tilde{G}_4 \cup \tilde{G}_4 = 0 \in H^8(Y^{11}; \mathbb{Z}_2)$.
- (iv) If $G_4 = 0$ and $dC_3 = 0$ can be lifted to an integral class \tilde{C}_3 , then we also have $\text{Sq}^3 \text{Sq}^1 \tilde{C}_3 = 0 \in H^7(Y^{11}; \mathbb{Z}_2)$.
- (v) If $dG_7 = G_4 \wedge G_4 = 0$ and G_7 can be lifted to an integral class \tilde{G}_7 , then we also have the condition $\text{Sq}^4 \tilde{G}_7 = 0 \in H^{11}(Y^{11}; \mathbb{Z}_2)$.

Consequences:

- 1 **Congruences for the action** The Chern-Simons term in the action

$$\frac{1}{6} \int_{Y^{11}} C_3 \wedge G_4 \wedge G_4 .$$

Since C_3 may not be globally defined in general, one may consider Y^{11} as the boundary of a 12-manifold Z^{12} and analyzes the globally well defined term

$$\frac{1}{6} \int_{Z^{12}} G_4 \wedge G_4 \wedge G_4 \quad (7)$$

[Witten]: usual quantization law of G_4 does not give rise to a well defined Chern-Simons action, as (7) might fail to be integral by a **factor of 6**.

Cohomotopy implies the added condition that

$$\tilde{G}_4^3 \equiv 0 \pmod{3} .$$

This, with $\tilde{G}_4^2 = \text{Sq}^4(\tilde{G}_4) \equiv 0 \pmod{2}$, gives result (without E_8 -gauge theory).

- 2 **The anomaly in the partition function** Quantization in cohomotopy yields the condition $\text{Sq}^2(\tilde{G}_4) = 0$ for some integral lift of G_4 .

- Implies the vanishing of the DMW anomaly $\text{Sq}^3(\tilde{G}_4) = 0$ [FSS].
- Obstruction theory for $S^4 \Rightarrow$ fields which contribute to the phase are just the field which lift to the first Postnikov stage in cohomotopy [GS].

Example (Flux compactification spaces)

Anti-de Sitter space $\text{AdS}_n \rightsquigarrow$ simply-connected cover $\widetilde{\text{AdS}}_n$ of AdS_n .

- 1 $\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2 \times T^2$: Supersymmetry without supersymmetry [Duff-Lu-Pope] and T-duality [Bouwknegt-Evslin-Mathai]. $\pi^4(\mathbb{C}P^2) \cong \mathbb{Z}$ while $H^4(\mathbb{C}P^2; \mathbb{Z}) \cong \mathbb{Z}$.
- 2 $\widetilde{\text{AdS}}_7 \times \mathbb{R}P^4$: M-theory on an orientifold [Witten][Hori]. $\pi^4(\mathbb{R}P^4) \cong \mathbb{Z}_2$ while $H^4(\mathbb{R}P^4; \mathbb{Z}) = 0$, indeed shows that cohomotopy detects more.
- 3 $\widetilde{\text{AdS}}_4 \times \mathbb{R}P^5 \times T^2$: $\pi^4(\mathbb{R}P^5)$ is cyclic of order 4, i.e. either \mathbb{Z}_4 or $\mathbb{Z}_2 \times \mathbb{Z}_2$, while $H^4(\mathbb{R}P^5; \mathbb{Z}) \cong \mathbb{Z}_2$.
- 4 $\widetilde{\text{AdS}}_4 \times \mathbb{C}P^3 \times S^1$: $\pi^4(\mathbb{C}P^3) \cong \mathbb{Z} \oplus \mathbb{Z}_2$ while $H^4(\mathbb{C}P^3; \mathbb{Z}) \cong \mathbb{Z}$, so that there is an extra contribution of \mathbb{Z}_2 present in cohomotopy.
- 5 For $\mathbb{H}P^2$: $\pi^4(\mathbb{H}P^2) \cong \mathbb{Z}$ while $H^4(\mathbb{H}P^2; \mathbb{Z}) \cong \mathbb{Z}$, and hence no new contribution,
- 6 For $\mathbb{O}P^2$: $\pi^4(\mathbb{O}P^2) \cong \mathbb{Z}$. while $H^4(\mathbb{O}P^2; \mathbb{Z}) = 0$, signaling a new effect. Important for bosonic M-theory ([Ramond][S]).

Interpretation and consequences? Work in progress (via Pontrjagin-Thom theory).

Twisted Cohomotopy theory [FSS]

In degree $d - 1$ there is a canonical twisting on Riemannian d -manifolds, given by the unit sphere bundle in the orthogonal tangent bundle:

$$\begin{array}{c}
 \text{J-twisted Cohomotopy theory } \pi^{TX^d}(X^d) := \left\{ \begin{array}{c}
 \begin{array}{ccc}
 & \begin{array}{c} \text{tangent} \\ \text{unit sphere bundle} \end{array} & \begin{array}{c} \text{universal tangent} \\ \text{unit sphere bundle} \end{array} \\
 & S(TX^d) & \longrightarrow S^{d-1} // O(d) \\
 \begin{array}{c} \text{continuous section} \\ \text{= twisted cocycle} \end{array} \nearrow & \downarrow p & \downarrow \\
 X & \xlongequal{\quad} X & \xrightarrow{TX^d} BO(d) \\
 & \text{classifying map of} & \\
 & \text{tangent/frame bundle} &
 \end{array} \right\} \Big/ \sim \frac{\text{homotopy}}{BO(d)}
 \end{array}
 \\
 \\
 \approx \left\{ \begin{array}{ccc}
 X & \overset{\text{continuous function}}{\dashrightarrow} & S^{d-1} // O(d) \\
 \swarrow & \searrow \text{homotopy} & \downarrow \\
 & & BO(d) \\
 \downarrow TX^d & & \swarrow \\
 \text{twist} & &
 \end{array} \right\} \Big/ \sim \frac{\text{homotopy}}{BO(d)}
 \end{array}$$

Since the canonical morphism $O(d) \rightarrow \text{Aut}(S^{d-1})$ is known as the *J-homomorphism*, we may call this *J-twisted Cohomotopy theory*, for short.

Twisted cohomotopy and anomalies [FSS]

Hypothesis H: *The C-field 4-flux & 7-flux forms in M-theory are subject to charge quantization in J-twisted Cohomotopy cohomology theory in that they are in the image of the non-abelian Chern character map from J-twisted Cohomotopy theory.*

⇒ Cancellation of main anomalies:

Half-integral flux quantization	$\underbrace{\left[G_4 + \frac{1}{4} p_1 \right]}_{=: \tilde{G}_4 \text{ integral flux}} \in H^4(X, \mathbb{Z})$
Background charge	$\underbrace{q(\tilde{G}_4)}_{\text{quadratic form}} = \tilde{G}_4 \left(\underbrace{\tilde{G}_4 - \frac{1}{2} p_1}_{=(\tilde{G}_4)_0} \right)$
DMW-anomaly cancellation	$W_7(TX) = 0$
Integral equation of motion	$\underbrace{\text{Sq}^3(\tilde{G}_4)}_{=\beta \text{Sq}^2} = 0$
M5-brane anomaly cancellation	$\underbrace{I_{\text{ferm}}^{\text{M5}}}_{\text{chiral fermion}} + \underbrace{I_{\text{sd}}^{\text{M5}}}_{\text{self-dual 3-flux}} + \underbrace{I_{\text{infl}}^{\text{bulk}}}_{\text{bulk inflow}} = 0$
M2-brane tadpole cancellation	$\underbrace{N_{\text{M2}}}_{\text{number of M2-branes}} + q(\tilde{G}_4) = \underbrace{I_8}_{\text{One loop polynomial}}$

Consequences for WZW model associated to M5-brane ⇒ [\[See talk by Domenico\]](#)

J-Twisted Cohomotopy and Topological G-Structure

- For every topological coset space realization G/H of an n -sphere, there is a canonical homotopy equivalence between the classifying spaces for G -twisted Cohomotopy and for topological H -structure (i.e., reduction of the structure group to H), as follows:



$$\begin{array}{ccc} \text{coset space structure} & & \text{G-twisted Cohomotopy /} \\ \text{on topological } n\text{-sphere} & & \text{topological } H\text{-structure} \\ \\ S^n \underset{\text{homeo}}{\simeq} G/H & \Rightarrow & S^n // G \underset{\text{htpy}}{\simeq} BH . \end{array}$$

(One may think of this as “moving G from numerator on the right to denominator on the left”.)



- Existence of a G -structure is a non-trivial topological condition, so is the existence of J -twisted Cohomotopy cocycles.
- Notice that this is a special effect of twisted non-abelian generalized Cohomology: A non-twisted generalized cohomology theory (abelian or non-abelian) always admits at least one cocycle, namely the trivial or zero-cocycle. But here for non-abelian J -twisted Cohomotopy theory on 8-manifolds, the existence of *any* cocycle is a non-trivial topological condition.

Equivalence for Spin 8-manifolds

$$\simeq S^7 // \text{Spin}(8) \\ \simeq B\text{Spin}(7)$$

 \Rightarrow

 \simeq


$$S^7 // \text{Sp}(2) \cdot \text{Sp}(1) \\ \simeq B\text{Sp}(1) \cdot \text{Sp}(1)$$

 \Rightarrow

 \simeq


Stable vs. unstable

The quaternionic Hopf fibration.

$$\begin{array}{c}
 \text{quaternionic Hopf fibration} \\
 h_{\mathbb{H}} \\
 \curvearrowright \\
 S^7 \xrightarrow{\simeq} S(\mathbb{H}^2) \xrightarrow{(q_1, q_2) \mapsto [q_1 : q_2]} \mathbb{H}P^1 \xrightarrow{\simeq} S^4, \\
 \text{unit sphere} \quad \text{quaternionic} \\
 \text{in quaternionic} \quad \text{projective} \\
 \text{2-space} \quad \text{1-space}
 \end{array}$$

which represents a generator of the non-torsion subgroup in the 4-Cohomotopy of the 7-sphere, as shown on the left here:

$$\begin{array}{c}
 \text{quaternionic Hopf fibration} \\
 \text{non-torsion generator} \\
 [S^7 \xrightarrow{h_{\mathbb{H}}} S^4] \quad \pi^4(S^7) \xrightarrow{\Sigma^\infty} S^4(S^7) \quad \Sigma^\infty [S^7 \xrightarrow{h_{\mathbb{H}}} S^4] \quad \text{stabilized quaternionic Hopf fibration} \\
 \text{non-abelian/unstable Cohomotopy group} \quad \text{stabilization} \quad \text{abelian/stable Cohomotopy group} \\
 (1, 0) \in \mathbb{Z} \times \mathbb{Z}_{12} \xrightarrow{(n, a) \mapsto (n \bmod 24)} \mathbb{Z}_{24} \ni 1 \quad \text{torsion generator}
 \end{array}$$

- So composition with the quaternionic Hopf fibration can be viewed as a *transformation* that translates deg-7 to deg-4 Cohomotopy classes:

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & & S^7 \\
 & \swarrow c & \downarrow h_{\mathbb{H}} \\
 X & \xrightarrow{\quad} & S^4 \\
 & \searrow (h_{\mathbb{H}})_*(c) & \\
 & &
 \end{array} & \begin{array}{l}
 \text{7-Cohomotopy} \\
 \text{reflects into} \\
 \text{4-Cohomotopy}
 \end{array} & \begin{array}{c}
 \pi^7(X) \\
 \downarrow (h_{\mathbb{H}})_* \\
 \pi^4(X)
 \end{array}
 \end{array}$$

Proposition (Differential form data underlying twisted Cohomotopy)

Let X be a simply connected smooth manifold and $\tau : X \rightarrow BO(n+1)$ a twisting for Cohomotopy in degree n . Let ∇_τ be any connection on the real vector bundle V classified by τ with Euler form $\chi_{2k+2}(\nabla_\tau)$ (see [Mathai-Quillen]).

(i) If $n = 2k + 1$ is odd $n \geq 3$: a cocycle defining a class in the rational τ -twisted Cohomotopy of X is equivalently given by

$$\pi_{\mathbb{Q}}^\tau(X) \simeq \left\{ G_{2k+1} \mid d G_{2k+1} = \chi_{2k+2}(\nabla_\tau) \right\} / \sim.$$

(ii) If $n = 2k$ is even, $n \geq 2$: a cocycle defining a class in the rational τ -twisted Cohomotopy of X is given by a pair of differential forms $G_{2k} \in \Omega^{2k}(X)$ and $G_{4k-1} \in \Omega^{4k-1}(X)$ such that

$$dG_{2k} = 0; \quad \pi^* G_{2k} = \frac{1}{2} \chi_{2k}(\nabla_{\hat{\tau}})$$

$$dG_{4k-1} = -G_{2k} \wedge G_{2k} + \frac{1}{4} p_k(\nabla_\tau),$$

where $p_k(\nabla_\tau)$ is the k -th Pontrjagin form of ∇_τ , $\pi : E \rightarrow X$ is the unit sphere bundle over X associated with τ , $\hat{\tau} : E \rightarrow BO(n)$ classifies the vector bundle \hat{V} on E defined by the splitting $\pi^* V = \mathbb{R}_E \oplus \hat{V}$ associated with the tautological section of $\pi^* V$ over E , and $\nabla_{\hat{\tau}}$ is the induced connection on \hat{V} . That is,

$$\pi_{\mathbb{Q}}^\tau(X) \simeq \left\{ (G_{2k}, G_{4k-1}) \mid \begin{array}{l} d G_{2k} = 0, \quad \pi^* G_{2k} = \frac{1}{2} \chi_{2k}(\nabla_{\hat{\tau}}) \\ d G_{4k-1} = -G_{2k} \wedge G_{2k} + \frac{1}{4} p_k(\nabla_\tau) \end{array} \right\} / \sim.$$

3. Differentially

Differential refinement

- Refine the topological lift (5) to a geometric lift at the level of smooth stacks of the form

$$\begin{array}{ccc} & & \widehat{S^4} \\ & \nearrow \text{Differential cohomology,} & \downarrow \\ & \text{prequantum and geometric} & \\ Y & \xrightarrow{\text{Differential cocycle,}} & B^3U(1)_{\nabla} \\ & \text{quantum and geometric} & \end{array} \quad (8)$$

where $\widehat{S^4}$ is the differential refinement of the 4-sphere and $B^3U(1)_{\nabla}$ is the smooth stack of 3-bundles with connections

- This would require a differential refinement of the *Postnikov tower* which uses refinement of cohomology operations, primary (such as Steenrod operations) and secondary (such as Massey products) [GS].

Differential cohomotopy [Fiorenza-S.-Schreiber]

- **\mathbb{H} -Hopf fibration:** $S^3 \rightarrow S^7 \rightarrow S^4 \rightarrow BSU(2) \xrightarrow{c_2} K(\mathbb{Z}, 4)$.
- Rationalize: $S_{\mathbb{Q}}^3 \rightarrow S_{\mathbb{Q}}^7 \rightarrow S_{\mathbb{Q}}^4 \rightarrow (BS^3)_{\mathbb{Q}}$ which is equivalent to
$$K(\mathbb{Q}, 7) \rightarrow S_{\mathbb{Q}}^4 \rightarrow K(\mathbb{Q}, 4)$$
- Rational homotopy of spaces can be modelled using L_{∞} -algebras.
- The Eilenberg-MacLane spaces $K(\mathbb{Q}, n) = B^n\mathbb{Q}$ can be modelled using algebras via chain complexes: $b^n\mathbb{Q} = \mathbb{Q}[n]$.
- Lie 7- algebra \mathfrak{s}^4 is defined by $\text{CE}(\mathfrak{s}^4) = \mathbb{R}[g_4, g_7]$ with g_k in degree k and with the differential defined by $dg_4 = 0, dg_7 = g_4 \wedge g_4$.
- Has a natural structure of infinitesimal $\mathbb{R}[2]$ -quotient of $\mathbb{R}[6]$, i.e., there exists a natural homotopy fiber sequence of L_{∞} -algebras

$$\begin{array}{ccc} \mathbb{R}[6] & \longrightarrow & \mathfrak{s}^4 \\ \downarrow & & \downarrow^P \\ 0 & \longrightarrow & \mathbb{R}[3] \end{array} \quad (9)$$

Theorem (FSS)

The system $(\widehat{G}_4, \widehat{G}_7)$ forms a cocycle in differential cohomotopy.

Differential refinements: $B^3U(1)_{\nabla}$ vs. \widehat{S}^4

- Let \mathfrak{s}^4 be the Lie 7-algebra whose corresponding Chevelley-Eilenberg algebra is the exterior algebra on generators g_4 and g_7 with relations

$$dg_4 = 0, \quad dg_7 = g_4 \wedge g_4.$$

- As a de Rham model for flat 1-forms with values in S^4 we take the sheaf on the site of Cartesian spaces given by the assignment

$$\Omega_{\mathbb{H}}^1(-; \mathfrak{s}^4) : U \longmapsto \text{hom}_{\text{dgcAlg}}(\text{CE}(\mathfrak{s}^4), \Omega^*(U)),$$

for each Cartesian space $U \cong \mathbb{R}^n$. (The homotopy type of $\Omega_{\mathbb{H}}^1(-; \mathfrak{s}^4)$ can be computed via the Sullivan construction as the \mathbb{R} -local 4-sphere $S_{\mathbb{R}}^4$).

- Then pulling back along the canonical map $S^4 \rightarrow S_{\mathbb{R}}^4$, we get a smooth stack

$$\begin{array}{ccc} \widehat{S}^4 & \longrightarrow & \Omega_{\mathbb{H}}^1(-; \mathfrak{s}^4) \\ \downarrow & & \downarrow \\ S^4 & \longrightarrow & S_{\mathbb{R}}^4. \end{array}$$

Definition (Differential unstable cohomotopy)

For a smooth manifold X , let $i(X)$ denote its embedding as a smooth stacks via its sheaf of smooth plots. Then the differential cohomotopy of X in degree 4 is defined as the pointed set $\widehat{\pi}_U^4(X) := \pi_0 \text{Map}(i(X), \widehat{S}^4)$ where the maps on the right are those of smooth stacks.

Differential cohomotopy: stably

- Stably, S^4 has only torsion groups in higher degrees and hence the canonical map $S^4 \rightarrow K(\mathbb{R}, 4)$ is a stable \mathbb{R} -local equivalence.
- Geometrically, the realification is modeled by closed 4-forms $\Omega_{\text{cl}}^4(-)$.
- Stable differential cohomotopy in degree 4 fits into a pullback square

$$\begin{array}{ccc} \widehat{\Sigma^\infty S^4} & \longrightarrow & H(\tau^{\leq 0} \Omega^{4+*}(-)) \\ \downarrow & & \downarrow \\ \Sigma^\infty S^4 & \longrightarrow & \Sigma^4 H\mathbb{R} . \end{array}$$

where $\Omega^{4+*}(-)$ denotes the de Rham complex, shifted so that Ω^4 is in degree zero, and $\tau^{\leq 0}$ truncates the complex in degree zero so that the complex is concentrated in negative degrees. The functor H denotes the Eilenberg-MacLane functor which turns a chain complex into a spectrum.

Definition (Differential stable cohomotopy)

Let X be a smooth manifold with $i(X)$ its associated smooth stack. The *stable* differential cohomotopy group of X is defined as

$$\widehat{\pi}_s^4(X) := \pi_0 \text{Map}(i(X); (\widehat{\Sigma^\infty S^4})_0).$$

where the subscript 0 denotes the deg 0 component of the sheaf of spectra $\widehat{\Sigma^\infty S^4}$.

Definition (Geometric cohomotopy cocycles [GS])

If X is a smooth manifold, a morphism $\hat{c} : X \rightarrow \widehat{S}^4$ can be identified with a triple (c, h, ω) where

- (i) $c : X \rightarrow S^4$ is a cocycle in ordinary cohomotopy,
- (ii) $\omega : \text{CE}(\mathfrak{s}^4) \rightarrow \Omega^*(X)$ is a DGA morphism, determined by specifying forms ω_4 and ω_7 on M satisfying $d\omega_7 = \omega_4^2$ and $d\omega_4 = 0$,
- (iii) and h is a homotopy interpolating between the rational cocycle represented by the form data and the rationalization of the classifying map $c : X \rightarrow S^4$. Thus, h exhibits a sort of *de Rham theorem* for cohomotopy.

Proposition (Differential refinement of Postnikov tower of the sphere)

$$\begin{array}{ccccc}
 K(\mathbb{Z}_{15}, 11) & \longrightarrow & (\widehat{S}^4)_7 & & \\
 & & \downarrow & & \\
 K(\mathbb{Z}_{24} \times \mathbb{Z}_3, 10) & \longrightarrow & (\widehat{S}^4)_6 & \longrightarrow & K(\mathbb{Z}_{15}, 12) \\
 & & \downarrow & & \\
 K(\mathbb{Z}_2 \times \mathbb{Z}_2, 9) & \longrightarrow & (\widehat{S}^4)_5 & \longrightarrow & K(\mathbb{Z}_{24} \times \mathbb{Z}_3, 11) \\
 & & \downarrow & & \\
 K(\mathbb{Z}_2 \times \mathbb{Z}_2, 8) & \longrightarrow & (\widehat{S}^4)_4 & \longrightarrow & K(\mathbb{Z}_2 \times \mathbb{Z}_2, 10) \\
 & & \downarrow & & \\
 K(\mathbb{Z}_{12}, 7) \times K(\mathbb{Z}, 7) & \longrightarrow & (\widehat{S}^4)_3 & \longrightarrow & K(\mathbb{Z}_2 \times \mathbb{Z}_2, 9) \\
 & & \downarrow & & \\
 K(\mathbb{Z}_2, 6) & \longrightarrow & (\widehat{S}^4)_2 & \xrightarrow{(\cdot, \iota_4^2)} & K(\mathbb{Z}_{12}, 8) \times B^7 U(1)_\nabla \\
 & & \downarrow & & \\
 K(\mathbb{Z}_2, 5) & \longrightarrow & (\widehat{S}^4)_1 & \xrightarrow{\alpha_{7^l}} & K(\mathbb{Z}_2, 7) \\
 & & \downarrow & & \\
 & & (\widehat{S}^4)_0 = B^3 U(1)_\nabla & \xrightarrow{Sq^2 \rho_{2^l}} & K(\mathbb{Z}/2, 6)
 \end{array}$$

where we have identified the first few obstructions.

Proposition (Differential cohomotopy vs. cohomology for the C-field)

Consider the differentially refined M-theory (shifted) C-field \widehat{G}_4 as an integral cohomology class in degree four. Then if \widehat{G}_4 lifts to a cohomotopy class $\mathcal{G}_4 \in \widehat{\pi}^4(Y^{11})$ the following obstructions necessarily vanish

- (i) $Sq^2 I(\widehat{G}_4) = 0 \in H^6(Y^{11}; \mathbb{Z}_2)$.
- (ii) $\mathcal{P}_3^1 I(\widehat{G}_4) = 0 \in H^8(Y^{11}; \mathbb{Z}_3)$.
- (iii) $Sq^4 I(\widehat{G}_4) = I(\widehat{G}_4 \cup_{DB} \widehat{G}_4) = 0 \in H^8(Y^{11}; \mathbb{Z}_2)$.
- (iv) If $\widehat{G}_4 = 0$ and C_3^{form} is quantized, with differential refinement \widehat{C}_3 , then we also have $Sq^3 Sq^1 I(\widehat{C}_3) = 0 \in H^7(Y^{11}; \mathbb{Z}_2)$.
- (v) If $dG_7^{\text{form}} = G_4^{\text{form}} \wedge G_4^{\text{form}} = 0$ and G_7^{form} is quantized, with differential refinement \widehat{G}_7 , then we also have the condition $Sq^4 I(\widehat{G}_7) = 0 \in H^{11}(Y^{11}; \mathbb{Z}_2)$.

Remark (Obstruction in M-theory via higher bundles with connections)

Déglise–Beilinson cup product in M-theory $\widehat{G}_4 \cup_{DB} \widehat{G}_4$ gives a 7-bundle with connection form locally given by $C_3^{\text{form}} \wedge G_4^{\text{form}}$ [FSS]. From identification of the k -invariant at 2nd stage (the DB square): to lift past the 2nd stage in the Postnikov tower for \widehat{S}^4 , this connection must be globally defined. In terms of differential cohomology, $a(C_3^{\text{form}} \wedge G_4^{\text{form}}) = \widehat{G}_4 \cup_{DB} \widehat{G}_4$, where $a : \Omega^7(Y^{11}) \rightarrow \widehat{H}^8(Y^{11})$ is the canonical map.

Example (Differential cohomotopy of flux compactification spaces)

LES in stable cohomotopy

$$\dots \longrightarrow \pi_s^3(X) \xrightarrow{\text{deg}} \Omega^3(X) \longrightarrow \widehat{\pi}_s^4(X) \longrightarrow \pi_s^4(X) \longrightarrow \dots$$

allows to compute some examples.

- (i) $\widetilde{\text{AdS}}_7 \times \mathbb{R}P^4$: $\widehat{\pi}_s^4(\widetilde{\text{AdS}}_7 \times \mathbb{R}P^4) \cong \widehat{H}^4(\widetilde{\text{AdS}}_7 \times \mathbb{R}P^4)$.
- (ii) $\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2$: $\widehat{\pi}_s^4(\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2) \cong \widehat{H}^4(\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2)$.
- (iii) $\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2 \times T^2$: $\widehat{\pi}_s^4(\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2 \times T^2) \cong \widehat{H}^4(\widetilde{\text{AdS}}_4 \times \mathbb{C}P^2 \times T^2)$.
- (iv) $\widetilde{\text{AdS}}_4 \times \mathbb{R}P^5 \times T^2$: $\pi^4(\mathbb{R}P^5)$ is order 4, either \mathbb{Z}_4 or $\mathbb{Z}_2 \times \mathbb{Z}_2$, while $H^4(\mathbb{R}P^5; \mathbb{Z}) \cong \mathbb{Z}_2$. Also $\pi^3(\mathbb{R}P^5)$ is finite. We therefore have a short exact sequence

$$0 \longrightarrow \Omega^3(\mathbb{R}P^5) \longrightarrow \widehat{\pi}^4(\mathbb{R}P^5) \longrightarrow \pi^4(\mathbb{R}P^5) \longrightarrow 0.$$

Since $\pi^4(\mathbb{R}P^5)$ is generated by $q_5\eta_4$, with $\eta_4 : S^5 \rightarrow S^4$ the two-fold suspension of the Hopf map, the induced map on H^4 necessarily vanishes. Hence, in this case, differential cohomotopy yields considerably different information than ordinary differential cohomology.

Back to




:

Cohomotopy \Rightarrow branes and gauge theory

Differential cohomotopy and D-brane gauge theories

Zoom in beyond foundational/structural M-theoretic considerations [SS]:

- ④ A differential refinement of Cohomotopy cohomology theory is given by *un-ordered configuration spaces of points*.
- ④ The fiber product of such differentially refined Cohomotopy cocycle spaces describing $D6 \perp D8$ -brane intersections is homotopy-equivalent to the *ordered configuration space of points* in the transversal space.
- ④ The higher observables on this moduli space are equivalently weight systems on horizontal chord diagrams.



Combining the above seemingly distinct mathematical areas reflect a multitude of effects expected on brane intersections in string theory. So aside from structural utility for M-theory, Hypothesis H implies:

- *M-theoretic observables on $D6 \perp D8$ -configurations* (cf. parametrized).
- *Chan-Paton observables.*
- *String topology operations.*
- *Multi-trace observables of BMN matrix model.*
- *Hanany-Witten states.*
- *BLG 3-Algebra observables.*
- *Bulk Wilson loop observables.*
- *Single-trace observables*
- *of SYK & BMN model.*
- *Fuzzy funnel observables.*
- *Supersymmetric indices.*
- *'t Hooft string amplitudes.*

[See talk by Urs]

Top-down M-theory via Hypothesis H: knowledge about gauge field theory and perturbative string theory is not used in deriving the algebras of observables of M-theory, but only to interpret them.

Thank you!