

Syllepsis in Homotopy Type Theory

Kristina Sojakova

July 19, 2021

Abstract

It is well-known that in homotopy type theory (HoTT), one can prove the *Eckmann-Hilton* theorem: given two 2-loops $\mathbf{p}, \mathbf{q} : 1_\star = 1_\star$ on the reflexivity path at an arbitrary point $\star : A$, we have $\mathbf{p} \cdot \mathbf{q} = \mathbf{q} \cdot \mathbf{p}$. If we go one dimension higher, *i.e.*, if \mathbf{p} and \mathbf{q} are 3-loops $\mathbf{p}, \mathbf{q} : 1_{1_\star} = 1_{1_\star}$, we show that a property classically known as *syllepsis* also holds in HoTT: namely, the Eckmann-Hilton proof for \mathbf{q} and \mathbf{p} is the inverse of the Eckmann-Hilton proof for \mathbf{p} and \mathbf{q} .

1 The Eckmann-Hilton Argument

Numerous short proofs of Eckmann-Hilton (EH) have been given by various people, including Favonia, Dan Christensen, and Mike Shulman. They are all propositionally equal but may have slightly different computational behavior. We start by outlining yet another such proof, which is particularly suitable for our proof of syllepsis. This proof is constructed differently from the aforementioned ones, but is again propositionally equal (it is in fact rather difficult to come up with a proof of EH that is *not* propositionally equal to the existing ones). This will also serve to establish some notation and preliminary definitions.

If we go one dimension lower, EH will not hold: it is not hard to see that loops $p, q : a = a$ in general do not commute (for example, endofunctions in the universe). So to prove EH, we must make use of the fact that \mathbf{p} and \mathbf{q} are based at a path (and an identity one at that). One obvious thing we can do with a path is to compose it with another path; in this case, the only 1-path we have at our disposal is the reflexivity path 1_\star itself. Since all functions in HoTT are functorial, this gives us 2-paths $\mathbf{whisk-L}(1_\star, \mathbf{p}), \mathbf{whisk-L}(1_\star, \mathbf{q}) : 1_\star \cdot 1_\star = 1_\star \cdot 1_\star$ if we concatenate with 1_\star on the left and $\mathbf{whisk-R}(\mathbf{p}, 1_\star), \mathbf{whisk-R}(\mathbf{q}, 1_\star) : 1_\star \cdot 1_\star = 1_\star \cdot 1_\star$ if we concatenate with 1_\star on the right:

Lemma 1. *For any points $a, b, c : A$, 1-paths $u : a = b$, $x, y : b = c$, and 2-path $q : x = y$, we have a term*

$$\mathbf{whisk-L}(u, p) : u \cdot x = u \cdot y$$

Pictorially:

$$\begin{array}{ccc} a & \xrightarrow{u} & b & \xrightarrow{x} & c \\ & & & \Downarrow q & \\ a & \xrightarrow{u} & b & \xrightarrow{y} & c \end{array}$$

Lemma 2. *For any points $a, b, c : A$, 1-paths $u, v : a = b$, $x : b = c$, and 2-path $p : u = v$, we have a term*

$$\mathbf{whisk-R}(q, x) : u \cdot x = v \cdot x$$

Pictorially:

$$\begin{array}{ccc} a & \xrightarrow{u} & b & \xrightarrow{x} & c \\ & \Downarrow p & & & \\ a & \xrightarrow{v} & b & \xrightarrow{x} & c \end{array}$$

Whiskering is a special case of a *parallel composition* of paths:

Lemma 3. For any points $a, b, c : A$, 1-loops $u, v : a = b$, $x, y : b = c$, and 2-paths $r : u = v$, $s : x = y$, we have a term

$$p \bullet \bullet q : u \bullet x = v \bullet y$$

Pictorially:

$$\begin{array}{ccc} a & \xrightarrow{u} & b & \xrightarrow{x} & c \\ & p \Downarrow & & q \Downarrow & \\ a & \xrightarrow{v} & b & \xrightarrow{y} & c \end{array}$$

As the picture above suggests, parallel composition and ordinary composition of paths satisfy an exchange law: given 1-paths $w : a = b$, $z : b = c$ and 2-paths $r : v = w$, $s : y = z$, we have $(p \bullet \bullet q) \bullet (r \bullet \bullet s) = (p \bullet r) \bullet \bullet (q \bullet s)$. It is not surprising, then, that the operations of whiskering on the left and whiskering on the right commute:

Lemma 4. For any points $a, b, c : A$, 1-paths $u, v : a = b$, $x, y : b = c$, and 2-paths $p : u = v$, $q : x = y$, we have a term

$$\text{whisk-L-R}(p, q) : \text{whisk-L}(u, q) \bullet \text{whisk-R}(p, y) = \text{whisk-R}(p, x) \bullet \text{whisk-L}(v, q)$$

Pictorially:

$$\begin{array}{ccc} u \bullet x & \xrightarrow{\text{whisk-R}(p, x)} & v \bullet x \\ \text{whisk-L}(u, q) \Big| & & \Big| \text{whisk-L}(v, q) \\ u \bullet y & \xrightarrow{\text{whisk-R}(p, y)} & v \bullet y \end{array}$$

In our case, this means the commuting diagram below:

$$\begin{array}{ccc} 1_\star \bullet 1_\star & \xrightarrow{\text{whisk-R}(q, 1_\star)} & 1_\star \bullet 1_\star \\ \text{whisk-L}(1_\star, p) \Big| & & \Big| \text{whisk-L}(1_\star, p) \\ 1_\star \bullet 1_\star & \xrightarrow{\text{whisk-R}(q, 1_\star)} & 1_\star \bullet 1_\star \end{array}$$

On the other hand, $1_\star \bullet 1_\star$ is just 1_\star , so a reasonable question might be whether $\text{whisk-L}(1_\star, p)$ is equal to p and $\text{whisk-R}(q, 1_\star)$ to q . Of course, the answer is yes, so the above diagram essentially proves EH. Nevertheless, since we will be reasoning about EH computationally, we need to give the proof in detail.

To prove the equalities $\text{whisk-L}(1_\star, p) = p$ and $\text{whisk-R}(q, 1_\star) = q$, we expand them into commuting diagrams:

Lemma 5. Concatenation on the left by reflexivity is natural: for any points $a, b : A$, 1-paths $u, v : a = b$, and 2-path $p : u = v$, we have a term

$$\blacksquare\text{-1-L-nat}(p) : \text{whisk-L}(1_a, p) \bullet \blacksquare\text{-1-L}(v) = \blacksquare\text{-1-L}(u) \bullet p$$

Pictorially:

$$\begin{array}{ccc} 1_a \bullet u & \xrightarrow{\blacksquare\text{-1-L}(u)} & u \\ \text{whisk-L}(1_a, p) \Big| & & \Big| p \\ 1_a \bullet v & \xrightarrow{\blacksquare\text{-1-L}(v)} & v \end{array}$$

Lemma 6. *Concatenation on the right by reflexivity is natural: for any points $b, c : A$, 1-paths $x, y : b = c$, and 2-path $q : x = y$, we have a term*

$$\blacksquare\text{-}1\text{-}R\text{-}nat(q) : whisk\text{-}R(q, 1_c) \cdot \blacksquare\text{-}1\text{-}R(y) = \blacksquare\text{-}1\text{-}R(x) \cdot q$$

Pictorially:

$$\begin{array}{ccc} x \cdot 1_c & \xrightarrow{\blacksquare\text{-}1\text{-}R(x)} & x \\ whisk\text{-}R(q, 1_c) \Big\downarrow & & \Big\downarrow q \\ y \cdot 1_c & \xrightarrow{\blacksquare\text{-}1\text{-}R(y)} & y \end{array}$$

Instantiating to the case $p, q : 1_\star = 1_\star$, we get terms

$$\begin{aligned} \blacksquare\text{-}1\text{-}L\text{-}nat(p) &: whisk\text{-}L(1_\star, p) \cdot 1_{1_\star} = 1_{1_\star} \cdot p \\ \blacksquare\text{-}1\text{-}R\text{-}nat(q) &: whisk\text{-}R(q, 1_\star) \cdot 1_{1_\star} = 1_{1_\star} \cdot q \end{aligned}$$

This clearly gives us $whisk\text{-}L(1_\star, p) = p$ and $whisk\text{-}R(q, 1_\star) = q$ as desired.

Squashing a commuting square whose opposing sides are reflexivities into an equality is so common that it will be useful to introduce a shorthand for this:

Lemma 7. *For any points $a, b : A$ and paths $p, q : a = b$, we have an equivalence*

$$\Downarrow : (p \cdot 1_b = 1_\star \cdot q) \simeq (p = q)$$

Lemma 8. *For any points $a, b : A$ and paths $p, q : a = b$, we have an equivalence*

$$\Rightarrow : (1_\star \cdot p = q \cdot 1_b) \simeq (p = q)$$

We can now formally prove Eckmann-Hilton:

Theorem 1 (Eckmann-Hilton). *For any point $\star : A$ and 2-loops $p, q : 1_\star = 1_\star$, we have a 3-path*

$$EH(p, q) : p \cdot q = q \cdot p$$

defined as the composition of the following three paths:

$$\begin{array}{c} p \cdot q \\ \Big\downarrow \left(\Downarrow (\blacksquare\text{-}1\text{-}L\text{-}nat(p)) \cdot \Downarrow (\blacksquare\text{-}1\text{-}R\text{-}nat(q)) \right)^{-1} \\ whisk\text{-}L(1_\star, p) \cdot whisk\text{-}R(q, 1_\star) \\ \Big\downarrow whisk\text{-}L\text{-}R(p, q) \\ whisk\text{-}R(q, 1_\star) \cdot whisk\text{-}L(1_\star, p) \\ \Big\downarrow \Downarrow (\blacksquare\text{-}1\text{-}R\text{-}nat(q)) \cdot \Downarrow (\blacksquare\text{-}1\text{-}L\text{-}nat(p)) \\ q \cdot p \end{array}$$

2 Eckmann-Hilton on Reflexivity

Since we want to prove something *about* the Eckmann-Hilton proof, and we intend to use path induction to do so, it is unsurprising that we will need to know how the EH proof behaves on identity paths. When both loops are reflexivities, the Eckmann-Hilton proof is very simple: $\text{EH}(1_{1_\star}, 1_{1_\star})$ automatically reduces to 1_{1_\star} . If one of the loops is a reflexivity, say \mathfrak{p} , there is an obvious candidate for $\text{EH}(1_{1_\star}, \mathfrak{q})$, namely

$$1_{1_\star} \cdot \mathfrak{q} \xrightarrow{\blacksquare\text{-}1\text{-}L(\mathfrak{q})} \mathfrak{q} \xrightarrow{\blacksquare\text{-}1\text{-}R(\mathfrak{q})^{-1}} \mathfrak{q} \cdot 1_{1_\star}$$

It is tempting (and correct) to instead use the path $\rightrightarrows^{-1}(1_{\mathfrak{q}})$ here. However, this may or may not give us what we want, depending on how exactly we define the equality \rightrightarrows . In one obvious implementation, $\rightrightarrows^{-1}(1_{\mathfrak{q}})$ reduces to

$$(\blacksquare\text{-}1\text{-}L(\mathfrak{q}) \cdot 1_{\mathfrak{q}}) \cdot \blacksquare\text{-}1\text{-}R(\mathfrak{q})^{-1}$$

and this, while equal to the path above, has a gratuitous $1_{\mathfrak{q}}$ term floating around. This would result in some extra work later on and extra work is best avoided.

So we want to construct a term

$$\text{EH-}1\text{-}L(\mathfrak{q}) : \text{EH}(1_{1_\star}, \mathfrak{q}) = \blacksquare\text{-}1\text{-}L(\mathfrak{q}) \cdot \blacksquare\text{-}1\text{-}R(\mathfrak{q})^{-1}$$

We cannot of course use induction on \mathfrak{q} right away. But we can work towards generalizing the situation until we can. The term $\text{EH}(1_{1_\star}, \mathfrak{q})$ reduces to the following composition:

$$\begin{array}{c} 1_{1_\star} \cdot \mathfrak{q} \\ \Big| \\ \left(1_{1_{1_\star}} \cdot \Downarrow (\blacksquare\text{-}1\text{-}R\text{-}nat(\mathfrak{q}))\right)^{-1} \\ \Big| \\ 1_{1_\star} \cdot \text{whisk-R}(\mathfrak{q}, 1_{1_\star}) \\ \Big| \\ \text{whisk-L-R}(1_{1_\star}, \mathfrak{q}) \\ \Big| \\ \text{whisk-R}(\mathfrak{q}, 1_{1_\star}) \cdot 1_{1_\star} \\ \Big| \\ \Downarrow (\blacksquare\text{-}1\text{-}R\text{-}nat(\mathfrak{q})) \cdot 1_{1_{1_\star}} \\ \Big| \\ \mathfrak{q} \cdot 1_{1_\star} \end{array}$$

Since whisk-L-R was defined by induction on both arguments, we first need to figure out what it does when only one of the arguments is reflexivity. Fortunately, this is very easy since we can just perform induction on the remaining argument:

Lemma 9. *For any point $a : A$, 1-paths $u : a = b$, $x, y : b = c$, and 2-path $q : x = y$, we have a term*

$$\text{whisk-L-R-}1\text{-}L(\mathfrak{q}) : \text{whisk-L-R}(1_u, \mathfrak{q}) = (\Downarrow^{-1}(1_{\text{whisk-L}(u, \mathfrak{q})}))$$

where the two paths in question witness the commutativity of the diagram below:

$$\begin{array}{ccc} u \cdot x & \xrightarrow{1_{u \cdot x}} & u \cdot x \\ \text{whisk-L}(u, \mathfrak{q}) \Big| & & \Big| \text{whisk-L}(u, \mathfrak{q}) \\ u \cdot y & \xrightarrow{1_{u \cdot y}} & u \cdot y \end{array}$$

Lemma 10. For any points $a, b, c : A$, 1-paths $u, v : a = b$, $x : b = c$, and 2-path $p : u = v$, we have a term

$$\mathit{whisk-L-R-1-R}(p) : \mathit{whisk-L-R}(p, 1_x) = (\Rightarrow^{-1} (1_{\mathit{whisk-R}(p,x)}))$$

where the two paths in question witness the commutativity of the diagram below:

$$\begin{array}{ccc} & \mathit{whisk-R}(p, x) & \\ u \cdot x & \xrightarrow{\quad} & v \cdot x \\ \left| 1_{u \cdot x} \right. & & \left. 1_{v \cdot x} \right| \\ u \cdot x & \xrightarrow{\quad \mathit{whisk-R}(p, x) \quad} & v \cdot x \end{array}$$

We can thus replace the middle segment $\mathit{whisk-L-R}(1_{1_\star}, q)$ in $\text{EH}(1_{1_\star}, q)$ as follows:

$$\begin{array}{c} 1_{1_\star} \cdot q \\ \left| \left(1_{1_{1_\star}} \cdot \cdot \Downarrow (\blacksquare\text{-1-R-nat}(q)) \right)^{-1} \right. \\ 1_{1_\star} \cdot \mathit{whisk-R}(q, 1_\star) \\ \left| \Rightarrow^{-1} (1_{\mathit{whisk-R}(q, 1_\star)}) \right. \\ \mathit{whisk-R}(q, 1_\star) \cdot 1_{1_\star} \\ \left| \Downarrow (\blacksquare\text{-1-R-nat}(q)) \cdot \cdot 1_{1_{1_\star}} \right. \\ q \cdot 1_{1_\star} \end{array}$$

Looking at the three segments, we see that it is possible to replace the diagram

$$\blacksquare\text{-1-R-nat}(q) : \mathit{whisk-R}(q, 1_\star) \cdot 1_{1_\star} = 1_{1_\star} \cdot q$$

with an abstract $\theta : \mathit{whisk-R}(q, 1_\star) \cdot 1_{1_\star} = 1_{1_\star} \cdot q$ of the same type:

$$\begin{array}{c} 1_{1_\star} \cdot q \\ \left| \left(1_{1_{1_\star}} \cdot \cdot \Downarrow (\theta) \right)^{-1} \right. \\ 1_{1_\star} \cdot \mathit{whisk-R}(q, 1_\star) \\ \left| \Rightarrow^{-1} (1_{\mathit{whisk-R}(q, 1_\star)}) \right. \\ \mathit{whisk-R}(q, 1_\star) \cdot 1_{1_\star} \\ \left| \Downarrow (\theta) \cdot \cdot 1_{1_{1_\star}} \right. \\ q \cdot 1_{1_\star} \end{array}$$

Having a path $\mathit{whisk-R}(q, 1_\star) \cdot 1_{1_\star} = 1_{1_\star} \cdot q$ is equivalent to having a path $\mathit{whisk-R}(q, 1_\star) = q$. If we managed to free the endpoints of the latter, we would be able to dispose of it by path induction. Fortunately, there is nothing that prevents us from doing so: we can just assume abstract loops $p, q : 1_\star = 1_\star$ in lieu of $\mathit{whisk-R}(q, 1_\star)$ and q , respectively. Our goal now becomes to prove that given $\theta : p \cdot 1_{1_\star} = 1_{1_\star} \cdot q$, the path

$$\begin{array}{c}
1_{1_\star} \cdot q \\
\left| (1_{1_\star} \cdot \Downarrow (\theta))^{-1} \right. \\
1_{1_\star} \cdot p \\
\left| \Rightarrow^{-1} (1_p) \right. \\
p \cdot 1_{1_\star} \\
\left| \Downarrow (\theta) \cdot 1_{1_\star} \right. \\
q \cdot 1_{1_\star}
\end{array}$$

equals $\blacksquare\text{-1-L}(q) \cdot \blacksquare\text{-1-R}(q)^{-1}$. This is already looking good but we can do even better: there is no longer anything in our goal that would require p and q to be loops, or even 2-paths! So we can fully generalize our goal: given points $a, b : A$, 1-paths $p, q : a = b$, and a 2-path $\theta : p \cdot 1_{1_\star} = 1_{1_\star} \cdot q$, the path

$$\begin{array}{c}
1_\star \cdot q \\
\left| (1_\star \cdot \Downarrow (\theta))^{-1} \right. \\
1_\star \cdot p \\
\left| \Rightarrow^{-1} (1_p) \right. \\
p \cdot 1_b \\
\left| \Downarrow (\theta) \cdot 1_b \right. \\
q \cdot 1_b
\end{array}$$

equals $\blacksquare\text{-1-L}(q) \cdot \blacksquare\text{-1-R}(q)^{-1}$. We can now convert the type of θ to the equivalent $p = q$: given points $a, b : A$, 1-paths $p, q : a = b$, and a 2-path $\theta : p = q$, the path

$$\begin{array}{c}
1_\star \cdot q \\
\left| (1_\star \cdot \theta)^{-1} \right. \\
1_\star \cdot p \\
\left| \Rightarrow^{-1} (1_p) \right. \\
p \cdot 1_b \\
\left| \theta \cdot 1_b \right. \\
q \cdot 1_b
\end{array}$$

equals $\blacksquare\text{-1-L}(q) \cdot \blacksquare\text{-1-R}(q)^{-1}$. But this is now trivial to prove: we first do induction on θ , which collapses p and q , and subsequently we do induction on p , which collapses a and b , and reduces everything to reflexivity.

Analogously, we construct a term

$$EH\text{-1-R}(p) : EH(p, 1_{1_\star}) = \blacksquare\text{-1-R}(p) \cdot \blacksquare\text{-1-L}(p)^{-1}$$

Our verbose explanation notwithstanding, the entire argument in this section can be coded up in just a few lines of Coq code, the most complex of which are the theorem statements themselves. As a further important bonus of formalization, we get that $EH\text{-1-L}(1_{1_\star})$ and $EH\text{-1-R}(1_{1_\star})$ reduce to 1_{1_\star} . It is of course also easy, albeit less convincing, to verify this explicitly by carefully examining the construction we gave.

3 Naturality of Eckmann-Hilton

As with all constructions in homotopy type theory, the Eckmann-Hilton proof *itself* respects equality:

Lemma 11. *For any 2-loops $u, v, x : 1_\star = 1_\star$, and 3-path $q : u = v$, we have a term*

$$EH\text{-L-nat}(q, x) : \text{whisk-R}(q, x) \cdot EH(v, x) = EH(u, x) \cdot \text{whisk-L}(x, q)$$

Pictorially:

$$\begin{array}{ccc} u \cdot x & \xrightarrow{EH(u, x)} & x \cdot u \\ \text{whisk-R}(q, x) \Big\downarrow & & \Big\downarrow \text{whisk-L}(x, q) \\ v \cdot x & \xrightarrow{EH(v, x)} & x \cdot v \end{array}$$

Lemma 12. *For any 2-loops $u, x, y : 1_\star = 1_\star$, and 3-path $p : x = y$, we have a term*

$$EH\text{-R-nat}(u, p) : \text{whisk-L}(u, p) \cdot EH(u, y) = EH(u, x) \cdot \text{whisk-R}(p, u)$$

Pictorially:

$$\begin{array}{ccc} u \cdot x & \xrightarrow{EH(u, x)} & x \cdot u \\ \text{whisk-L}(u, p) \Big\downarrow & & \Big\downarrow \text{whisk-R}(p, u) \\ u \cdot y & \xrightarrow{EH(u, y)} & y \cdot u \end{array}$$

In our case all 2-loops are reflexivities 1_{1_\star} . The terms $EH\text{-L-nat}(q, 1_{1_\star})$ and $EH\text{-R-nat}(1_{1_\star}, p)$, however, are not going to compute directly, as p and q are nontrivial and even loops. As we now show, however, this is not a problem - it turns out that both $EH\text{-L-nat}(q, 1_{1_\star})$ and $EH\text{-R-nat}(1_{1_\star}, p)$ can be constructed explicitly by pasting together a few commutative squares! So why even bother with an inductive definition? The key is precisely the word *inductive* - replacing the explicit construction we give below by the inductive definition we gave above effectively amounts to an abstraction that lies at the very heart of the entire proof of syllepsis.

To this end, we recall the following standard constructions on commutative squares:

Lemma 13. *For any points $a, b, c, d, e, f : A$, 1-paths $p : a = b$, $q : b = c$, $r : d = e$, $s : e = f$, $u : a = d$, $v : b = e$, $w : c = f$, and 2-paths $\gamma : p \cdot v = u \cdot r$, $\delta : q \cdot w = v \cdot s$ as in the diagram*

$$\begin{array}{ccc} a & \xrightarrow{u} & d \\ p \Big\downarrow & \gamma & \Big\downarrow r \\ b & \xrightarrow{v} & e \\ q \Big\downarrow & \delta & \Big\downarrow s \\ c & \xrightarrow{w} & f \end{array}$$

we have a term

$$\gamma \boxminus \delta : (p \cdot q) \cdot w = u \cdot (r \cdot s)$$

Lemma 14. For any points $a, b, c, d, e, f : A$, 1-paths $p : a = b$, $q : b = c$, $r : d = e$, $s : e = f$, $u : a = d$, $v : b = e$, $w : c = f$, and 2-paths $\gamma : u \cdot r = p \cdot v$, $\delta : v \cdot s = q \cdot w$ as in the diagram

$$\begin{array}{ccccc}
 a & \xrightarrow{p} & b & \xrightarrow{q} & c \\
 u \downarrow & & \gamma & & v \downarrow & & \delta & & w \downarrow \\
 d & \xrightarrow{r} & e & \xrightarrow{s} & f
 \end{array}$$

we have a term

$$\gamma \square \delta : u \cdot (r \cdot s) = (p \cdot q) \cdot w$$

Lemma 15. For any points $a, b, c, d : A$, 1-paths $p : a = b$, $q : c = d$, $r : a = c$, $s : b = d$, and 2-path $\gamma : p \cdot s = r \cdot q$ as in the diagram

$$\begin{array}{ccc}
 a & \xrightarrow{r} & c \\
 p \downarrow & & \gamma & & q \downarrow \\
 b & \xrightarrow{s} & d
 \end{array}$$

we have a term

$$\gamma^\uparrow : p^{-1} \cdot r = s \cdot q^{-1}$$

We also recall that the higher path $\text{EH-L-nat}(q, 1_{1_*})$ witnesses the commutativity of the following square:

$$\begin{array}{ccc}
 1_{1_*} \cdot 1_{1_*} & \xrightarrow{\text{EH}(1_{1_*}, 1_{1_*})} & 1_{1_*} \cdot 1_{1_*} \\
 \text{whisk-R}(q, 1_{1_*}) \downarrow & & \text{EH-L-nat}(q, 1_{1_*}) & & \text{whisk-L}(1_{1_*}, q) \downarrow \\
 1_{1_*} \cdot 1_{1_*} & \xrightarrow{\text{EH}(1_{1_*}, 1_{1_*})} & 1_{1_*} \cdot 1_{1_*}
 \end{array}$$

There is another way to fill this square - we horizontally compose $\blacksquare\text{-1-R-nat}(q)$ with the horizontal inverse of $\blacksquare\text{-1-L-nat}(q)$:

$$\begin{array}{ccccc}
 1_{1_*} \cdot 1_{1_*} & \xrightarrow{\blacksquare\text{-1-R}(1_{1_*})} & 1_{1_*} & \xrightarrow{\blacksquare\text{-1-L}(1_{1_*})^{-1}} & 1_{1_*} \cdot 1_{1_*} \\
 \text{whisk-R}(q, 1_{1_*}) \downarrow & & \blacksquare\text{-1-R-nat}(q) & & q & & \blacksquare\text{-1-L-nat}(q)^{\Leftarrow} & & \text{whisk-L}(1_{1_*}, q) \downarrow \\
 1_{1_*} \cdot 1_{1_*} & \xrightarrow{\blacksquare\text{-1-R}(1_{1_*})} & 1_{1_*} & \xrightarrow{\blacksquare\text{-1-L}(1_{1_*})^{-1}} & 1_{1_*} \cdot 1_{1_*}
 \end{array}$$

This indeed works because $\text{EH}(1_{1_*}, 1_{1_*})$, $\blacksquare\text{-1-R}(1_{1_*})$, and $\blacksquare\text{-1-L}(1_{1_*})$ are all definitionally 1_{1_*} . We now show that the two higher paths filling the square are equal too:

$$\text{EH-L-nat}(q, 1_{1_*}) = \blacksquare\text{-1-R-nat}(q) \square \blacksquare\text{-1-L-nat}(q)^{\Leftarrow}$$

We would like to prove this by induction on q ; in fact we don't have much choice since $\text{EH-L-nat}(q, 1_{1_*})$ is *defined* by induction on q . But, as mentioned above, q is a loop. So we need to free at least one of its endpoints. This is not so easy, however: the entire reason why the above equality even type-checks is that both endpoints of q are themselves reflexivities.

So we need to do some adjustment in our goal. Suppose we free the right endpoint, *i.e.*, we replace \mathfrak{q} by an abstract $q : 1_{1_\star} = y$. We could also try to free both endpoints, of course, to get $q : x = y$ but the one-sided version has a distinct advantage: after performing induction on q , *both* q and y reduce to reflexivity. The more reflexivities, the more things compute, so we go for this option.

Comparing the two sides of our goal, the left-hand side now witnesses the commutativity of the following square:

$$\begin{array}{ccc}
 1_{1_\star} \cdot 1_{1_\star} & \xrightarrow{EH(1_{1_\star}, 1_{1_\star})} & 1_{1_\star} \cdot 1_{1_\star} \\
 \text{whisk-R}(q, 1_{1_\star}) \Big\downarrow & EH\text{-L-nat}(q, 1_{1_\star}) & \Big\downarrow \text{whisk-L}(1_{1_\star}, q) \\
 1_{1_\star} \cdot y & \xrightarrow{EH(y, 1_{1_\star})} & 1_{1_\star} \cdot y
 \end{array}$$

On the right-hand side we have the following pasting of squares:

$$\begin{array}{ccccc}
 1_{1_\star} \cdot 1_{1_\star} & \xrightarrow{\blacksquare\text{-I-R}(1_{1_\star})} & 1_{1_\star} & \xrightarrow{\blacksquare\text{-I-L}(1_{1_\star})^{-1}} & 1_{1_\star} \cdot 1_{1_\star} \\
 \text{whisk-R}(q, 1_{1_\star}) \Big\downarrow & \blacksquare\text{-I-R-nat}(q) & q \Big\downarrow & \blacksquare\text{-I-L-nat}(q)^{\Leftarrow} & \Big\downarrow \text{whisk-L}(1_{1_\star}, q) \\
 1_{1_\star} \cdot 1_{1_\star} & \xrightarrow{\blacksquare\text{-I-R}(y)} & y & \xrightarrow{\blacksquare\text{-I-L}(y)^{-1}} & y \cdot 1_{1_\star}
 \end{array}$$

As we can see, the two diagrams are no longer identical - the former has $EH(y, 1_{1_\star})$ on the bottom and the latter has $\blacksquare\text{-I-R}(y) \cdot \blacksquare\text{-I-L}(y)^{-1}$ instead. But not all hope is lost! In the previous section, we constructed a higher path for precisely such an occasion, namely

$$EH\text{-I-R}(y) : EH(y, 1_{1_\star}) = \blacksquare\text{-I-R}(y) \cdot \blacksquare\text{-I-L}(y)^{-1}$$

So all we need to do is to adjust our goal as follows:

$$\begin{aligned}
 EH\text{-L-nat}(q, 1_{1_\star}) &= \text{whisk-L}(\text{whisk-R}(q, 1_{1_\star}), EH\text{-I-R}(y)) \cdot \\
 &\quad (\blacksquare\text{-I-L-nat}(q) \square \blacksquare\text{-I-L-nat}(q)^{\Leftarrow})
 \end{aligned}$$

This is now easy to show by induction on q . Specializing to the case of interest $\mathfrak{q} : 1_{1_\star} = 1_{1_\star}$ yields the desired equality

$$EH\text{-L-nat}(q, 1_{1_\star}) = \blacksquare\text{-I-R-nat}(q) \square \blacksquare\text{-I-L-nat}(q)^{\Leftarrow}$$

since in this case the path $EH\text{-I-R}(1_{1_\star})$ reduces to $1_{1_{1_\star}}$, as remarked in the previous section. Entirely analogously, we have

$$EH\text{-R-nat}(1_{1_\star}, p) = \blacksquare\text{-I-L-nat}(p) \square \blacksquare\text{-I-L-nat}(p)^{\Leftarrow}$$

4 Syllepsis

We can now proceed with our proof of syllepsis. Formally, we wish to show the following:

Theorem 2 (Syllepsis). *For any point $\star : A$ and 3-loops $p, q : 1_{1_\star} = 1_{1_\star}$, we have*

$$EH(q, p) = EH(q, p)^{-1}$$

Pictorially:

$$\begin{array}{ccc}
& p \cdot q & \\
\left(\Downarrow (\blacksquare -1-L\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(q)) \right)^{-1} & \swarrow & \searrow \left(\Downarrow (\blacksquare -1-R\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(q)) \right)^{-1} \\
whisk-L(1_{1_*}, p) \cdot whisk-R(q, 1_{1_*}) & & whisk-R(p, 1_{1_*}) \cdot whisk-L(1_{1_*}, q) \\
\downarrow whisk-L-R(p, q) & & \downarrow whisk-L-R(q, p)^{-1} \\
whisk-R(q, 1_{1_*}) \cdot whisk-L(1_{1_*}, p) & & whisk-L(1_{1_*}, q) \cdot whisk-R(p, 1_{1_*}) \\
\Downarrow (\blacksquare -1-R\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(p)) & \searrow & \swarrow \Downarrow (\blacksquare -1-L\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(p)) \\
& q \cdot p &
\end{array}$$

Step 1 The natural first step towards proving syllepsis is to split the hexagon into two triangles and a square:

$$\begin{array}{ccc}
& p \cdot q & \\
\left(\Downarrow (\blacksquare -1-L\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(q)) \right)^{-1} & \swarrow & \searrow \left(\Downarrow (\blacksquare -1-R\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(q)) \right)^{-1} \\
whisk-L(1_{1_*}, p) \cdot whisk-R(q, 1_{1_*}) & \xrightarrow{??} & whisk-R(p, 1_{1_*}) \cdot whisk-L(1_{1_*}, q) \\
\downarrow whisk-L-R(p, q) & & \downarrow whisk-L-R(q, p)^{-1} \\
whisk-R(q, 1_{1_*}) \cdot whisk-L(1_{1_*}, p) & \xrightarrow{??} & whisk-L(1_{1_*}, q) \cdot whisk-R(p, 1_{1_*}) \\
\Downarrow (\blacksquare -1-R\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(p)) & \searrow & \swarrow \Downarrow (\blacksquare -1-L\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(p)) \\
& q \cdot p &
\end{array}$$

We would like to prove the commutativity of the square in the middle by induction on p and q . We thus want to generalize the situation so that we have $p : x = y$ and $q : u = v$ for arbitrary 2-loops $x, y, u, v : 1_* = 1_*$. The obvious attempt at this leads us to the following:

$$\begin{array}{ccc}
whisk-L(u, p) \cdot whisk-R(q, y) & \xrightarrow{??} & whisk-R(p, u) \cdot whisk-L(y, q) \\
\downarrow whisk-L-R(p, q) & & \downarrow whisk-L-R(q, p)^{-1} \\
whisk-R(q, x) \cdot whisk-L(v, p) & \xrightarrow{??} & whisk-L(x, q) \cdot whisk-R(p, v)
\end{array}$$

However, the vertices on the left are paths $u \cdot x = v \cdot y$ whereas those on the right are paths $x \cdot u = y \cdot v$. To make the endpoints align, we insert an Eckmann-Hilton proof on both sides. We are able to do this precisely because the p and q we started with were 3-loops; syllepsis does *not* hold if we go one dimension lower (this observation is due to Jamie Vicary).

Our square now has the following form, where the vertical paths simply leave the EH term unchanged:

$$\begin{array}{ccc}
(whisk-L(u, p) \cdot whisk-R(q, y)) \cdot EH(v, y) & \xrightarrow{??} & EH(u, x) \cdot (whisk-R(p, u) \cdot whisk-L(y, q)) \\
\downarrow whisk-L-R(p, q) \cdot \cdot 1_{EH(v, y)} & & \downarrow 1_{EH(u, x)} \cdot \cdot whisk-L-R(q, p)^{-1} \\
(whisk-R(q, x) \cdot whisk-L(v, p)) \cdot EH(v, y) & \xrightarrow{??} & EH(u, x) \cdot (whisk-L(x, q) \cdot whisk-R(p, v))
\end{array}$$

Before we fill the horizontal paths and the square itself, we check that we have not lost touch with what we originally set out to prove: in the special case when all of the 2-paths are reflexivities, the EH terms $\text{EH}(u, x)$ and $\text{EH}(v, y)$ reduce to $1_{1_{1_\star}}$, as observed in the previous section. So the square we have is not *exactly* the one we wanted - all the vertices now contain an extra reflexivity path - but it is close enough.

To construct the horizontal paths, we need to fill the following two diagrams:

$$\begin{array}{ccc}
& \text{EH}(u, x) & \\
u \cdot x & \xrightarrow{\quad} & x \cdot u \\
\text{whisk-L}(u, p) \Big| & & \Big| \text{whisk-R}(p, u) \\
& & \\
u \cdot y & & y \cdot u \\
\text{whisk-R}(q, y) \Big| & & \Big| \text{whisk-L}(y, q) \\
& & \\
v \cdot y & \xrightarrow{\quad \text{EH}(v, y) \quad} & y \cdot v
\end{array}
\qquad
\begin{array}{ccc}
& \text{EH}(u, x) & \\
u \cdot x & \xrightarrow{\quad} & x \cdot u \\
\text{whisk-R}(q, x) \Big| & & \Big| \text{whisk-L}(x, q) \\
& & \\
v \cdot x & & x \cdot v \\
\text{whisk-L}(v, p) \Big| & & \Big| \text{whisk-R}(p, v) \\
& & \\
v \cdot y & \xrightarrow{\quad \text{EH}(v, y) \quad} & y \cdot v
\end{array}$$

The obvious way to do this is to split each diagram into two squares as follows:

$$\begin{array}{ccc}
& \text{EH}(u, x) & \\
u \cdot x & \xrightarrow{\quad} & x \cdot u \\
\text{whisk-L}(u, p) \Big| & & \Big| \text{whisk-R}(p, u) \\
& & \\
u \cdot y & \xrightarrow{\quad \text{EH}(u, y) \quad} & y \cdot u \\
\text{whisk-R}(q, y) \Big| & & \Big| \text{whisk-L}(y, q) \\
& & \\
v \cdot y & \xrightarrow{\quad \text{EH}(v, y) \quad} & y \cdot v
\end{array}
\qquad
\begin{array}{ccc}
& \text{EH}(u, x) & \\
u \cdot x & \xrightarrow{\quad} & x \cdot u \\
\text{whisk-R}(q, x) \Big| & & \Big| \text{whisk-L}(x, q) \\
& & \\
v \cdot x & \xrightarrow{\quad \text{EH}(v, x) \quad} & x \cdot v \\
\text{whisk-L}(v, p) \Big| & & \Big| \text{whisk-R}(p, v) \\
& & \\
v \cdot y & \xrightarrow{\quad \text{EH}(v, y) \quad} & y \cdot v
\end{array}$$

Each small square now commutes by the naturality of Eckmann-Hilton, so we can fill the horizontal paths in our big square as follows:

$$\begin{array}{ccc}
(\text{whisk-L}(u, p) \cdot \text{whisk-R}(q, y)) \cdot \text{EH}(v, y) & \xrightarrow{\text{EH-R-nat}(u, p) \boxminus \text{EH-L-nat}(q, y)} & \text{EH}(u, x) \cdot (\text{whisk-R}(p, u) \cdot \text{whisk-L}(y, q)) \\
\text{whisk-L-R}(p, q) \cdot \cdot 1_{\text{EH}(v, y)} \Big| & & \Big| 1_{\text{EH}(u, x)} \cdot \cdot \text{whisk-L-R}(q, p)^{-1} \\
(\text{whisk-R}(q, x) \cdot \text{whisk-L}(v, p)) \cdot \text{EH}(v, y) & \xrightarrow{\text{EH-L-nat}(q, x) \boxminus \text{EH-R-nat}(v, p)} & \text{EH}(u, x) \cdot (\text{whisk-L}(x, q) \cdot \text{whisk-R}(p, v))
\end{array}$$

The above commutes by induction on p and q . Specializing the generalization $p : x = y$ and $q : u = v$ to our situation yields the following commuting square:

$$\begin{array}{ccc}
(\text{whisk-L}(1_{1_\star}, p) \cdot \text{whisk-R}(q, 1_{1_\star})) \cdot 1_{1_{1_\star}} & \xrightarrow{\text{EH-R-nat}(1_{1_\star}, p) \boxminus \text{EH-L-nat}(q, 1_{1_\star})} & 1_{1_{1_\star}} \cdot (\text{whisk-R}(p, 1_{1_\star}) \cdot \text{whisk-L}(1_{1_\star}, q)) \\
\text{whisk-L-R}(p, q) \cdot \cdot 1_{1_{1_\star}} \Big| & & \Big| 1_{1_{1_\star}} \cdot \cdot \text{whisk-L-R}(q, p)^{-1} \\
(\text{whisk-R}(q, 1_{1_\star}) \cdot \text{whisk-L}(1_{1_\star}, p)) \cdot 1_{1_{1_\star}} & \xrightarrow{\text{EH-L-nat}(q, 1_{1_\star}) \boxminus \text{EH-R-nat}(1_{1_\star}, p)} & 1_{1_{1_\star}} \cdot (\text{whisk-L}(1_{1_\star}, q) \cdot \text{whisk-R}(p, 1_{1_\star}))
\end{array}$$

Step 2 We now want to somehow retrofit the square from the previous step into the original diagram for syllepsis. Of course we cannot do this quite literally, since our square has an extra reflexivity path in each vertex. But we can fit the following triangles into the upper and lower part of the hexagon, respectively:

$$\begin{array}{ccc}
& p \cdot q & \\
\left(\Downarrow (\blacksquare -1-L\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(q)) \right)^{-1} & & \left(\Downarrow (\blacksquare -1-R\text{-nat}(p)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(q)) \right)^{-1} \\
& \swarrow & \searrow \\
\text{whisk-L}(1_{1_*}, p) \cdot \text{whisk-R}(q, 1_{1_*}) & \xrightarrow{\quad} & \text{whisk-R}(p, 1_{1_*}) \cdot \text{whisk-L}(1_{1_*}, q) \\
& \Downarrow (EH-R\text{-nat}(1_{1_*}, p) \boxminus EH-L\text{-nat}(q, 1_{1_*})) & \\
& \xrightarrow{\quad} & \\
& \text{whisk-R}(q, 1_{1_*}) \cdot \text{whisk-L}(1_{1_*}, p) & \xrightarrow{\quad} & \text{whisk-L}(1_{1_*}, q) \cdot \text{whisk-R}(p, 1_{1_*}) \\
& \Downarrow (EH-L\text{-nat}(q, 1_{1_*}) \boxminus EH-R\text{-nat}(1_{1_*}, p)) & & \\
& \swarrow & & \searrow \\
& q \cdot p & \\
\Downarrow (\blacksquare -1-R\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-L\text{-nat}(p)) & & \Downarrow (\blacksquare -1-L\text{-nat}(q)) \cdot \cdot \Downarrow (\blacksquare -1-R\text{-nat}(p))
\end{array}$$

If we show that these triangles do in fact commute, we will be (almost) done. To do so, we again wish to suitably generalize the situation. Starting with the first triangle, we can put together the four commuting squares $\blacksquare -1-L\text{-nat}(p)$, $\blacksquare -1-R\text{-nat}(p)$, $\blacksquare -1-L\text{-nat}(q)$, $\blacksquare -1-R\text{-nat}(q)$ as follows:

$$\begin{array}{ccccc}
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-L}(1_{1_*}, p) \Big| & \blacksquare -1-L\text{-nat}(p) & p & (\blacksquare -1-R\text{-nat}(p))^{\Leftarrow} & \text{whisk-R}(p, 1_{1_*}) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-R}(q, 1_{1_*}) \Big| & \blacksquare -1-R\text{-nat}(q) & q & (\blacksquare -1-L\text{-nat}(q))^{\Leftarrow} & \text{whisk-L}(1_{1_*}, q) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*}
\end{array}$$

On the other hand, we also have the two commuting squares $EH-L\text{-nat}(q, 1_{1_*})$ and $EH-R\text{-nat}(1_{1_*}, p)$:

$$\begin{array}{ccccc}
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-L}(1_{1_*}, p) \Big| & & EH-R\text{-nat}(1_{1_*}, p) & & \text{whisk-R}(p, 1_{1_*}) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-R}(q, 1_{1_*}) \Big| & & EH-L\text{-nat}(q, 1_{1_*}) & & \text{whisk-L}(1_{1_*}, q) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} & \xrightarrow{1_{1_*}} & 1_{1_*} \cdot 1_{1_*}
\end{array}$$

We have already established a relationship between these in a previous section:

$$\begin{aligned}
EH-R\text{-nat}(1_{1_*}, p) &= \blacksquare -1-L\text{-nat}(p) \boxplus \blacksquare -1-R\text{-nat}(p)^{\Leftarrow} \\
EH-L\text{-nat}(q, 1_{1_*}) &= \blacksquare -1-R\text{-nat}(q) \boxplus \blacksquare -1-L\text{-nat}(q)^{\Leftarrow}
\end{aligned}$$

Thus, we can abstractly summarize the entire situation in the following lemma:

Lemma 16. *Assume points $a, b, c : A$, 1-paths $p, q, r : a = b$, $u, v, w : b = c$, and 2-paths $\alpha : p \cdot 1_b = 1_a \cdot q$, $\beta : r \cdot 1_b = 1_a \cdot q$, $\gamma : u \cdot 1_c = 1_b \cdot v$, $\delta : w \cdot 1_c = 1_b \cdot v$ as in the diagram below:*

$$\begin{array}{ccccc}
 & a & \xrightarrow{1_a} & a & \xrightarrow{1_a} & a \\
 p & \downarrow & & \downarrow & & \downarrow & r \\
 & & \alpha & & \beta^{\Leftarrow} & & \\
 & b & \xrightarrow{1_b} & b & \xrightarrow{1_b} & b \\
 u & \downarrow & & \downarrow & & \downarrow & w \\
 & & \gamma & & \delta^{\Leftarrow} & & \\
 & c & \xrightarrow{1_c} & c & \xrightarrow{1_c} & c
 \end{array}$$

Furthermore, assume 2-paths $\theta : p \cdot 1_b = 1_a \cdot r$ and $\phi : u \cdot 1_c = 1_b \cdot w$ as in the diagram below:

$$\begin{array}{ccccc}
 & a & \xrightarrow{1_a} & a & \xrightarrow{1_a} & a \\
 p & \downarrow & & \downarrow & & \downarrow & r \\
 & & \theta & & & & \\
 & b & \xrightarrow{1_b} & b & \xrightarrow{1_b} & b \\
 u & \downarrow & & \downarrow & & \downarrow & w \\
 & & \phi & & & & \\
 & c & \xrightarrow{1_c} & c & \xrightarrow{1_c} & c
 \end{array}$$

If $\theta = \alpha \sqcup \beta^{\Leftarrow}$ and $\phi = \gamma \sqcup \delta^{\Leftarrow}$, then the following triangle commutes:

$$\begin{array}{ccc}
 & q \cdot v & \\
 (\Downarrow (\alpha) \cdot \Downarrow (\gamma))^{-1} \swarrow & & \searrow (\Downarrow (\beta) \cdot \Downarrow (\delta))^{-1} \\
 p \cdot u & \xrightarrow{\Downarrow (\theta \sqcup \phi)} & r \cdot w
 \end{array}$$

To prove the above lemma, we first perform path induction on the two hypotheses, thereby eliminating the 2-paths θ and ϕ . Next we reformulate the goal so that we can get rid of the remaining 2-paths by induction - given points $a, b, c : A$, 1-paths $p, q, r : a = b$, $u, v, w : b = c$, and 2-paths $\alpha : p = q$, $\beta : r = q$, $\gamma : u = v$, $\delta : w = v$, the following triangle commutes:

$$\begin{array}{ccc}
 & q \cdot v & \\
 (\Downarrow (\Downarrow^{-1} (\alpha)) \cdot \Downarrow (\Downarrow^{-1} (\gamma)))^{-1} \swarrow & & \searrow (\Downarrow (\Downarrow^{-1} (\beta)) \cdot \Downarrow (\Downarrow^{-1} (\delta)))^{-1} \\
 p \cdot u & \xrightarrow{\Downarrow ((\Downarrow^{-1} (\alpha) \sqcup (\Downarrow^{-1} (\beta))^{\Leftarrow}) \sqcup (\Downarrow^{-1} (\gamma) \sqcup (\Downarrow^{-1} (\delta))^{\Leftarrow}))} & r \cdot w
 \end{array}$$

Now we perform path induction on $\alpha, \beta, \gamma, \delta$, which also eliminates the 1-paths p, r, v, w . The only two remaining 1-paths are $p : a = b$ and $u : b = c$, and these beg for further path induction after which there is nothing left to do.

The lemma now immediately implies the commutativity of the upper triangle. In fact, it also implies the commutativity of the lower triangle - the four commuting squares \blacksquare -1-L-nat(p), \blacksquare -1-R-nat(p), \blacksquare -1-L-nat(q), \blacksquare -1-R-nat(q) can now be put together as follows:

$$\begin{array}{ccccc}
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-R}(q, 1_{1_*}) \Big| & \blacksquare\text{-1-R-nat}(q) & q & \blacksquare\text{-1-L-nat}(q)^{\Leftarrow} & \text{whisk-L}(1_{1_*}, q) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-L}(1_{1_*}, p) \Big| & \blacksquare\text{-1-L-nat}(p) & p & \blacksquare\text{-1-R-nat}(p)^{\Leftarrow} & \text{whisk-R}(p, 1_{1_*}) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*}
\end{array}$$

The two commuting squares $\text{EH-L-nat}(1_{1_*}, p)$ and $\text{EH-R-nat}(q, 1_{1_*})$ can be stacked as in the diagram below:

$$\begin{array}{ccccc}
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-R}(q, 1_{1_*}) \Big| & & \text{EH-L-nat}(q, 1_{1_*}) & & \text{whisk-L}(1_{1_*}, q) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*} \\
\text{whisk-L}(1_{1_*}, p) \Big| & & \text{EH-R-nat}(1_{1_*}, p) & & \text{whisk-R}(p, 1_{1_*}) \\
1_{1_*} \cdot 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} & \xrightarrow{1_{1_{1_*}}} & 1_{1_*} \cdot 1_{1_*}
\end{array}$$

And as before, we have

$$\begin{aligned}
\text{EH-L-nat}(q, 1_{1_*}) &= \blacksquare\text{-1-R-nat}(q) \square \blacksquare\text{-1-L-nat}(q)^{\Leftarrow} \\
\text{EH-R-nat}(1_{1_*}, p) &= \blacksquare\text{-1-L-nat}(p) \square \blacksquare\text{-1-R-nat}(p)^{\Leftarrow}
\end{aligned}$$


The same lemma thus implies the commutativity of the following triangle:

$$\begin{array}{ccc}
& q \cdot p & \\
\swarrow & & \searrow \\
\left(\Downarrow (\blacksquare\text{-1-R-nat}(q)) \cdot \Downarrow (\blacksquare\text{-1-L-nat}(p)) \right)^{-1} & & \left(\Downarrow (\blacksquare\text{-1-L-nat}(q)) \cdot \Downarrow (\blacksquare\text{-1-R-nat}(p)) \right)^{-1} \\
\text{whisk-R}(q, 1_{1_*}) \cdot \text{whisk-L}(1_{1_*}, p) & \xrightarrow{\Downarrow (\text{EH-L-nat}(q, 1_{1_*}) \boxplus \text{EH-R-nat}(1_{1_*}, p))} & \text{whisk-L}(1_{1_*}, q) \cdot \text{whisk-R}(p, 1_{1_*})
\end{array}$$


Flipping the triangle along the horizontal axis yields precisely the desired lower triangle, and we are done.

Step 3 It remains to somehow combine the two commuting triangles on the top and bottom with the commuting square in the middle. We cannot literally paste them together because, as we recall, the vertices of the square contain an extra reflexivity path originating from the Eckmann-Hilton term. But this is not a problem because we have a suitable generalization up our sleeve:

Lemma 17. *Assume points $a, b : A$, 1-paths $p, q, u, v, x, y : a = b$, and 2-paths $\alpha : u = p$, $\beta : v = p$, $\gamma : x = q$, $\delta : y = q$, $\theta : u = x$, $\phi : y = v$ as in the diagram below:*



Assume further 2-paths $\eta : u \cdot 1_b = 1_a \cdot v$ and $\varepsilon : x \cdot 1_b = 1_a \cdot y$. Then the hexagon above commutes provided the triangles and square below do:



To prove this lemma, we first perform path induction on $\theta, \phi, \beta, \delta$. This eliminates the 1-paths p, q, v, y . The commutative square hypothesis now becomes equivalent to $\eta = \varepsilon$, so we can perform induction and get rid of ε . The commutative triangle hypotheses now become equivalent to $\Downarrow (\eta) = \alpha$ and $\Downarrow (\eta) = \gamma$, so we can perform induction and get rid of α and γ . Among the 2-paths this only leaves $\eta : u \cdot 1_b = 1_a \cdot y$, which is equivalent to $\eta : u = y$, so we can perform induction to get rid of y . The sole remaining 1-path is $u : a = b$, and we perform one last induction on it. We have managed to reduce everything in sight to reflexivity and thus made the hexagon trivially commute.

The conclusion of the above lemma clearly implies syllepsis, and the previous two steps show that the hypotheses hold, so \checkmark .