

Engineering of Anyons on M5-Probes via Flux Quantization

Hisham Sati^{a,b}

hsati@nyu.edu

Urs Schreiber^a

us13@nyu.edu

March 13, 2025

Abstract

These extended lecture notes survey a novel derivation of anyonic topological order (as seen in fractional quantum Hall systems) on single magnetized M5-branes probing Seifert orbi-singularities (“geometric engineering” of anyons), which we motivate from fundamental open problems in the field of quantum computing.

The rigorous construction is non-Lagrangian and non-perturbative, based on previously neglected global completion of the M5-brane’s tensor field by flux-quantization consistent with its non-linear self-duality and its twisting by the bulk C-field. This exists only in little-studied non-abelian generalized cohomology theories, notably in a twisted equivariant (and “twistorial”) form of unstable Cohomotopy (“Hypothesis H”).

As a result, topological quantum observables form Pontrjagin homology algebras of mapping spaces from the orbi-fixed worldvolume into a classifying 2-sphere. Remarkably, results from algebraic topology imply from this the quantum observables and modular functor of abelian Chern–Simons theory, as well as braid group actions on defect anyons of the kind envisioned as hardware for topologically protected quantum gates.

prepared for the lecture series

Introduction to Hypothesis H

held at

45th Winter School GEOMETRY AND PHYSICS

Srní, Czechia (18-25 Jan 2024)

Contents

1	Motivation: Better Anyon Theory	2
2	Flux-Quantization on M5-Probes	7
3	Cohomotopy Charge of Solitons	15
4	The topological Quantum States	20
5	Conclusion: Better Anyon Theory	26
6	Digest for Algebraic Topologists	30
A	Background on Homotopy Theory	33
B	Background on TED Cohomotopy	35

^a Mathematics, Division of Science; and
Center for Quantum and Topological Systems,
NYUAD Research Institute,
New York University Abu Dhabi, UAE.

^b The Courant Institute for Mathematical Sciences, NYU, NY.




1 Motivation: Better Anyon Theory

While the hopes associated with the idea of *quantum computing* [102][58] are hard to over-state [51][10][116], there are good arguments that commercial-value quantum computing will ultimately require quantum hardware exhibiting *anyonic topological order* [160][129]. But microscopic theoretical derivations, from first principles, of such anyonic quantum states in strongly-coupled quantum systems had remained sketchy, which may explain the dearth of experimental realizations to date.

What we review here (based on [132][135][53]) is a rigorous theoretical account via “geometric engineering on M-branes” subject to a previously neglected step of “flux-quantization” (the latter surveyed in [130]).

First, we expand on the motivation a little further:

Ultimate need for Topological Quantum Protection. Despite the fascinating reality of presently available Noisy Intermediate-Scale Quantum computers (NISQ [114]) and despite the mid-term prospect of their stabilization at the software-level via Quantum Error Correction (QEC [87][115], at heavy cost of available system scale), serious arguments [75][30][85][31][32][67][50][146] and experience [22] suggest that large-scale quantum computation is hardly attainable by incremental optimization of NISQ architectures, but [23]¹ that more fundamental quantum principles will need to be exploited – notably *topological error protection* already at the hardware-level [80][46][137][136] in order to suppress quantum errors occurring in the first place.



While topological quantum protection is thus possibly indispensable for achieving commercial-value quantum computing, its ambitious development, in theory and practice, is in fact far from mature, is in need of new ideas and of further analysis, and leaves much room for development. Since this is not always made clear, to amplify this point:

(i) **Theoretical challenges:** While quantum theorists now routinely deal with the algebraic structure (namely: braided fusion categories) commonly *expected* [81] to describe interaction of anyon species *in toto*, the *microscopic* first-principles understanding of the formation of anyonic topological order as solitonic states in the many-body (electron) dynamics of quantum materials has remained at most sketchy, even in the best-understood case of the fractional quantum Hall effect [140], cf. [70].²

In fact, this is an instance of the general open problem of analytically establishing gapped bound states in any strongly coupled/correlated quantum system: The problem of formulating non-perturbative quantum field theory [6][34]. The analogous issue in particle physics (there called the *Yang-Mills mass gap* problem [103]) has been recognized as being profound enough to be declared one of seven “Millennium Problems” [20].

(ii) **Practical challenges:** But without a robust theoretical prediction of anyonic solitons in actual quantum materials, it remains unclear where and how to look for them. As an unfortunate result, experimentalists have turned attention to mere stand-ins, such as “Majorana zero modes” at the ends of super/semi-conducting nanowires ([79][92] which, even if the doubts about their detection were to be removed [24], are by construction immobile and hence do not serve as hardware-protected quantum braid gates) and quantum-simulation of anyons on NISQ architectures ([69][43, Fig. 5], which might serve as software-level QEC but again offers no hardware-level protection.

¹[23]: “The qubit systems we have today are a tremendous scientific achievement, but they take us no closer to having a quantum computer that can solve a problem that anybody cares about. [...] What is missing is the breakthrough [...] bypassing quantum error correction by using far-more-stable qubits, in an approach called topological quantum computing.”


²[70, p. 3]: “Though the Laughlin function very well approximates the true ground state at $\nu = 1/q$, the physical mechanism of related correlations and of the whole hierarchy of the FQHE remained, however, still obscure. [...] The so-called HH (Halperin–Haldane) model of consecutive generations of Laughlin states of anyonic quasiparticle excitations from the preceding Laughlin state has been abandoned early because of the rapid growth of the daughter quasiparticle size, which quickly exceeded the sample size. [...] the Halperin multicomponent theory and of the CF model advanced the understanding of correlations in FQHE, however, on phenomenological level only. CFs were assumed to be hypothetical quasi-particles consisting of electrons and flux quanta of an auxiliary fictitious magnetic field pinned to them. The origin of this field and the manner of attachment of its flux quanta to electrons have been neither explained nor discussed.”

In short: **Foundation and implementation** of topological quantum computing as a plausible long-term pathway to actual quantum value **deserves and admits thorough re-investigation**.

Concretely, the intrinsic tension haunting the traditional quantum computing paradigm is (cf. [17, p 272][146, p 3]) that:


- (i) quantum gates are *implemented via interaction* of subsystems,
- (ii) while quantum coherence requires *avoiding all interaction*.

The idea of topological protection is to cut this Gordian knot by *quantum gates operating without interaction*. The physical principle that allows this to work [3][4][46, p 6][108, p 50] is the *quantum adiabatic theorem* [118]: Gapped quantum systems frozen at absolute zero in one of several ground states, but dependent on external parameters, will defy interaction with noise quanta below the energy gap and yet have their ground state transformed by sufficiently gentle tuning of the parameters: a *holonomic quantum gate*. This is *topological* if it is invariant under local deformations of parameter paths, and thus protected also against classical noise. For an *anyonic braid gate* the parameters in question are the positions of defects in a 2-dimensional transverse space within a quantum material.



The remaining problem is to develop a precise mathematical theory describing these *anyons*.


Improved Anyon Models via Geometric Engineering on M-branes. A remarkable approach to the otherwise elusive microscopic analysis of such strongly-coupled/correlated quantum systems emerges in the guise of “geometric engineering” [78][13] of quantum fields on “M-branes” probing orbifold singularities, whereby the given dynamics is (partially) mapped onto the fluctuations of Membranes (whence *M-theory* [28]), and of higher-dimensional “M5-branes” [53], propagating within an auxiliary higher-dimensional gravitating spacetime orbifold [122].



This procedure is most famous in the (unrealistic) limit of large rank and hence of large numbers $N \rightarrow \infty$ of coincident such branes, where it extracts quantum correlators and quantum phase transitions entirely from classical gravitational asymptotics (“holographic duality” [1]). The application to quantum materials [159][63] is now well-studied, notably in the case of quantum critical superconductors engineered in M-theory [66][48][49][59][26][27][2].

But we have established [53][132][133][135] that after implementing a previously neglected step of “flux quantization” [130] on the M5-brane worldvolume, there provably appear general solitonic and specifically anyonic quantum states already in the more realistic situation of single ($N = 1$) coincident branes. (Similar results for $N = 2$ had previously only been conjectured [18] by appeal to the expected but notoriously undefined effective quantum field theory on coincident M5-branes.)

Brane diagram for geometric engineering of anyons on single M5-branes wrapping an orbifold singularity [135]: It is a subtle mechanism of *flux-quantization* [130] of the self-dual tensor-field on the M5 [53] that stabilizes [132] its anyonic soliton configurations.



Here, we review and explain how this works for an audience assumed to be familiar with the general mechanism of *flux quantization* as surveyed in [130].

But first to recall the traditional theory of fractional quantum Hall anyons:

Quantum Hall effect (cf. [113][15][140][109]). In a very thin (atomic multi-layer) and hence effectively 2-dimensional sheet Σ^2 of (semi-)conducting material carrying magnetic flux density B , the energy of electron states is (cf. [145, (4-12)]) quantized by *Landau levels* $i \in \mathbb{N}$ as

$$E = \hbar\omega_B \left(i + \frac{1}{2} \right),$$

where each level comprises of one state per magnetic flux quantum:

$$n_{\text{deg}} = B/\Phi_0;$$

the Lorentz force on a longitudinal electron current J_x at filling fraction ν is compensated in equilibrium by an electric *Hall field*

$$E_y = \frac{1}{\nu} J_x.$$

Integer quantum Hall effect. Therefore, Fermi’s theory of idealized *free* electrons predicts the system to be a conductor away from the energy gaps between a completely filled and the next empty Landau level, hence away from the number of electrons being integer multiples of the number of flux quanta, where longitudinal conductivity should vanish.


$$n_{\text{el}} = \nu B/\Phi_0, \quad \nu \in \mathbb{N}.$$

This is indeed observed — in fact, the vanishing conductivity is observed in sizeable neighborhoods of the critical filling fractions (“Hall plateaux”, attributed to subtle disorder effects).

Fractional quantum Hall effect (FQHE). But in reality, the electrons are far from free. While there is little theory for strongly interacting quantum systems, experiment shows that the Fermi idealization breaks down at low enough temperature, where longitudinal conductivity also decreases in neighborhoods of certain *fractional* filling factors ν .

$$\nu \in \mathbb{Q}, \quad \text{prominently for } \nu = 1/k, \quad k \in 2\mathbb{N} + 1.$$

The traditional heuristic idea is that at these filling fractions the interacting electrons each form a kind of bound state with k flux quanta, making “composite bosons” (cf. [161]) that, as such, condense to produce an insulating mass gap, even inside the Landau level.



Anyonic quasi-particles. This heuristic model suggests that in the Hall plateau neighborhood *around* such filling fraction, there are *unpaired* flux quanta effectively “bound to” $1/k$ th of a (missing) electron: called “quasi-particles” (“quasi-holes”). These quasi-particles/holes evidently have fractional charge $\pm e/k$ and are expected to be anyonic with fractional pair exchange phase $e^{i\pi/k}$. This phase has been experimentally observed [101].

Effective abelian Chern-Simons theory. The traditional proposal for an effective field theory description of k -fractional quantum Hall systems postulates that the effective field is a 1-form potential a for the electric current density 2-form J , itself minimally coupled to the *quasi-hole current* j , and with effective dynamics encoded by the level= k Chern-Simons (CS) Lagrangian [161][151]:

$$\begin{aligned}
 \text{Electron current} & \quad J = \overrightarrow{J} \lrcorner \text{dvol} \quad =: \text{d} a \quad \text{Effective gauge field} \\
 \text{density 2-form} & \\
 \text{Quasi-particle current} & \quad j = \overrightarrow{j} \lrcorner \text{dvol} \\
 \text{density 2-form} & \\
 \text{Background flux} & \quad F = \text{d} A \quad \text{External gauge field} \\
 \text{density 2-form} & \\
 \text{Effective Lagrangian} & \quad L := k \underbrace{\frac{1}{2} a \text{d} a}_{\text{CS}(a)} - \underbrace{A \text{d} a}_{A J} + a j \quad [151, (2.11)] \\
 \text{density 3-form} &
 \end{aligned}$$

Its Euler-Lagrange equations of motion

$$\frac{\delta L}{\delta a} = 0 \quad \Leftrightarrow \quad J = \frac{1}{k} (F - j)$$

in the case of longitudinal electron current and static quasi-particles

$$\begin{aligned}
 J & \equiv J_0 \text{d}x \text{d}y - J_x \text{d}t \text{d}y \\
 j & \equiv j_0 \text{d}x \text{d}y \\
 F & \equiv B \text{d}x \text{d}y - E_y \text{d}t \text{d}y
 \end{aligned}$$

express just the hallmark properties of the FQHE that we saw above, at filling fraction $\nu = 1/k$:

$$\Leftrightarrow \begin{cases} J_x = \frac{1}{k} E_y & \Leftrightarrow \text{Hall conductivity law at } 1/k \text{ filling} \\ J_0 = \frac{1}{k} B & \Leftrightarrow \text{each electron binds to } k \text{ flux quanta, but} \\ -\frac{1}{k} j_0 & \text{ } 1/k\text{th electron missing for each quasi-hole.} \end{cases}$$

Conceptual problems. However this can only be a *local* description on a single chart (as is common for Lagrangian field theories): Neither J nor F may admit global coboundaries a and A , respectively. Instead, both must be subjected to some kind of flux-quantization. For F this must be classical Dirac charge quantization, which however is incompatible with integrality of J when $k \neq 1$ (cf. [154, p. 35][144, p 159]). But without this, the implications break concerning topological order from abelian CS theory (ground state degeneracy, modular functoriality, ...).

Therefore we must ask:

Question: *Is there a non-Lagrangian theory for quasi-particles of properly flux-quantized FQH systems?*

Answer: Yes!:

The main result to be discussed here is that the key features of the anyonic topological order as seen in fractional quantum Hall systems are consistently, rigorously and naturally reflected by the topological light-cone quantization of the self-dual tensor field on M5-brane probes of certain orbi-singularities in 11D supergravity — once the subtle (non-abelian) flux-quantization of this field is properly taken care of, which is the key step that has not previously received attention. This is what we explain below.

Further aspects. In fact, fractional quantum Hall systems exhibit further remarkable properties which have not previously been reflected in their effective (Chern-Simons) descriptions, but which are naturally reflected in the M5-brane model, among them *hidden supersymmetry*. We close this introduction by briefly indicating this phenomenon.

N -Electron ground states of quantum Hall systems. While a microscopic derivation of fractional quantum Hall ground states Ψ remains missing, phenomenologically successful Ansätze exist:³

- At odd filling fraction $\nu = 1/q$, $q \in 2\mathbb{N} + 1$, the **Laughlin wavefunction**

$$\Psi_{\text{La}}(z^1, \dots, z^N) := \prod_{i < j} (z^i - z^j)^q \exp\left(-\frac{1}{\ell_B^2} \sum_i |z^i|^2\right)$$

³For N electrons in an effectively 2D material, and assumed to be completely spin-polarized by the transverse magnetic field, their wavefunction Ψ is a skew-symmetric (by Pauli exclusion) \mathbb{C} -valued function of N complex numbers $(z^i \in \mathbb{C})_{i=1}^N$. We omit normalization. For the Read-Moore state N must (for $\text{Pf}(-)$ to be defined) be even (which is harmless since N is a macroscopic number of electrons).

- At even filling fraction $\nu = 1/q$, $q \in 2\mathbb{N}$, the **Read-Moore wavefunction**

$$\Psi_{\text{RM}}(z^1, \dots, z^N) := \text{Pf} \left(\frac{1}{z^{\bullet_1} - z^{\bullet_2}} \right) \Psi_{\text{La}}(z^1, \dots, z^N).$$

Here the *Pfaffian* Pf of a skew-symmetric $N \times N$ matrix A is the Bererzinian integral over anti-commuting variables $(\theta^i)_{i=1}^N$:

$$\text{Pf}(A) := \int (\prod_i d\theta^i) \exp \left(\frac{1}{2} A_{ij} \theta^i \theta^j \right).$$


pick coefficient of
top θ -power

Hidden super-geometry of quantum Hall systems. This suggests to promote the plane \mathbb{C}^1 to the super-space $\mathbb{C}^{1|1}$ with its *super-translation group* structure

$$(z, \theta) + (z', \theta') = (z + z' + \theta\theta', \theta + \theta')$$

Here the **super-Laughlin state** exhibits the Read-Moore state as a super-partner to the Laughlin state (up to normalization) [64][57, (13)]:

$$\Psi_{\text{sLa}}((z^1, \theta^1), \dots, (z^N, \theta^N)) := \prod_{i < j} (z^i - z^j - \theta^i \theta^j)^q \exp \left(-\frac{1}{\ell_B^2} \sum_i |z^i|^2 \right)$$



Collective excitations. The Moore-Read state is known to have two *density-wave excitations* for wave-vectors $k \in \mathbb{C}$:

- (i) The **magneto-roton state**

$$\Psi_{\text{MR},k}(z^1, \dots, z^N) := \sum_i \exp(-i\bar{k}\partial_{z^i}) \exp(-\frac{i}{2}\bar{k}z^i) \Psi_{\text{MR}}(z^1, \dots, z^N)$$


- (ii) The **neutral fermion state**

$$\Psi_{\text{NF},k} \text{ which originally did not have a closed expression}$$

However, lifting the magneto-roton state to super-space, for super-wavevector $(k, \kappa) \in \mathbb{C}^{1,1}$

$$\Psi_{\text{MR},(k,\kappa)}(z^1, \dots, z^N) := \int (\prod_i d\theta^i) \sum_i \exp(-i\bar{k}\partial_{z^i}) \exp(-\frac{i}{2}\bar{k}z^i) \exp(-\frac{i}{2}\bar{\kappa}\theta^i) \Psi_{\text{sLa}}((z^1, \theta^1), \dots, (z^N, \theta^N))$$

it reproduces the magneto-roton state for even N , and the neutral fermion mode when an $(N + 1)$ st electron is added [57]:



Hidden super-symmetry in fractional quantum Hall systems. This super-unification predicts hidden supersymmetry in fractional quantum Hall systems — which is indeed (numerically) observed [117][89] (also [5, §5]).

This all suggests that an accurate model for fractional quantum Hall systems should in fact itself *originate on superspace*, and this is what we start with now.

2 Flux-Quantization on M5-Probes

The first task now is to understand the flux-quantization on M5-brane probes, according to [39][41][134]. We will not (need to) explain in full detail the (super-)geometry of probe branes nor of their (super-)gravity backgrounds (full discussion is in [52][53]), but do offer the following broad dictionary, for orientation: ⁴

M5-Brane probes (namely *sigma-model* branes, in contrast to *black branes*) are 5-dimensional objects propagating in a gravitational target space X (the “bulk”), along trajectories that are modeled by (super-)immersions of their 6D (and $\mathcal{N} = (2, 0)$) worldvolume (super-)manifolds Σ

$$\begin{array}{ccc} \text{probe M5-brane} & \xrightarrow{\phi_s} & X^{1,10} | \mathbf{32} \\ \text{(super-)worldvolume} & \text{trajectory} & \text{target/background} \\ \Sigma^{1,5} | 2 \cdot \mathbf{8}_+ & \text{(super-)immersion} & \text{(super-)spacetime} \end{array} \quad (1)$$

Here the admissible (“on-shell”, meaning: satisfying the appropriate equations of motion) immersions ϕ_s are controlled by the (super-)geometry of X – namely the brane’s trajectory is subject to the gravitational- and Lorentz-forces exerted by the field content of X – but X itself remains unaffected by the choice of ϕ_s – meaning that the (gravitational) *back-reaction* of the brane on its ambient spacetime is neglected; this is what makes the brane but a *probe* of the *background* X .

Thereby the probe brane (Σ, ϕ_s) plays a double role:

- (i) on the one hand it is like a (higher-dimensional) fundamental particle, an “observer” of the bulk X in the sense of mathematical relativity,
- (ii) on the other hand it is itself a (super-)spacetime with its own (quantum) field content:

Remarkably, the magic of super-geometry makes such purely super-geometric immersions ϕ_s (1) embody not just the naïve (temporal-)spatial worldvolume trajectory, but also a 3-flux density H_3^s on Σ [53, §3.3]. This is (on-shell) the notorious “self-dual” flux density whose accurate quantization (traditionally neglected) is our main concern here.

This second aspect is what we are concerned with for the purpose of modeling strongly-coupled quantum systems: The (1+3)D worldvolume $M^{1,3}$ of a quantum material – or, for the intent of modeling anyons, the effectively (1 + 2) D -worldvolume $M^{1,2}$ of a sheet-like material (e.g. an atomic mono-layer akin to graphene) – is to be identified with a sub-quotient of the brane worldvolume, typically with a fixed locus (orbifold singularity) inside the base of a fibration (Kaluza-Klein reduction).

$$\begin{array}{ccc} \text{M5-brane} & & \text{ambient bulk} \\ \text{worldvolume} & & \text{spacetime} \\ \Sigma^{1,5} & \xrightarrow{\phi} & Y^{1,10} \\ \downarrow & & \downarrow \text{M/IIA-} \\ \text{orbi-} & & \text{fibration} \\ \text{singularity} & & \\ M^{1,2} & \xleftarrow{\text{orbi-}} & \Sigma^{1,4} \xrightarrow{\text{orbi-}} X^{1,9} \\ \text{quantum material} & & \text{D4-brane} \\ \text{worldvolume} & & \text{worldvolume} \end{array}$$


Their flux quantization (to recall from [130]) is then encoded in a choice of a fibration $\mathcal{A} \xrightarrow{p} \mathcal{B}$ of classifying spaces, subject to the constraint that the Bianchi identities for the (duality-symmetric) flux densities on bulk and brane are the closure/flatness condition on l -valued differential forms, where $l(-)$ forms *Whitehead L_∞ -algebras* of these classifying fibrations (dual to their minimal relative Sullivan model).

Given such a choice, the topological sector of the higher gauge fields on bulk and brane are given by maps from the brane-immersion into the classifying fibration:

With these comments on perspective out of the way, **the plan of this section** are the following topics:

- (i) [Bianchi identities on magnetized M5-probes](#)
- (ii) [Flux quantization in Twistorial Cohomotopy](#)
- (iii) [Aside: Projective Spaces and their Fibrations](#)
- (iv) [Orbi-worldvolumes and Equivariant charges](#)

⁴All brane concepts we consider are well-defined and all conclusions have proofs – at no point do we rely on informal string theory folklore beyond motivation.



The first step of flux quantization is to identify the Bianchi identities satisfied by the flux densities:

Bianchi identities on M5-Probes of 11D SuGra via super-geometry. Consider the 11D super-tangent space

$$\mathbb{R}^{1,10|\mathbf{32}} \begin{array}{c} \text{super-Minkowski} \\ \longleftarrow \\ \text{isom}(\mathbb{R}^{1,10|\mathbf{32}}) \\ \text{super-Poincaré} \\ \longrightarrow \\ \mathfrak{so}(1,10) \\ \text{Lorentz} \end{array}$$

with its super-invariant 1-forms (cf. [52, §2.1]):

$$\text{CE}(\mathbb{R}^{1,10|\mathbf{32}}) \simeq \underbrace{\Omega_{\text{dR}}^\bullet(\mathbb{R}^{1,10|\mathbf{32}})^{\text{li}}}_{\text{super-transl. invar. forms}} \simeq \mathbb{R}_{\text{d}} \left[\begin{array}{l} (\Psi^\alpha)_{\alpha=1}^{32} \\ (E^a)_{a=0}^{10} \end{array} \right] / \left(\begin{array}{l} \text{d}\Psi^\alpha = 0 \\ \text{d}E^a = (\bar{\Psi}\Gamma^a\Psi) \end{array} \right).$$

Remarkably, the quartic Fierz identities entail that [25][100][52, Prop. 2.73]:

$$\left. \begin{array}{l} G_4^0 := \frac{1}{2}(\bar{\Psi}\Gamma_{a_1 a_2}\Psi)E^{a_1}E^{a_2} \\ G_7^0 := \frac{1}{5!}(\bar{\Psi}\Gamma_{a_1 \dots a_5}\Psi)E^{a_1} \dots E^{a_5} \end{array} \right\} \in \text{CE}(\mathbb{R}^{1,10|\mathbf{32}})^{\text{Spin}(1,10)} \quad \text{satisfy:} \quad \begin{array}{l} \text{d}G_4^0 = 0 \\ \text{d}G_7^0 = \frac{1}{2}G_4^0 G_4^0 \end{array}$$

fully super-invariant forms

To globalize this situation, say that an **11D super-spacetime** X is a super-manifold equipped with a super-Cartan connection, locally on an open cover $\tilde{X} \rightarrow X$ given by

$$\left. \begin{array}{l} (\Psi^\alpha)_{\alpha=1}^{32} \\ (E^a)_{a=0}^{10} \\ (\Omega^{ab} = -\Omega^{ba})_{a,b=0}^{10} \end{array} \right\} \in \Omega_{\text{dR}}^1(\tilde{X}) \quad \text{such that the super-torsion vanishes} \quad \text{d}E^a - \Omega^a_b E^b = (\bar{\Psi}\Gamma^a\Psi),$$

and say that **C-field super-flux** on such a super-spacetime are super-forms with these co-frame components:

$$\boxed{\begin{array}{l} G_4^s := G_4 + G_4^0 := \frac{1}{4!}(G_4)_{a_1 \dots a_4} E^{a_1} \dots E^{a_4} + \frac{1}{2}(\bar{\Psi}\Gamma_{a_1 a_2}\Psi)E^{a_1}E^{a_2} \\ G_7^s := G_7 + G_7^0 := \frac{1}{7!}(G_7)_{a_1 \dots a_7} E^{a_1} \dots E^{a_7} + \frac{1}{5!}(\bar{\Psi}\Gamma_{a_1 \dots a_5}\Psi)E^{a_1} \dots E^{a_5} \end{array}}$$

Theorem [52, Thm. 3.1]: On an 11D super-spacetime X with C-field super-flux (G_4^s, G_7^s) :

$$\text{The duality-symmetric super-Bianchi identity} \quad \left\{ \begin{array}{l} \text{d}G_4^s = 0 \\ \text{d}G_7^s = \frac{1}{2}G_4^s G_4^s \end{array} \right\} \text{ is equivalent to} \quad \text{the full 11D SuGra equations of motion!}$$

Next, on the super-subspace $\mathbb{R}^{1,5|2\cdot\mathbf{8}_+} \xrightarrow{\phi_0} \mathbb{R}^{1,10|\mathbf{32}}$ fixed by the involution $\Gamma_{012345} \in \text{Pin}^+(1,10)$ we have:

$$H_3^0 := 0 \in \text{CE}(\mathbb{R}^{1,5|2\cdot\mathbf{8}_+})^{\text{Spin}(1,5)} \quad \text{satisfies:} \quad \text{d}H_3^0 = \phi_0^* G_4^0$$

To globalize this situation, say that a super-immersion $\Sigma^{1,5|2\cdot\mathbf{8}_+} \xrightarrow{\phi_s} X^{1,10|\mathbf{32}}$ is $1/2$ **BPS M5** if it is ‘‘locally like’’ ϕ_0 , and say that **B-field super-flux** on such an M5-probe is a super-form with these co-frame components:

$$\boxed{H_3^s := H_3 + H_3^0 := \frac{1}{3!}(H_3)_{a_1 a_2 a_3} e^{a_1} e^{a_2} e^{a_3} + 0} \quad (e^{a<6} := \phi_s^* E^a)$$

Theorem [53, §3.3]: On a super-immersion ϕ_s with B-field super-flux H_3^s :

The super-Bianchi identity $\{dH_3^s = \phi_s^* G_4^s\}$ is equivalent to the M5's B-field equations of motion.

In particular, the (non-linear self-)duality conditions on the ordinary fluxes are implied: $G_4 \leftrightarrow G_7$ and $H_3 \leftrightarrow H_3$.

Seeing from this that also trivial tangent super-cochains may have non-trivial globalization, observe next that:

$$F_2^0 := (\bar{\psi}\psi) = 0 \in \text{CE}(\mathbb{R}^{1,5|2\cdot\mathbf{8}_+})^{\text{Spin}(1,5)} \quad \text{satisfies :} \quad dF_2^0 = 0$$

Globalizing this to $\Sigma^{1,5|2\cdot\mathbf{8}_+}$ via

$$F_2^s := F_2 + F_2^s := \frac{1}{2}(F_2)_{a_1 a_2} e^{a_1} e^{a_2} + 0$$

we have on top of the above:

Theorem [135, p 7]:

The super-Bianchi identity $\{dF_2^s = 0\}$ is equivalent to the Chern-Simons E.O.M.: $F_2 = 0$.

Flux quantization in Twistorial Cohomotopy. In summary, a remarkable kind of higher super-Cartan geometry locally modeled on the 11D super-Minkowski spacetime $\mathbb{R}^{1,10|32}$ entails that on-shell 11D supergravity probed by magnetized $1/2$ BPS M5-branes implies and is entirely governed by these Bianchi identities on super-flux densities:

A-field	$dF_2^s = 0$	$dG_4^s = 0$	C-field
self-dual B-field	$dH_3^s = \phi_s^* G_4^s + \theta F_2^s F_2^s$	$dG_7^s = \frac{1}{2} G_4^s G_4^s$	dual C-field
M5 probe	$\Sigma^{1,5 2\cdot\mathbf{8}_+}$	$\xrightarrow[\text{1/2BPS immersion}]{\phi_s}$	$X^{1,10 32}$
			SuGra bulk

Here we have observed that the Green-Schwarz term $F_2^s F_2^s$ may equivalently be included for any theta-angle $\theta \in \mathbb{R}$ without affecting the equations of motion (since, recall, the CS e.o.m. $F_2^s = 0$ is already implied by $dF_2^s = 0$).

However, non-vanishing theta-angle does affect the admissible flux-quantization laws and hence the global solitonic and torsion charges of the fields. The choice of flux quantization according to *Hypothesis H* [39][41] is the following:

Admissible fibrations of classifying spaces for cohomology theories with the above character images (2).

The homotopy quotient of S^7 is

- (i) for $\theta = 0$ by the trivial action and
- (ii) for $\theta \neq 0$ by the principal action of the complex Hopf fibration.

$$\begin{array}{ccccc}
 \boxed{\theta = 0} & S^7 //_{\theta} U(1) \simeq S^7 \times \mathbb{C}P^\infty & \longrightarrow & S^7 & \xrightarrow[\text{\mathbb{H-Hopf fibration}}]{h_{\mathbb{H}}} & \mathbb{H}P^1 \\
 & \downarrow & & \downarrow & & \parallel \\
 & & & \text{C-Hopf fibration} & & \\
 \boxed{\theta \neq 0} & S^7 // U(1) & \xrightarrow{\sim} & \mathbb{C}P^3 & \xrightarrow[\text{\textit{Twistor fibration}}]{t_{\mathbb{H}}} & \mathbb{H}P^1
 \end{array}$$

Proof. This may be seen as follows [41, Lem. 2.13]:

Since the real cohomology of projective space is a truncated polynomial algebra,

$$\begin{array}{ll}
 H^\bullet(\mathbb{C}P^n; \mathbb{R}) \simeq \mathbb{R}[\underbrace{c_1}_{\text{deg}=2}] / (c_1^{n+1}) & H^\bullet(\underbrace{\mathbb{C}P^\infty}_{\simeq BU(1)}; \mathbb{R}) \simeq \mathbb{R}[c_1] \\
 H^\bullet(\mathbb{H}P^n; \mathbb{R}) \simeq \mathbb{R}[\underbrace{\frac{1}{2}p_1}_{\text{deg}=4}] / (p_1^{n+1}) & H^\bullet(\underbrace{\mathbb{H}P^\infty}_{\simeq BSp(1) \simeq BSU(2) \simeq BSpin(3)}; \mathbb{R}) \simeq \mathbb{R}[\frac{1}{2}p_1],
 \end{array}$$

the minimal dgc-algebra model for $\mathbb{C}P^n$ needs a closed generator f_2 to span the cohomology and a generator h_{2n+1} in order to truncate it; analogously for $\mathbb{H}P^n$. Since these generators also form a graded linear basis for the rationalized homotopy groups of these spaces, they give the minimal Sullivan models (cf [130, Prop. 3.7]):

$$\begin{aligned} \text{CE}(\mathbb{I}CP^n) &\simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ h_{2n+1} \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d h_{2n+1} = (f_2)^{n+1} \end{array} \right) \\ \text{CE}(\mathbb{I}HP^n) &\simeq \mathbb{R}_d \left[\begin{array}{c} g_4 \\ g_{4n+3} \end{array} \right] / \left(\begin{array}{l} d g_4 = 0 \\ d g_{4n+3} = (g_4)^{n+1} \end{array} \right). \end{aligned}$$

Furthermore, since the second Chern class of a $U(1) \simeq S(U(1)^2) \subset SU(2)$ -bundle is minus the cup square of the first Chern class (by the Whitney sum rule), so that (cf. [124, (216)])

$$\begin{array}{ccc} \mathbb{C}P^3 & \hookrightarrow & \mathbb{C}P^\infty \simeq BU(1) & & -(c_1)^2 \\ \downarrow t_{\mathbb{H}} & & \downarrow & & \downarrow B(c \rightarrow \text{diag}(c, c^*)) \\ \mathbb{H}P^1 & \hookrightarrow & \mathbb{H}P^\infty \simeq BSU(2) & & \frac{1}{2}p_1 = c_2, \end{array}$$

the minimal model of $\mathbb{C}P^3$ relative to that of $\mathbb{H}P^1 \simeq S^4$ (cf. [42, Prop. 4.24]) needs to adjoin to the latter not only f_2 but also a generator h_3 imposing this relation in cohomology, whence it must be

$$\text{CE}(\mathbb{I}_{\mathbb{H}P^1} \mathbb{C}P^3) \simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ h_3 \\ g_4 \\ g_7 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d h_3 = g_4 + f_2 f_2 \\ d g_4 = 0 \\ d g_7 = \frac{1}{2} g_4 g_4 \end{array} \right),$$

which is clearly quasi-isomorphic to $\text{CE}(\mathbb{I}CP^3)$. □

The resulting fibration of L_∞ -algebras is manifestly just that classifying the desired Bianchi identities (2) (we are showing the case $\theta \neq 0$, which by isomorphic rescaling may be taken to be $\theta = 1$):

$$\begin{array}{ccccc} \Sigma^6 & \dashrightarrow & \Omega_{\text{dR}}^1(-; \mathbb{I}_{\mathbb{H}P^1} \mathbb{C}P^3)_{\text{clsd}} & \xleftarrow{\quad} & \text{CE}(\mathbb{I}_{\mathbb{H}P^1} \mathbb{C}P^3) & \xrightarrow{F_2} & \Omega_{\text{dR}}^\bullet(\Sigma^6) & \left| \begin{array}{l} d F_2 = 0 \\ d H_3 = G_4 + F_2 F_2 \end{array} \right. \\ \downarrow \phi & & \downarrow (t_{\mathbb{H}})_* & \Leftrightarrow & \downarrow (t_{\mathbb{H}})^* & \Leftrightarrow & \downarrow \phi^* & \\ \Sigma^{11} & \dashrightarrow & \Omega_{\text{dR}}^1(-; \mathbb{I}HP^1)_{\text{clsd}} & \xleftarrow{\quad} & \text{CE}(\mathbb{I}HP^1) & \xrightarrow{G_4} & \Omega_{\text{dR}}^\bullet(X^{11}) & \left| \begin{array}{l} d G_4 = 0 \\ d G_7 = \frac{1}{2} G_4 G_4 \end{array} \right. \end{array}$$

Aside: Projective Spaces and their Fibrations – Here we used the following classical facts. Consider:

division algebras $\mathbb{R} \hookrightarrow \mathbb{C} \hookrightarrow \mathbb{H}$ generically denoted $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$

groups of units $\mathbb{K}^\times := \mathbb{K} \setminus \{0\}$ understood with the multiplicative group structure

projective spaces $\mathbb{K}P^n := (\mathbb{K}^{n+1} \setminus \{0\})/\mathbb{K}^\times$

higher spheres $S^n \simeq (\mathbb{R}^{n+1} \setminus \{0\})/\mathbb{R}_{>0}$

\mathbb{K} -Hopf fibrations are the quotient co-projections induced by $\iota : \mathbb{R}_{>0} \hookrightarrow \mathbb{K}$

The classical Hopf fibrations $h_{\mathbb{K}}$ are:

$$\begin{array}{ccc} \begin{array}{c} S^0 \simeq \mathbb{R}^\times / \mathbb{R}_{>0} \\ \downarrow \ker \\ S^1 \simeq (\mathbb{R}^2 \setminus \{0\}) / \mathbb{R}_{>0} \\ \downarrow h_{\mathbb{R}} \\ S^1 \simeq \underbrace{(\mathbb{R}^2 \setminus \{0\}) / \mathbb{R}^\times}_{\mathbb{R}P^1} \end{array} & \begin{array}{c} S^1 \simeq \mathbb{C}^\times / \mathbb{R}_{>0} \\ \downarrow \ker \\ S^3 \simeq (\mathbb{C}^2 \setminus \{0\}) / \mathbb{R}_{>0} \\ \downarrow h_{\mathbb{C}} \\ S^2 \simeq \underbrace{(\mathbb{C}^2 \setminus \{0\}) / \mathbb{C}^\times}_{\mathbb{C}P^1} \end{array} & \begin{array}{c} S^3 \simeq \mathbb{H}^\times / \mathbb{R}_{>0} \\ \downarrow \ker \\ S^7 \simeq (\mathbb{H}^2 \setminus \{0\}) / \mathbb{R}_{>0} \\ \downarrow h_{\mathbb{H}} \\ S^4 \simeq \underbrace{(\mathbb{H}^2 \setminus \{0\}) / \mathbb{H}^\times}_{\mathbb{H}P^1} \end{array} \end{array}$$

The Hopf fibrations in higher dimensions are the attaching maps exhibiting the topological cell-complex structure of projective spaces [104], from which the (cellular) cohomology follows readily.

Further factor-fibrations arise by factoring the Hopf fibrations via the stage-wise quotienting along

$$\mathbb{R}_{>0} \hookrightarrow \mathbb{R} \hookrightarrow \mathbb{C} \hookrightarrow \mathbb{H}.$$

Notably, the classical quaternionic Hopf fibration $h_{\mathbb{H}}$ factors through a higher-dimensional complex Hopf fibration followed by the **Calabi-Penrose twistor fibration** $t_{\mathbb{H}}$ [41, §2].

Equivariantization: Since the quotienting is by right actions, these fibrations are equivariant under the left action of

$$\text{Spin}(5) \simeq \text{Sp}(2) := \{g \in \text{GL}_2(\mathbb{H}) \mid g^\dagger \cdot g = e\}.$$

$$\begin{array}{ccc} S(\mathbb{K}^{n+1}) & \longrightarrow & * \\ h_{\mathbb{K}} \downarrow & \swarrow (po) & \downarrow \\ \mathbb{K}P^n & \hookrightarrow & \mathbb{K}P^{n+1} \end{array}$$

$$\begin{array}{ccc} S^1 \simeq \mathbb{C}^\times / \mathbb{R}_{>0} & & \\ & \searrow & \\ & & S^7 \simeq (\mathbb{H}^2 \setminus \{0\}) / \mathbb{R}_{>0} \\ & \swarrow & \downarrow h_c \text{ complex Hopf fibration} \\ S^2 \simeq \mathbb{H}^\times / \mathbb{C}^\times & & \mathbb{C}P^3 \simeq (\mathbb{H}^2 \setminus \{0\}) / \mathbb{C}^\times \\ & \searrow h_{\mathbb{H}} \text{ quaternionic Hopf fibration} & \downarrow t_{\mathbb{H}} \text{ Calabi-Penrose twistor fibration} \\ & & \mathbb{H}P^1 \simeq (\mathbb{H}^2 \setminus \{0\}) / \mathbb{H}^\times \end{array}$$

For example, the involution $\sigma := \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \in \text{Sp}(2)$ swaps the two copies of \mathbb{H} :

$$\begin{array}{ccc} \mathbb{C}P^3 & \xrightarrow{t_{\mathbb{H}}} & \mathbb{H}P^1 \\ \downarrow \sigma & \Downarrow & \downarrow \sigma \\ (\mathbb{H} \times \mathbb{H} \setminus \{0\}) / \mathbb{C}^\times & \rightarrow & (\mathbb{H} \times \mathbb{H} \setminus \{0\}) / \mathbb{H}^\times \\ \downarrow \sigma & \Downarrow & \downarrow \sigma \\ (\mathbb{H} \oplus \mathbb{H} \setminus \{0\}) / \mathbb{C}^\times & \rightarrow & (\mathbb{H} \oplus \mathbb{H} \setminus \{0\}) / \mathbb{H}^\times \\ \downarrow \sigma & \Downarrow & \downarrow \sigma \\ \mathbb{C}P^3 & \xrightarrow{t_{\mathbb{H}}} & \mathbb{H}P^1 \end{array}$$

The resulting \mathbb{Z}_2 -fixed locus is the 2-sphere:

$$\begin{array}{ccc} (\mathbb{C}P^3)^{\mathbb{Z}_2} \simeq (\mathbb{H} \setminus \{0\}) / \mathbb{C}^\times \simeq S^2 & & \\ \downarrow (t_{\mathbb{H}})^{\mathbb{Z}_2} & \downarrow & \downarrow \\ (\mathbb{H}P^1)^{\mathbb{Z}_2} \simeq (\mathbb{H} \setminus \{0\}) / \mathbb{H}^\times \simeq * & & \end{array}$$

This is the 2-sphere coefficient that will end up being responsible for stabilizing anyons on orbi-worldvolumes! We next discuss how this comes about.

Aside: Implications of Hypothesis H, in view of traditional expectations for M-theory.

The **plain Hypothesis H** in the bulk says that the non-perturbative completion of the C-field in 11d supergravity is a cocycle in *differential Cohomotopy* $\widehat{\pi}^4$ [36, §4][55, §3.1][42, Ex. 9.3] and as such involves (exposition in [130, §3.3]) a map χ from spacetime to the homotopy type of the 4-sphere, with the C-field gauge potentials $(\widehat{C}_3, \widehat{C}_6)$ exhibiting the flux densities (G_4, G_7) as \mathbb{R} -rational representatives of χ .

$$\begin{array}{ccc} \begin{array}{c} \text{canonical differential} \\ \text{non-abelian (unstable)} \\ 4\text{-Cohomotopy} \\ (\widehat{C}_3, \widehat{C}_6) \\ \text{full nonperturbative} \\ \text{11d SuGra C-field} \end{array} \in \widehat{\pi}^4(X) & \xrightarrow[\text{topological sector}]{\chi} & \begin{array}{c} \text{plain} \\ \text{non-abelian (unstable)} \\ 4\text{-Cohomotopy} \\ \pi^4(X) \end{array} \\ \downarrow \text{flux densities} & & \downarrow \text{Cohomotopical charge sector} \\ H_{\text{dR}}(X; \mathbb{I}S^4) & & \text{Maps}(X; S^4) \\ \downarrow \text{\mathbb{I}S^4\text{-valued de Rham cohomology}} & & \downarrow \text{ch} \\ \Omega_{\text{dR}}^1(X; \mathbb{I}S^4)_{\text{clsd}} & \xrightarrow{\eta^f} & \int \Omega_{\text{dR}}^1(X; \mathbb{I}S^4)_{\text{clsd}} \\ \downarrow \text{C-field flux densities} & \mapsto & \downarrow \text{character image} \\ (G_4, G_7) & \mapsto & \eta^f(G_4, G_7) \xrightarrow[\text{gauge potentials}]{(\widehat{C}_3, \widehat{C}_6)} \end{array}$$

As an immediate plausibility check, from the well-known homotopy groups of spheres in low degrees this implies (cf. [68, (22-3)][124, (22)]):


- Integral quantization of charges carried by singular M5-brane branes

$$\begin{aligned}\pi^4(\mathbb{R}^{10,1} \setminus \mathbb{R}^{5,1}) &= \pi^4(\mathbb{R}^{5,1} \times \mathbb{R}_+ \times S^4) \\ &= \pi^4(S^4) = \pi_4(S^4) = \mathbb{Z}.\end{aligned}$$

- Integral quantization of charges carried by singular M2-branes... plus a torsion-contribution (a first prediction of Hypothesis H: fractional M2-branes):

$$\begin{aligned}\pi^4(\mathbb{R}^{10,1} \setminus \mathbb{R}^{2,1}) &= \pi^4(\mathbb{R}^{2,1} \times \mathbb{R}_+ \times S^7) \\ &= \pi^4(S^7) = \pi_7(S^4) = \mathbb{Z} \oplus \mathbb{Z}_{12}.\end{aligned}$$

Hypothesis H with curvature corrections. More generally, curvature corrections from the coupling to the background gravity are postulated to be reflected in *tangentially twisted* 4-Cohomotopy [39], analogous to the well-known twisting of the RR-field flux-quantization in K-theory by its background B-field:



To distinguish M2/M5-charge, the tangential twisting needs to preserve the \mathbb{H} -Hopf fibration \Rightarrow tangential $\text{Sp}(2) \hookrightarrow \text{Spin}(8)$ -structure [39, §2.3]. With this, integrality of M2's Page charge & anomaly-cancellation of the M5's Hopf-WZ term follows from trivialization of the Euler 8-class, which means lift to the *Fivebrane* 6-group $\widehat{\text{Sp}}(2) \rightarrow \text{Sp}(2)$ [38, §4].

This implies [39, Prop. 3.13][38, Thm. 4.8]:

- (i) half-integrally shifted quantization of M5-brane charge in curved backgrounds,

$$[\tilde{G}_4] := \underbrace{[G_4]}_{\text{C-field 4-flux}} + \frac{1}{2} \underbrace{\left(\frac{1}{2} p_1(TX^8)\right)}_{\text{integral Spin-Pontrjagin class}} \in H^4(X^8; \mathbb{Z})$$

- (ii) integral quantization of the Page charge of M2-branes.

$$2[\tilde{G}_7] := 2([G_7] + \frac{1}{2}[H_3 \wedge \tilde{G}_4]) \in H^7(\hat{X}^8; \mathbb{Z})$$

Both of these quantization conditions on M-brane charge Previously, item (i) had remained enigmatic and item are thought to be crucial for M-theory to make any sense. (ii) had remained wide open.

But there is more:

Provable implications from Hypothesis H of subtle effects expected in M-theory:

- half-integral shift of 4-flux [39, Prop. 3.13]
- DMW anomaly cancellation [39, Prop. 3.7]
- the C-field's "integral EoM" [39, §3.6]
- M2 Page charge quantization [38, Thm. 4.8]
- integrality of $\frac{1}{6}(G_4)^3$ [55, Rem. 2.9]
- M5-brane anomaly cancellation [123]
- non-abelian gerbe field on M5 [40]

It is these and further results that suggest that Hypothesis H goes towards the correct flux-quantization law for the C-field in M-theory.

Yet more generally, Hypothesis H applies to orbifold spacetimes, where it postulates flux quantization in (twisted and) *equivariant* Cohomotopy [121][14]. This is what we turn to next.

Orbi-worldvolumes and Equivariant charges. Flux-quantization generalizes to *orbifolds*⁵ by generalizing the cohomology of the charges to *equivariant cohomology* [122]. In terms of classifying spaces this simply means that all spaces are now equipped with the action of a finite group G and all maps are required to be G -equivariant.

We take $G := \mathbb{Z}_2$ and the classifying fibration to be the **twistor fibration** $p := t_{\mathbb{H}}$ equivariant under swapping the \mathbb{H} -summands,

$$\begin{array}{ccccc}
 \text{orbi-worldvolume} & \begin{array}{c} \langle G \rangle \\ \Sigma \end{array} & \xrightarrow{\text{orbi-brane charges}} & \begin{array}{c} \langle G \rangle \\ \mathcal{A} \end{array} & \xlongequal{\quad} & \begin{array}{c} \langle \mathbb{Z}_2 \rangle \\ \mathbb{C}P^3 \end{array} \\
 \downarrow \phi & \swarrow \text{orbi-brane charges} & & \downarrow p \text{ equivariant classifying fibration...} & & \downarrow t_{\mathbb{H}} \text{ ...for equivariant twistorial Cohomotopy} \\
 \text{orbi-spacetime} & \begin{array}{c} X \\ \langle G \rangle \end{array} & \xrightarrow{\text{orbi-bulk charges}} & \begin{array}{c} \mathcal{B} \\ \langle G \rangle \end{array} & \xlongequal{\quad} & \begin{array}{c} S^4 \\ \langle \mathbb{Z}_2 \rangle \end{array}
 \end{array}$$

and the brane/bulk orbifold we take to be as on p. 3:

The orbi-brane diagram for a flat M5-brane wrapped on a trivial Seifert-fibered orbi-singularity. Shaded is the \mathbb{Z}_2 -fixed locus/orbi-singularity.

We are adjoining the *point at infinity* to the space $\mathbb{R}_{\cup\{\infty\}}^2 \xrightarrow{\text{homeo}} S^2$ which is thereby designated as transverse to any worldvolume solitons to be measured in reduced cohomology.

$$\begin{array}{c}
 \begin{array}{c} \langle \mathbb{Z}_2 \rangle \\ \Sigma \end{array} := \mathbb{R}^{1,0} \times \mathbb{R}_{\cup\{\infty\}}^2 \times S^1 \times \mathbb{R}_{\text{sgn}}^2 \langle \mathbb{Z}_2 \rangle \\
 \downarrow \phi \\
 \begin{array}{c} X \\ \langle \mathbb{Z}_2 \rangle \end{array} := \mathbb{R}^{1,0} \times \mathbb{R}_{\cup\{\infty\}}^2 \times S^1 \times \mathbb{R}_{\text{sgn}}^2 \langle \mathbb{Z}_2 \rangle \times \mathbb{R}^5
 \end{array}$$

time
trnsvrs space to solitons
M/IIA-circle
orbi-cone
trnsvrs space to M5-brane

But since the cone $\mathbb{Z}_2 \curvearrowright \mathbb{R}_{\text{sgn}}^2$ is equivariantly contractible, the inclusion of the \mathbb{Z}_2 -fixed loci is actually a homotopy equivalence

$$\begin{array}{ccc}
 \begin{array}{c} \mathbb{Z}_2 \curvearrowright * \\ \text{cone} \end{array} & \xrightarrow{\text{hmtp}} & \mathbb{Z}_2 \curvearrowright * \\
 \Rightarrow & & \begin{array}{ccc}
 \Sigma^{\mathbb{Z}_2} & \xrightarrow{\sim \text{hmtp}} & \begin{array}{c} \langle \mathbb{Z}_2 \rangle \\ \Sigma \end{array} \\
 \phi^{\mathbb{Z}_2} \downarrow & & \downarrow \phi \\
 X^{\mathbb{Z}_2} & \xrightarrow{\sim \text{hmtp}} & \begin{array}{c} X \\ \langle G \rangle \end{array}
 \end{array}
 \end{array}$$

Therefore, our equivariant classifying maps are determined up to equivariant homotopy by their restriction to the fixed-locus and hence the charges are *localized on the orbi-singularity* where they take values in 2-Cohomotopy:

$$\left\{ \begin{array}{ccc}
 \begin{array}{c} \langle \mathbb{Z}_2 \rangle \\ \Sigma \end{array} \xrightarrow{\quad} \begin{array}{c} \langle \mathbb{Z}_2 \rangle \\ \mathbb{C}P^3 \end{array} \\
 \downarrow \phi \swarrow \text{orbi-brane charges} \quad \downarrow t_{\mathbb{H}} \\
 \begin{array}{c} X \\ \langle \mathbb{Z}_2 \rangle \end{array} \xrightarrow{\quad} \begin{array}{c} S^4 \\ \langle \mathbb{Z}_2 \rangle \end{array} \\
 \text{charges on orbifold}
 \end{array} \right\} \simeq \left\{ \begin{array}{ccc}
 \Sigma^{\mathbb{Z}_2} \xrightarrow{\quad} (\mathbb{C}P^3)^{\mathbb{Z}_2} \xlongequal{\quad} S^2 \\
 \phi^{\mathbb{Z}_2} \downarrow \swarrow \text{orbi-brane charges} \quad \downarrow t_{\mathbb{H}}^{\mathbb{Z}_2} \quad \downarrow \\
 X^{\mathbb{Z}_2} \xrightarrow{\quad} (S^4)^{\mathbb{Z}_2} \xlongequal{\quad} * \\
 \text{charges localized on orbi-singularity}
 \end{array} \right\} \simeq \left\{ \begin{array}{ccc}
 \mathbb{R}_{\cup\{\infty\}}^2 \times S^1 \xrightarrow{\quad} S^2 \\
 \text{charges in 2-Cohomotopy of B-field solitons on M5 orbi-singularity}
 \end{array} \right\}$$


Moduli space of worldvolume solitons. To be precise, the solitonic charges are to be measured in the *reduced* 2-Cohomotopy classified by *pointed maps*, enforcing the condition that solitonic fields *vanish at infinity* [130, §2.2]. In the strongly-coupled situation, where the M/IIA circle de-compactifies to \mathbb{R}^1 , the vanishing-at-infinity must also be applied here, whence (cf. [131, §A.2]) the moduli space of topological solitons is the loop space of the reduced 2-Cohomotopy moduli of the transverse space:

$$\begin{array}{ccc}
 \text{Moduli space of solitons on M5 orbi-singularity} & \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2 \wedge S^1, S^2) & \simeq & \Omega \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2) & \text{Loop space of moduli space of solitons on D4 orbi-singularity} \\
 & \simeq & & \simeq & \\
 & \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, \Omega S^2) & & &
 \end{array}$$


(The algebraic topology of maps to ΩS^2 have also found some attention by [97][98].)

⁵For brevity we consider here only “very good” orbifolds, namely global quotients of manifolds by the action of a finite group G . This is sufficient for the present purpose and anyways the case understood by default in the string theory literature.

Outlook. Strikingly, as we explain next, this moduli space is equivalently a space of *worldsheets of strings* in \mathbb{R}^3 with unit *charged endpoints* forming oriented *framed links*! [132]




Such link diagrams are just the envisioned topological quantum circuit protocols, and their framing regularizes the anyonic phase observables (“Wilson loop observables”).




3 Cohomotopy Charge of Solitons

Remarkably, there is an equivalence between *Cohomotopy* of spacetime/worldvolumes and *Cobordism* classes of submanifolds behaving like solitonic branes carrying the corresponding Cohomotopy charge [124, §2.2] [121, §2.1]:


The **Pontrjagin theorem** [112][83, §IX] identifies the unstable n -Cohomotopy of a closed manifold with the cobordism classes of its normally framed submanifolds of co-dimension n .




The **Cohomotopy charge** of a normally framed submanifold (aka *scanning map* or *Pontrjagin-Thom collapse*) is represented by mapping points of the ambient space to their directed distance if inside a tubular neighborhood, else to ∞ . Conversely, every Cohomotopy class is represented by a smooth map with 0 a regular value, whose pre-image is a normally framed submanifold with that Cohomotopy charge.




Under this relation, homotopy of charge maps corresponds to normal framed **cobordism** of submanifolds. The cobordism relation exhibits a form of pair creation/annihilation of submanifolds carrying opposite Cohomotopy charges.



When making more ambient dimensions available, the cobordism classes eventually (quickly) exhibit **stabilization** on abelian cobordism cohomology groups. (This might relate *Hypothesis H* to Vafa's *cobordism conjecture* cf. [124, §4]).




This “linearized” Cohomotopy/Cobordism is a **form of K-theory**: algebraic K-theory over the “absolute base field \mathbb{F}_1 ” (cf. [19, Thm. 5.9][9, Cor. 2.25]).




Thus flux quantization in Cohomotopy lifts to M-theory the same arguments that motivated topological K-theory in type II string theory: its character map reproduces the Bianchi identities & its equivalence relation models (anti-)brane pair-creation/annihilation.

Moduli space of soliton configurations. But the Pontrjagin theorem concerns only the total cohomotopical charge, identifying it with the *net* (anti-)brane content. Beyond that we have the whole *moduli space* of charges (considered now specialized to our 2D transverse space), and **Segal's theorem** [139] says that the cohomotopy charge map (scanning map) identifies this with a moduli space of brane positions, namely with the *group-completed configuration space of points* [21][153][77]:




where the *configuration space of points* is the space of finite subsets of \mathbb{R}^2 – here understood as the space of positions of cores of solitons of unit charge +1,




and its *group completion* $\mathbb{G}(-)$ is the topological completion of the topological partial monoid structure given by disjoint union of soliton configurations.

Naïvely this is given by also including **anti-solitons** in the form of configurations of \pm -charged points, topologized such as to allow for their pair annihilation/creation as shown in the left column on the right.


Remarkably, closer analysis reveals [105] that the group completion $\mathbb{G}(-)$ produces configurations of **strings** (extending parallel to one axis in \mathbb{R}^3) **with charged endpoints** whose pair annihilation/creation is smeared-out to string worldsheets as shown in the right column ([132, Fig. 2]):



This means (cf. [132, Prop. 3.14]) that the **vacuum-to-vacuum soliton scattering processes**, forming the loop space $\Omega \mathbb{G} \text{Conf}(\mathbb{R}^2)$, are identified with *framed links* ([106, p 15]); for instance



subject to *link cobordism* (cf. [91]):



Shown on the right are evident continuous deformations of paths in the above configuration space of charged strings, hence in the group-completed configuration space of points. Shown on the left are the local deformations of corresponding framed links, generating the relation of *link cobordism*.

It follows [132, Thm 3.17] that the charge of a soliton scattering process L is the sum over crossings of the *crossing number*

$$\# \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) = +1, \quad \# \left(\begin{array}{c} \nwarrow \\ \nearrow \end{array} \right) = -1,$$

which equals the linking+framing number:

$$\begin{array}{ccc} \Omega \mathbb{G} \text{Conf}(\mathbb{R}^2) & \xrightarrow{\sim} & \Omega \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2) \xrightarrow{[-]} \pi_3(S^2) \simeq \mathbb{Z} \\ L & \mapsto & \begin{array}{l} \text{total crossing number} = \\ \text{linking} + \text{framing number} \end{array} \rightarrow \#L \end{array}$$

But this is precisely the **Wilson loop observable** of L in (abelian) **Chern-Simons theory!** [132, §4]. This is what we explain next.


The k -Soliton sector. More generally, we may consider loops based in the k th connected component of the moduli space (cf. [132, Rem. 3.20]), corresponding to scattering process from k to k net number of solitons.

$$\begin{array}{ccc} \text{net charge } k & & \text{Hopf degree } k \\ \mathbb{G} \text{Conf}_k(\mathbb{R}^2) & \xrightarrow{\sim} & \text{Maps}_k^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2) \\ \downarrow & & \downarrow \\ \mathbb{G} \text{Conf}(\mathbb{R}^2) & \xrightarrow{\sim} & \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2) \end{array}$$


Since the double loop space $\text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2)$ admits the structure of a topological group, all these connected components have the same homotopy type, and hence these scattering processes L are again classified by the integer total crossing number $\#L$ which is the abelian Chern-Simons Wilson-loop observable.

$$\begin{array}{ccc} \Omega_k \mathbb{G} \text{Conf}(\mathbb{R}^2) & & \\ \downarrow & \searrow L \mapsto \#L & \\ \pi_0 \Omega_k \mathbb{G} \text{Conf}(\mathbb{R}^2) & \xrightarrow{\sim} & \mathbb{Z} . \end{array}$$


For instance, a generic $k = 3$ process looks like this:



and via the framed cobordism moves



it computes to the trivial scattering process accompanied by $\#L$ vacuum pair braiding processes:



Chern-Simons level. We will see below further meanings of the number k :

This integer k is equivalently $\left\{ \begin{array}{l} \text{the } \textit{number} \text{ of fractional quasi-hole vortices in a quantum Hall system,} \\ \text{the } \textit{level} \text{ of their effective abelian Chern-Simons theory,} \\ \text{the } \textit{maximal denominator} \text{ for filling fractions of their quantum states.} \end{array} \right.$

Generally, we will recover in a novel *non-Lagrangian* way the features of quantum Chern-Simons theory that are traditionally argued starting with the k th multiple of the local Lagrangian density $a \wedge da$ for a gauge potential 1-form a .

The situation on the 2-Sphere. Furthermore, consider k solitons on the actual 2-sphere S^2 . Here, the 2-Cohomotopy moduli space satisfies (cf. [61]):

$$\pi_0 \Omega_k \text{Maps}(S^2, S^2) \simeq \mathbb{Z}_{2|k|},$$

and the long homotopy fiber sequence induced by point evaluation shows that the generator of this cyclic group is again identified with the basic half-braiding operation:

$$\begin{array}{ccccccc} \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2) & \xrightarrow{\text{fiber of...}} & \text{Maps}(S^2, S^2) & \xrightarrow{\text{point-evaluation}} & S^2 & & \\ \underbrace{\pi_2(S^2)}_{\mathbb{Z}} & \xrightarrow{-2k} & \underbrace{\pi_0 \Omega_k \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^2, S^2)}_{\mathbb{Z}} & \longrightarrow & \underbrace{\pi_0 \Omega_k \text{Maps}(S^2, S^2)}_{\mathbb{Z}_{2|k|}} & \longrightarrow & \underbrace{\pi_1(S^2)}_1 \end{array}$$

$$\begin{array}{ccc} \begin{array}{c} \nearrow \\ \searrow \end{array} & \longmapsto & \left[\begin{array}{c} \nearrow \\ \searrow \end{array} \right] \end{array}$$

With flux-quantized fields being equipped with a classifying space \mathcal{A} , there is a neat way to directly obtain the topological quantum observables – via the following observation:

Topological flux observables in Yang-Mills theory – Theorem [131, §1]. For G -Yang-Mills theory on $\mathbb{R}^{1,1} \times \Sigma^2$, with a choice of Ad-invariant lattice $\Lambda \subset \mathfrak{g}$:

- (i) Non-perturbative quantization of the algebra of flux observables through the closed surface Σ^2 is given by the group C^* -algebra $\mathbb{C}[-]$ of the Fréchet-Lie group of smooth maps $\Sigma^2 \rightarrow G \times (\mathfrak{g}/\Lambda)$.
- (ii) The corresponding group algebra of topological observables (observing only the connected components of flux) coincides with the Pontrjagin homology algebra of pointed maps $(\mathbb{R}^1 \times \Sigma^2)_{\cup\{\infty\}} \rightarrow B(G \times (\mathfrak{g}/\Lambda))$:

$$\mathbb{C} \left[C^\infty(\Sigma^2, G) \times C^\infty(\Sigma^2, (\mathfrak{g}/\Lambda)) \right] \xrightarrow{\pi_0} \mathbb{C} \left[H^0(\Sigma^2; G) \times H^1(\Sigma^2; \Lambda) \right] \simeq H_0 \left(\text{Maps}^* \left((\mathbb{R}^1 \times \Sigma^2) \cup \{\infty\}, B(G \times (\mathfrak{g}/\Lambda)) \right); \mathbb{C} \right)$$

Non-perturbative quantum algebra of observables on flux through Σ^2
corresponding algebra of topological observables
Pontrjagin homology algebra of moduli space of soliton charges

For example, in electromagnetism, with $G = U(1)$ and $\Lambda := \mathbb{Z} \hookrightarrow \mathbb{R}$ this gives [131, §2]:


$$\mathbb{C} \left[\underbrace{H^1(\Sigma^2; \mathbb{Z})}_{\text{electric}} \times \underbrace{H^1(\Sigma^2; \mathbb{Z})}_{\text{magnetic}} \right] \simeq H_0 \left(\text{Maps}^* \left((\mathbb{R}^1 \times \Sigma^2) \cup \{\infty\}, \underbrace{BU(1) \times BU(1)}_{\text{classifying space for Dirac flux quantization}} \right); \mathbb{C} \right).$$

This allows us to generalize [131, §3,4]:

Topological flux observables of any higher gauge theory. For a higher gauge theory flux-quantized in \mathcal{A} -cohomology, the quantum algebra of topological flux observables on a spacetime of the form $\mathbb{R}^{1,1} \times \Sigma^{D-2}$ is the Pontrjagin homology algebra of the soliton moduli, hence in $\text{deg} = 0$ is the group algebra of vacuum soliton processes “on the light-cone”:

$$\begin{aligned} \text{Obs}_\bullet &:= H_\bullet \left(\text{Maps}^* \left((\mathbb{R}^1 \times \Sigma^{D-2}) \cup \{\infty\}, \mathcal{A} \right); \mathbb{C} \right) \\ &\simeq H_\bullet \left(\Omega \text{Maps}(\Sigma^{D-2}, \mathcal{A}); \mathbb{C} \right) \\ \text{Obs}_0 &= \mathbb{C} \left[\pi_0 \Omega \text{Maps}(\Sigma^{D-2}, \mathcal{A}) \right] \end{aligned}$$

For this, note that the star-involution is given by the *combination* of complex conjugation (time reversal) and loop reversal (hence x -reversal), where $\mathbb{R}^{1,1} \simeq \mathbb{R}\langle t, x \rangle$, and the operator product is given by loop concatenation:



4 The topological Quantum States

To summarize so far, we have seen that the topological sector of the flux-quantized phase space of solitons on magnetized M5-probes Σ wrapping Seifert orbi-singularities is

$$\text{Maps} \left(\begin{array}{c} \Sigma \quad \mathbb{C}P^3 \\ \downarrow, \downarrow \\ X \quad S^4 \end{array} \right)^{\mathbb{Z}_2} \simeq \text{Maps}^*/(\mathbb{R}^2_{\cup\{\infty\}} \wedge S^1, S^2) \simeq \Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2) \xrightarrow{[-]} \pi_0 \Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2) \simeq \mathbb{Z}$$

$$L \longmapsto \#L$$

topological sector of flux-quantized phase space
2-Cohomotopy cocycle space
loop space of group-completed configuration space
net charge


The **topological quantum states of this system** now follow [131][132, §4] by general algebraic quantum theory: The gauge-invariant topological **observables** form the (higher) homology of this space

$$\text{Obs}_\bullet := H_\bullet(\Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2); \mathbb{C})$$

making a (star-)algebra under concatenation (reversion) of loops — the *Pontrjagin algebra*.

$$\begin{array}{ccc} \Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2) & \xrightarrow[\text{rev}]{\text{loop reversal}} & \Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2) \\ H_\bullet(\Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2); \mathbb{C}) & \xrightarrow[\text{rev}_*]{\text{Pontr. antipode}} & H_\bullet(\Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2); \mathbb{C}) \\ & \searrow^{\text{Hermitian conjugation of quantum observables}} & \downarrow^{\overline{(-)} \text{ cmplx cnjgtn}} \\ & & H_\bullet(\Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2); \mathbb{C}) \end{array}$$

This means that time-reversal goes along with the reversal of looping around the M/IIA-circle, whence we are dealing with a version of *discrete light-cone quantization* in their topological sectors.



The basic ordinary (degree=0) observables detect the deformation class of a framed link L .


$$\begin{array}{l} \text{Obs}_0 \xrightarrow{\sim} \mathbb{C}[\pi_0(\Omega_0 \mathbb{G}\text{Conf}(\mathbb{R}^2))] \xrightarrow{\sim} \mathbb{C}[\mathbb{Z}] \\ \mathcal{O}_L := \delta_{[L]} = \delta_{\#L} \\ \mathcal{O}_L \cdot \mathcal{O}_{L'} = \delta_{L \sqcup L'} = \delta_{\#L + \#L'} \end{array} \quad (3)$$

Since these observables commute among each other, their *pure* topological quantum states are their (real & positive) algebra homomorphisms:

$$\begin{array}{l} \text{PureQStates}_0 \simeq \left\{ \rho : \text{Obs}_0 \xrightarrow{\text{homo}} \mathbb{C} \mid \rho \in \text{MixedQStates}_0 \right\} \\ \downarrow \text{on commuting observables} \\ \text{MixedQStates}_0 := \left\{ \rho : \text{Obs}_0 \xrightarrow{\text{linear}} \mathbb{C} \mid \forall_{\mathcal{O} \in \text{Obs}_\bullet} \left(\rho(\mathcal{O}^*) = \rho(\mathcal{O})^*, \rho(\mathcal{O}^* \cdot \mathcal{O}) \geq 0 \in \mathbb{R} \hookrightarrow \mathbb{C} \right), \rho(1) = 1 \right\}. \end{array}$$

reality
(semi-)positivity
normalization

Therefore pure topological states $|m\rangle$ are determined by an **anyonic phase** $\exp(\pi i/m)$ assigned to any crossing,



accumulating to the exponentiated crossing number

$$\begin{array}{ccc} \text{Obs}_0 & \xrightarrow{\langle m | - | m \rangle} & \mathbb{C} \\ \mathcal{O}_L & \mapsto & e^{\frac{\pi i}{m} \#L} \end{array}$$

The resulting expectation values

$$\langle m | \mathcal{O}_L | m \rangle = \exp\left(\frac{\pi i}{m} \#L\right) = \exp\left(\frac{\pi i}{m} \left(\sum_{i \neq j \in \pi_0(L)} \underset{\substack{\text{linking} \\ \text{numbers}}}{\text{lnk}(L_i, L_j)} + \sum_{i \in \pi_0(L)} \underset{\substack{\text{framing} \\ \text{numbers}}}{\text{frm}(L_i)} \right)\right)$$

are [132, §4] just those of *Wilson loop observables* in “spin” *Chern-Simons theory*, as **expected for abelian anyons!** For example:

$$\left\langle m \left| \begin{array}{c} \text{link} \\ \text{diagram} \end{array} \right| m \right\rangle = \left\langle m \left| \begin{array}{c} \text{link} \\ \text{diagram} \end{array} \right| m \right\rangle = \exp\left(\pi i \frac{3}{m}\right).$$

Applying the GNS-construction to such state produces a 1-dimensional Hilbert space

$$\overbrace{\mathbb{C}[\mathbb{Z}]} / (e^{\pi i/m} - \theta) \simeq \mathbb{C}, \quad (4)$$

which is as expected for the quantum states of abelian Chern-Simons theory on $\mathbb{R}_{U\{\infty\}}^2$. (More on this on p 21.)

Remark. At this point, $m \in \mathbb{R} \neq 0$ may be irrational, but its rationality will be enforced by requiring compatibility with states on more general domain surfaces, see p. 21 and p. 23.

Remark. These *solitonic* anyons are *not* yet the controllable/parameterized defect anyons that could be used for topological braid quantum gates operating by adiabatic movement of anyonic *defects* or (quasi-)holes. But the latter arise as defect points among the former, we come to this on p. 24.

Remark. The appearance of framed links along just the above lines is known in the condensed matter theory of anyonic defect lines in the 3D “8-band model” ([45, pp 15], following [142]): From this perspective, the Cohomotopy classifying space S^2 plays the role of the classifying space for electron band Hamiltonians on a crystal lattice.

Anyonic topological order on Flux-quantized M5-probes. We now identify the promised topological order on M5-probes flux-quantized in equivariant twistorial Cohomotopy, by considering M5s wrapping closed surfaces:

Anyonic quantum observables on closed surfaces. Consider now a *closed orientable surface* Σ_g^2 of genus $g \in \mathbb{N}$ to replace the previous factor $\mathbb{R}_{U\{\infty\}}^2$ in the brane diagram:

$$\Sigma^{1,5} := \mathbb{R}^{1,0} \times \Sigma_g^2 \times S^1 \times \mathbb{R}_{\text{sgn}}^2$$

Directly analogous analysis as before gives that the topological quantum observables on the flux-quantized self-dual tensor field form the group algebra of the fundamental group of the 2-cohomotopy moduli space in the k th connected component

$$\text{Obs}_0(\Sigma_g^2) := H_0\left(\Omega_k \text{Maps}(\Sigma_g^2, S^2); \mathbb{C}\right) \simeq \mathbb{C}\left[\pi_0 \Omega_k \text{Maps}(\Sigma_g^2, S^2)\right], \quad (5)$$

where $k \in \mathbb{N}$ is the degree of the classifying maps, corresponding under the Pontrjagin theorem to a net number of k (anti-)solitons on Σ_g^2 .

Theorem (using [61, Thm 1][84, Thm 1][76, Cor 7.6]). This group of 2-cohomotopy charge sectors is identified as *twice* the integer Heisenberg group extension (cf. [86]) of \mathbb{Z}^{2g} by $\mathbb{Z}_{2|k|}$:⁶

$$\pi_0 \Omega_k \text{Maps}(\Sigma_g^2, S^2) \simeq \left\{ \left(\vec{a}, \vec{b}, [n] \right) \in \mathbb{Z}^g \times \mathbb{Z}^g \times \mathbb{Z}_{2|k|}, \quad \begin{array}{l} (\vec{a}, \vec{b}, [n]) \cdot (\vec{a}', \vec{b}', [n']) = \\ (\vec{a} + \vec{a}', \vec{b} + \vec{b}', [n + n' + \vec{a} \cdot \vec{b}' - \vec{a}' \cdot \vec{b}]) \end{array} \right\} =: \widehat{\mathbb{Z}^{2g}}$$

Ground state degeneracy. Hence the observable group-algebra Obs_0 for $g = 1$, $\Sigma_1^2 = T^2$, has generators

$$\left\{ \begin{array}{l} W_a := (1, 0, [0]) \\ W_b := (0, 1, [0]) \\ \zeta := (0, 0, [1]) \end{array} \right\}$$

⁶Here $\mathbb{Z}_n := \mathbb{Z}/(n)$ (with $\mathbb{Z}_0 = \mathbb{Z}$) are the (in-)finite cyclic groups.

subject to the relations

$$\left\{ \begin{array}{l} W_a \cdot W_b = \zeta^2 W_b \cdot W_a \\ \zeta^{2k} = 1 \\ [\zeta, -] = 0 \end{array} \right\}.$$

This algebra is just the observable algebra expected [144, (5.28)] for anyonic topological order on the torus as described by abelian Chern-Simons theory at *level* k . The non-trivial irreps have:

- dimension k , this being the expected *ground state degeneracy* on the torus,
- are labeled by $\nu := p/k$, $p \in \{1, 2, \dots, k\}$, as expected for fractional *filling factors*.

Hilbert space of
quantum states
on the torus

$$\mathcal{H}_{T^2} := \text{Span}(|[n]\rangle, [n] \in \mathbb{Z}_{|k}|) \in \text{Obs}_0(T^2)\text{Modules}, \quad \dim(\mathcal{H}_{T^2}) = k,$$

$$W_a|[n]\rangle := e^{2\pi i\nu} |[n]\rangle$$

$$W_b|[n]\rangle := |[n+1]\rangle$$

$$\zeta|[n]\rangle := e^{\pi i\nu} |[n]\rangle.$$

Modular equivariance. Strikingly, in this construction modular symmetry **is manifest**, since the looped mapping space is canonically acted on by the mapping class group MCG of Σ_g^2 (cf. [33, §2.1]), simply by precomposition of maps! Inspection of the above theorem (cf. [61, bottom of p 153]) shows that this MCG -action identifies indeed as the canonical action of $\text{Sp}_{2g}(\mathbb{Z})$ on $\widehat{\mathbb{Z}^{2g}}$.

$$\begin{array}{ccc} \overbrace{\pi_0 \text{Homeos}_{\text{or}}(\Sigma_g^2)}^{\text{MCG}(\Sigma_g^2)} & \hookrightarrow & \pi_0 \Omega_k \text{Maps}(\Sigma_g^2, S^2) \\ \downarrow [33, \S 6.3] & & \wr \\ \text{Sp}_{2g}(\mathbb{Z}) & \hookrightarrow & \widehat{\mathbb{Z}^{2g}} \end{array}$$

Hence, we may ask for a lift of the $\widehat{\mathbb{Z}^{2g}}$ action on quantum states to an action of the semidirect product $\widehat{\mathbb{Z}^{2g}} \rtimes \text{Sp}_{2g}(\mathbb{Z})$. For $g = 1$ one readily checks that this gives the modular transformations of states known [93, p 65] from abelian Chern-Simons theory:

$$\begin{array}{l} \text{modular} \\ \text{action on observables} \\ \text{and on states} \end{array} \quad m(W) \cdot m(|[n]\rangle) = m(W|[n]\rangle), \quad \forall \left\{ \begin{array}{l} m \in \text{Sp}_{2g}(\mathbb{Z}) \\ W \in \widehat{\mathbb{Z}^{2g}} \\ |[n]\rangle \in \mathcal{H}_g \end{array} \right. \\ S(|[n]\rangle) = \frac{1}{\sqrt{|k|}} \sum_{[\hat{n}]} e^{2\pi i \frac{n\hat{n}}{k}} |[\hat{n}]\rangle, \quad T(|[n]\rangle) = e^{i\pi \frac{n^2}{k}} |[n]\rangle.$$

Generally, writing $(\vec{e}_i \in \mathbb{Z}^g)_{i=1}^g$ for the canonical basis vectors, the observable group-algebra Obs_0 for general g has generators

$$\left\{ \begin{array}{l} W_a^i := (\vec{e}_i, 0, [0]) \\ W_b^i := (0, \vec{e}_j, [0]), 1 \leq i \leq g \\ \zeta := (0, 0, [1]) \end{array} \right\}$$

subject to the relations

$$\left\{ \begin{array}{l} W_a^i \cdot W_b^j = \delta^{ij} \zeta^2 W_b^j \cdot W_a^i \\ \zeta^{2k} = 1 \\ \text{all other commutators vanish} \end{array} \right\}.$$


Requiring the reps \mathcal{H}_g of this algebra to analogously support modular equivariance requires them to have dimension $|k|^g$ — which is the result expected [93, p 40] for abelian topological order on Σ_g^2 :

Hilbert space of
quantum states
on genus= g surface

$$\mathcal{H}_{\Sigma_g^2} \in \text{Obs}_0(\Sigma_g^2)\text{Modules}, \quad \dim(\mathcal{H}_{\Sigma_g^2}) = |k|^g,$$

Here, the generators $W_{a,b}^i$ correspond to the classical generators of the surface's fundamental group. **Oriented closed surfaces** are all obtained (cf. [47, p 100]) by identifying in the regular $4g$ -gon, for *genus* $g \in \mathbb{N}$:

- (i) all boundary vertices with a single point;
and, going clockwise for $r \in \{0, \dots, g-1\}$,
- (ia) the $(4r+1)$ st boundary edge with the reverse of the $(4r+3)$ rd,
- (iib) the $(4r+2)$ nd boundary edge with the reverse of the $(4r+4)$ th.



In other words, the homotopy type of the surface sits in a (pointed) homotopy co-fiber sequence of this form:

$$S^1 \xrightarrow{\prod_i [a_i, b_i]} \bigvee_g (S_a^1 \vee S_b^1) \longrightarrow \Sigma_g^2 \xrightarrow{\delta} S^2$$

whence its fundamental group is the quotient of the free group on $2g$ generators $(a_i, b_i)_{i=1}^g$ by the normal subgroup generated by that polygon's boundary:

$$\pi_1(\Sigma_g^2) \simeq \langle a_1, b_1, \dots, a_g, b_g \rangle / \prod_i [a_i, b_i]$$

2-Cohomotopy moduli of oriented closed surfaces. Mapping this co-fiber sequence into S^2 and applying $\pi_0 \Omega_k$, it collapses [61, Prop. 2] to twice [84, Thm 1] the integer Heisenberg central extension of \mathbb{Z}^{2g} by $\mathbb{Z}_{2|k|}$:


$$1 \rightarrow \underbrace{\pi_0 \Omega_k \text{Maps}(S^2, S^2)}_{\mathbb{Z}_{2|k|}} \xrightarrow{\delta^*} \underbrace{\pi_0 \Omega_k \text{Maps}(\Sigma_g^2, S^2)}_{\text{integer Heisenberg group}} \longrightarrow \underbrace{\pi_0 \Omega_* \text{Maps}^*(\bigvee_g (S_a^1 \vee S_b^1), S^2)}_{\mathbb{Z}^{2g}} \rightarrow 1.$$

The phase generators. Hence these integer Heisenberg groups inject into each other as the surfaces are surjected onto each other by collapsing pairs of 1-cycles.

$$\begin{array}{ccc} \Sigma_g^2 & \xleftarrow{p} & \Sigma_{g+1}^2 \\ \pi_0 \Omega_k \text{Maps}(\Sigma_g^2, S^2) & \xrightarrow{\pi_1(p^*, k)} & \pi_0 \Omega_k \text{Maps}(\Sigma_{g+1}^2, S^2) \\ \wr & & \wr \\ \widehat{\mathbb{Z}^{2g}} & \xleftarrow{\quad} & \widehat{\mathbb{Z}^{2(g+1)}} \end{array}$$


Thereby, their central generator ζ represents the previously identified half-braiding operation of solitons on these surfaces. This is the “reason” for the central extension being by $\mathbb{Z}_{2|k|}$ instead of just $\mathbb{Z}_{|k|}$:

The phase generator ζ does not correspond to full rotations (such as around the square on the right) but to “particle exchange” by half-braiding — as expected for anyons.



Non-orientable closed surfaces are all obtained by identifying in the regular $2h$ -gon, for *crosscap number* $h \in \mathbb{N}_{\geq 1}$:

- (i) all boundary vertices with a single point and, going clockwise for $r \in \{0, \dots, h-1\}$,
- (ii) the $(2r+1)$ st boundary edge with the reverse of the $(2r+2)$ nd.



In other words, the homotopy type of the surface sits in a (pointed) homotopy co-fiber sequence of this form:

$$S^1 \xrightarrow{\prod_i a_i^2} \bigvee_h S^1 \longrightarrow \Sigma_h^2 \xrightarrow{\delta} S^2$$

2-Cohomotopy moduli of non-orientable closed surfaces. Mapping this co-fiber sequence into S^2 and applying $\pi_0 \Omega_k$, it induces [61, Prop. 3] an extension of \mathbb{Z}^{h-1} by \mathbb{Z}_2 which as such is trivial [84, Thm. 2]:

$$1 \rightarrow \underbrace{\text{coker}\left(\left(\Sigma \prod_i a_i^2\right)^*\right)}_{\mathbb{Z}_2} \xrightarrow{\delta^*} \underbrace{\pi_0 \Omega_k \text{Maps}^*\left(\Sigma_h^2, S^2\right)}_{\mathbb{Z}_2 \times \mathbb{Z}^{h-1}} \longrightarrow \underbrace{\ker\left(\left(\prod_i a_i^2\right)^*\right)}_{\mathbb{Z}^{h-1}} \rightarrow 1.$$


Again, the exponent appearing, $h-1$, is just that expected for abelian Chern-Simons ground state degeneracy, where (cf. [16, (73)]):

$$\dim(\mathcal{H}_{\Sigma_h^2}) = |k|^{h-1}.$$

Defects via punctured worldvolumes. It is now immediate to bring *adiabatically movable defect anyons* into the picture, missing in traditional discussion but crucially needed for topological quantum gates (cf. [99, §3]). Namely, we may simply further generalize the surfaces Σ_g^2 to their *n-punctured* versions, obtained by deleting the positions of a subset of points – thus literally creating defects!

$$\Sigma_{g,n}^2 := \Sigma_{g,n}^2 \setminus \{s_1, \dots, s_n\}$$

for $\{s_1, \dots, s_n\} \subset \Sigma_g^2$



That these defects are void of the dynamical solitons is elegantly enforced by identifying all their positions with the point-at-infinity (where, recall, the soliton's very nature is to not be present):


$$\text{domain for solitons in the presence of } n \text{ defects} = \text{one-point compactification of } n\text{-punctured surface} = (\Sigma_{g,n}^2)_{\cup\{\infty\}} \quad \text{e.g.:} \quad (\Sigma_{0,1}^2)_{\cup\{\infty\}} \simeq \mathbb{R}_{\cup\{\infty\}}^2.$$

In this generality, our previous brane diagram now is the following, with algebra of soliton quantum observables as shown, by the same kind of argument as before:

$$\left(\begin{array}{c} \mathbb{Z}_2 \\ \downarrow \\ \Sigma^{1,6} \end{array}\right) := \mathbb{R}^{1,0} \times (\Sigma_{g,n}^2)_{\cup\{\infty\}} \times S^1 \times \mathbb{R}_{\text{sgn}}^2 \rightsquigarrow \text{Obs}_0(\Sigma_{g,n}^2) := H_0\left(\Omega_k \text{Maps}^*\left((\Sigma_{g,n}^2)_{\cup\{\infty\}}, S^2\right); \mathbb{C}\right).$$

Braid group action. This algebra of observables is faithfully acted on by the mapping class group of the punctured surface – again simply by precomposition of maps.

But, with punctures, that group is now an extension (cf. [94, Thm. 3.13]) of the plain mapping class group by the *surface braid group* that acts by (“adiabatically”) moving the defects around each other!



$$1 \rightarrow \text{Br}_n(\Sigma_g^2) \hookrightarrow \pi_0 \text{Homeos}_{\text{or}}^*\left((\Sigma_{g,n}^2)_{\cup\{\infty\}}\right) \twoheadrightarrow \text{MCG}(\Sigma_g^2) \rightarrow 1$$

surface braid group
mapping class group of punctured surface
mapping class group of plain surface

In deducing this, we used that $\text{Homeos}^*\left((\Sigma_{g,n}^2)_{\cup\{\infty\}}\right) \simeq \text{Homeos}(\Sigma_{g,n}^2)$, since $(-)\cup\{\infty\}$ is functorial on homeos.

Concretely, observe that the homotopy type of the one-point compactification of a punctured closed surface is the wedge sum of the original surface with $n-1$ circles (cf. [65, p 11], whose graphics we are adapting):

$$(\Sigma_{g,n}^2)_{\cup\{\infty\}} \simeq \Sigma_g^2 \vee \bigvee_{n-1} S^1 \quad (6)$$

This means that the punctures are effectively topology changing defects, as such similar to the *genon*-ic anyon

defects considered in [8], and it implies that their bare quantum observables are (the group algebra) of:

$$\pi_1 \text{Maps}^* \left((\Sigma_{g,n}^2)_{\cup \{\infty\}}, S^2 \right) \simeq \pi_1 \text{Maps}^* (\Sigma_g^2, S^2) \times \mathbb{Z}^{n-1} \underset{g=0}{\simeq} \mathbb{Z}^n, \quad (7)$$

where on the right we recognize that associated with each of the n punctures is one copy of the braid phase observable algebra (3) for the nearby solitonic anyons. This observable algebra of defects gets further enhanced by the corresponding mapping class group:

Side-remark: Defect-braiding on M5s as a quantum-gravitational effect. Noting that the mapping class group is equivalently the group of large *diffeomorphisms* of the punctured surface (cf. [33, p 45]),

$$\pi_0 \text{Homeos}_{\text{or}}^* \left((\Sigma_{g,n}^2)_{\cup \{\infty\}} \right) \simeq \pi_0 \text{Diffeos}_{\text{or}} (\Sigma_{g,n}^2), \quad (8)$$

we see that braiding of anyonic defects is reflected in equipping the moduli spaces of cohomotopical charges on the brane worldvolume with the action by diffeomorphisms, hence by passing to the action *groupoid* of moduli quotiented by diffeos:

$$\text{GnrlCovariantModuli}(\Sigma) \simeq \text{Moduli}(\Sigma) // \text{Diffeos}(\Sigma).$$

But this is the hallmark of *generally covariant* systems (cf. [29]), such as are our probe M5-branes.

Ultimately we are to consider surfaces $\Sigma_{g,n,b}^2$ that may feature $b \in \mathbb{N}$ boundary components, and then determine the normal subgroup *pure gauge* diffeomorphisms inside (8) which are trivial on the boundary. The resulting quotient group will be the (ADM/BSM-like) group of diffeos that serve in practice as experimentally observable boundary charges.

Observables on soliton + defect anyons. So the covariantized quantum observables on the sphere, but now in the presence of n defects is the group algebra of the spherical *framed braid group* [82], namely of the *wreath product* $\mathbb{Z} \wr \text{Br}_n$ (cf. [11, §8])

- of the braid-phase group \mathbb{Z} of the solitonic anyons as before in (3)
- with the actual braid group $\text{Br}_n(\Sigma_0^2)$ of the defect anyons,

namely of the semidirect product group

$$\mathbb{C}[\text{FBr}_n(\Sigma_0^2)] = \mathbb{C}[\mathbb{Z} \wr \text{Br}_n(\Sigma_0^2)] = \mathbb{C}[\mathbb{Z}^n \rtimes \text{Br}_n(\Sigma_0^2)], \quad (9)$$

where the braid group acts on \mathbb{Z}^n through permutation of the factors Sym_n of the defect anyons:

$$\mathbb{Z}^n \rtimes \text{Br}_n(\Sigma_0^2) \longrightarrow \mathbb{Z}^n \rtimes \text{Sym}_n \simeq \left\{ ((n_i)_{i=1}^n, \sigma) \mid ((n_\bullet), \sigma) \cdot ((n'_\bullet), \sigma') = ((n_\bullet + n'_{\sigma(\bullet)}), \sigma\sigma') \right\}. \quad (10)$$

Just such *para-statistical* (cf. [149]) wreath-group statistics of defect anyons is seen in condensed matter [45].

Anyons as seen in Cohomotopy	Nature	Number	Braiding
Solitonic anyons	concentrations of flux density	net charge, CS-level: k	by (LC-)time evolution
Defect anyons	punctures in worldvolume	n in $\Sigma_{g,n}^2$	by worldvolume diffeomorphisms

The diagram shows a 2D worldvolume surface with a dashed boundary. On the left, a blue arrow points to a green oval labeled 'field solitons/quasi-particles/-holes/vortices: fmd submanifolds'. On the right, a red arrow points to a white circle labeled 'flux-expelling defects: punctures in the surface'. There are three green ovals and one white circle on the surface.

Quantum states of defect anyons. By (6) the worldvolume domains with n defects retract onto those with a single defect, whence so do the corresponding quantum observables (7):

$$\begin{array}{c} \xrightarrow{\text{id}} \\ \text{Obs.} \left((\Sigma_{0,1}^2)_{\cup \{\infty\}} \longleftarrow (\Sigma_{0,n}^2)_{\cup \{\infty\}} \longrightarrow (\Sigma_{0,1}^2)_{\cup \{\infty\}} \right) \\ \mathbb{Z} \longleftarrow \mathbb{Z}^n \longrightarrow \mathbb{Z} \end{array}$$

from which the 1-dimensional state spaces (4) associated with a single defect make the Hilbert space for n defects be their direct sum:

$$\mathcal{H}_{\Sigma_{0,n}^2} \simeq \mathbb{C}^n \simeq \langle e_1, e_2, \dots, e_n \rangle$$

with the braid group acting by permutation of summands (10). This *permutation representation* is evidently unitary with respect to the standard inner product on \mathbb{C}^n and as such reducible to the direct sum of the 1d trivial and the unitarization of the “*standard representation*”

$$\mathcal{H}_{\Sigma_{0,n}^2} \simeq \langle e_1 + e_2 + \dots + e_n \rangle \oplus \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & & & \\ \hline \end{array}$$

Topologically protected rotation gates. Via the cyclic subgroup $\mathbb{Z}_n \subset \text{Sym}_n$ of cyclic permutations, this standard representation contains what in quantum computing are known as qdit-based *rotation gates* [158]. For example, for $n = 3$ inspection readily shows that the unitarized standard representation is generated from a Pauli *Z-gate* and a qbit-rotation around the (conventional) *y-axis*, like this:


$$\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array} \simeq \left\{ (213) \mapsto \underbrace{\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}}_Z, (231) \mapsto \underbrace{\begin{bmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix}}_{R_y(2\alpha)} \text{ where } \alpha = 4\pi/3 \right\},$$

which implies at once that in the standard rep $\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \square & & & & \\ \hline \end{array}$ of Sym_6 we find also the corresponding qbit-*controlled* rotation, and so on.

Together with the global phase rotations of solitonic anyons given by the first wreath factor in (9), (controlled), such rotation gates are the workhorse in the quantum Fourier transform [102, §5][148, §3.2.1] (hence notably in Shor’s algorithm for prime factorization) and their precision and error protection is a major bottleneck in the implementation of useful quantum algorithms (cf. [44, §III]).

Here we see that our geometric engineering predicts the relevant gates to have topologically error-protected implementation by braiding of defects in FQH systems.

Topological rotation gates, obtained by cyclic braiding of defect anyons, combined with the global phase rotations given by braiding of solitonic anyons, would provide intrinsically exact and topologically protected gates of the kind that make up the quantum Fourier transform (in qdit-bases), and with it many other quantum algorithms.



Side-remark: “Parastatistics” as the most stable anyon braid gates. Such braid representations on irreps of the symmetric group have traditionally received little to no attention in topological quantum computing (popular are instead solutions to the Knizhnik-Zamolodchikov equation, cf. [129], and of the Yang-Baxter equation). Elsewhere they are discussed as speculative *parastatistics* [62][111] of fundamental particles instead of as adiabatic Berry-transformations of defect anyons. Therefore Jordan 2010 [73], who is the first to propose symmetric irreps as a model for quantum computation – aka *permutational quantum computing* [74] –, admitted “not [to] worry too much about the physical justification for the model” [73, p 109].


This seems to be a blind spot in the literature: Irreps of Sym_n are in particular surface braid representations via the surjection $\text{Br}_n(\Sigma_g^2) \twoheadrightarrow \text{Sym}_n$ — regarded as anyon braid gates they are in fact the *most stabilized* such, in that the gate operation is independent not just of isotopy but even of homotopy of the adiabatic transformation.

5 Conclusion: Better Anyon Theory

New theory of anyonic topological order, engineered on flux-quantized M5s. In summary, we have seen that global completion by flux-quantization of 11D supergravity with M5-probes (here: in equivariant twistorial cohomotopy – “Hypothesis H”), makes the quantized topological sector of the self-dual tensor field on M5-probes (wrapping Seifert orbi-singularities) reproduce key phenomena of abelian Chern-Simons theory thought of as an effective field theory for abelian anyons in fractional quantum Hall (FQH) systems:

(i) **Flux tubes bound to anyons.** The central assumption in the traditional heuristic understanding of the FQHE is that the anyonic solitons have flux quanta “attached” to them [140, p 883]. It is crucially this assumption that motivates and justifies abelian Chern-Simons theory as an effective field theory for FQH anyons, since variation of the sum of the abelian Chern-Simons term with the standard source term predicts that the gauge field flux is localized at the source particles (cf. [144, (5.25)][154, (3.6)]).

In contrast, in the present approach **this effect is a consequence of cohomotopical flux-quantization**, via the Pontrjagin theorem: The classifying map of the 2-Cohomotopy charge identifies an open neighborhood of each anyon with the 2-sphere minus its point at infinity, and the flux density F_2 is the pullback of the sphere's volume form along this map (cf. p 39), hence supported on just these open neighborhoods.



(ii) **Anyons subject to each other's Aharonov-Bohm phases.** Traditional discussion furthermore assumes from these attached flux tubes that the anyons must pick up Aharonov-Bohm quantum phases when circling around each other. While this is plausible, rigorous quantum field-theoretic derivation of this statement may not have found much attention.


In contrast, in the approach discussed here, this effect is again a direct consequence of cohomotopical flux-quantization, now via algebro-topological theorems of Segal and others, which serve to identify the cohomotopy charge moduli space with configuration spaces of soliton cores, whose fundamental group reflects the anyon braid phases (and thereby also the ground state degeneracy / topological order).

$$\begin{array}{ccc}
 \pi_0 \text{Maps}^*(\mathbb{R}^2_{\cup\{\infty\}}, S^2) & \xrightarrow{\sim} & \pi_0 \text{Maps}^*(\mathbb{R}^2_{\cup\{\infty\}}, B^2\mathbb{Z}) \\
 \downarrow \cong & \text{same net charges...} & \downarrow \cong \\
 \mathbb{Z} & & \mathbb{Z} \\
 \\
 \pi_1 \text{Maps}^*(\mathbb{R}^2_{\cup\{\infty\}}, S^2) & \xrightarrow{\quad} & \pi_1 \text{Maps}^*(\mathbb{R}^2_{\cup\{\infty\}}, B^2\mathbb{Z}) \\
 \downarrow \cong & \text{...but different moduli} & \downarrow \cong \\
 \pi_1 \mathbb{G}\text{Conf}(\mathbb{R}^2) & & 1 \\
 \text{config space} & & \text{no structure}
 \end{array}$$

Note how both these effects come about by changing the traditional flux-quantization of the Chern-Simons field from the classifying space for complex line bundles to just its first “cell”. This preserves the quantization of charges but makes their moduli exhibit anyonic effects.

$$\begin{array}{ccc}
 S^2 \simeq \mathbb{C}P^1 & \xleftarrow{\text{1st cell inclusion}} & \mathbb{C}P^\infty \simeq B^2\mathbb{Z} \\
 \text{classifying space} & & \text{classifying space for} \\
 \text{for 2-Cohomotopy} & & \text{ordinary 2-cohomology}
 \end{array}$$

(iii) **Topological order.** The traditional way of establishing topological order is by applying geometric quantization to Wilson line observables, with respect to some effective action, which is a somewhat convoluted process involving ad-hoc choices and regularizations. In contrast, in the approach discussed here, the quantum observables obtain immediately, without further choices, from the topological light-cone quantization of the flux-quantized moduli space (as its Pontrjagin homology algebra).



Here the looping Ω_k that drives this quantum dynamics reflects dependence of moduli on the M/IIA circle.(!)

(iv) **Defect anyons** — as opposed to the solitonic anyons tracing out “Wilson lines” — seem to have previously found little to no attention in quantum Hall theory in general and its effective abelian Chern-Simons theories in particular. And yet, it is only such classically parameterized and hence, in principle, externally controllable defect anyons which may support braid quantum gates as envisioned in topological quantum computation.

In our approach, defect braiding emerges just as readily as the solitonic anyons, as a mild kind of quantum gravitational effect on M5-worldvolumes having a punctured surface factor space. This may be seen as a theoretical prediction of defect anyons in quantum Hall systems which might inform future search for experimental realization.

Summary of results:

On super-space, the equations of motion of **11D supergravity** with magnetic $1/2$ BPS **M5-brane** probes are equivalent to these Bianchi identities on the super-flux densities:

A-field	$d F_2^s = 0$	$d G_4^s = 0$	C-field
self-dual B-field	$d H_3^s = \phi_s^* G_4^s + \theta F_2^s F_2^s$	$d G_7^s = \frac{1}{2} G_4^s G_4^s$	dual C-field
M5 probe	$\Sigma^{1,5} 2\cdot 8_+ \xrightarrow[1/2\text{BPS immersion}]{\phi_s} X^{1,10} 32$		SuGra bulk

One admissible choice of **flux-quantization** law (the simplest in number of CW cells) is **twistorial Cohomotopy**, where the charges are classified by dashed maps like this:

$$\begin{array}{ccc}
 \text{M5-brane worldvolume} & \Sigma^{1,5} & \xrightarrow[\text{A- \& B-field charge}]{(a_1, b_2)} \mathbb{C}P^3 \\
 \downarrow \phi & \downarrow \text{immersion} & \downarrow t_H \\
 \text{bulk spacetime} & X^{1,10} & \xrightarrow[\text{C-field charge}]{(c_3, c_6)} S^4
 \end{array}$$

For (very good) $G \subset \text{Sp}(2)$ -orbifold domains, these maps are to be G -equivariant.

This flux-quantization implies a list of topological effects expected in M-theory.
 \Rightarrow **Hypothesis H:** This is the right choice of flux-quantization for M-theory.

Choosing (“engineering”) the M5-probe to be:

M5-brane probe worldvolume	2-brane worldvolume hosting anyonic solitons	M-theory circle	cone orbifold
$\Sigma^{1,5}$	$= \mathbb{R}^{1,0} \times \Sigma_{g,n}^2$	$\times S^1$	$\times \mathbb{R}_{\text{sgn}}^2 // \mathbb{Z}_2$

the **moduli space of solitons** becomes:
 $\text{Moduli} \simeq \text{Maps}^* \left((\mathbb{R}^1 \times \Sigma_{g,n}^2) \cup \{\infty\}, S^2 \right)$

The algebra of **topological quantum observables** on these solitons is:

$$\text{Obs}_0 := \underset{\text{topological quantum observables}}{H_0} \left(\underset{\text{Pontrjagin homology algebra}}{\text{Maps}^* \left((\mathbb{R}^1 \times \Sigma_{g,n}^2) \cup \{\infty\}, S^2 \right)}; \mathbb{C} \right) \simeq \underset{\text{group algebra}}{\mathbb{C} \left[\pi_1 \text{Maps}^* (\Sigma_{g,n}^2, S^2) \right]},$$

acted on by large diffeomorphisms (**general covariance** on the brane):

$$1 \xrightarrow{\text{braid group}} \text{Br}_n(\Sigma_g^2) \xleftarrow{\text{large diffeomorphism group}} \pi_0 \text{Homeos}_{\text{or}}^* \left((\Sigma_{g,n}^2) \cup \{\infty\} \right) \xrightarrow{\text{mapping class group}} \text{MCG}(\Sigma_g^2) \xrightarrow{1} 1$$

The corresponding **topological quantum states**:

- | | | |
|---|---|-----------------------------------|
| on $\Sigma_{0,0}^2 = S^2$ | reflect abelian braiding of solitonic anyons | } as for abelian Chern-Simons QFT |
| on $\Sigma_{g,0}^2 = \Sigma_{1,0}^2 \# \dots \# \Sigma_{1,0}^2$ | have k^g -fold degeneracy: topological order | |
| on $\Sigma_{1,0}^2 = \mathbb{T}^2$ | exhibit irred $\text{SL}_2(\mathbb{Z})$ - modular equivariance | |
| on $\Sigma_{0,n}^2 = S^2 \setminus \{z^1, \dots, z^n\}$ | reflect abelian braiding of defect anyons | |
- new & needed for topological quantum gates!

A broad lesson following immediately from our successful geometric engineering of topological qubits is the plausible existence of more exotic anyonic states than traditionally envisioned: Namely the “duality symmetry” [110][28, §6] of M-theory predicts that any geometrically engineered quantum system has “dual” incarnations with isomorphic quantum observables but entirely different geometric realization, where ordinary space is replaced by more abstract parameter spaces. Notably “T-duality” [147][37][54] applied to topological quantum materials has been argued [95][96][60] to exchange the roles of ordinary space with that of reciprocal “momentum space”.


(2) Novel experimental pathways towards anyons. Indeed, while anyonic solitons are traditionally envisioned as being localized in “position space” (meaning that the anyon cores are points in the plane of the crystal lattice) the physical principle behind topological quantum gates — namely [3][4][46, p 6][108, p 50] the *quantum adiabatic theorem* [118] — is unspecific to position space and only requires the material’s Hamiltonian to *depend on any continuous parameters* (such as external voltage or strain) varying in any abstract parameter space.

The general physical conditions for topological quantum gates given by the *quantum adiabatic theorem*, listed (a) - (e) on the right, are much more general than traditionally considered for anyon braid gates — the latter are only the special case where the parameters are configurations of points in the plane of the 2D crystal lattice.

- (a) **Ground state degeneracy** (when frozen at absolute zero, the system still has more than one state to be in, even up to phase).
- (b) **Spectral gap** (quanta of energy smaller than a given gap $\epsilon > 0$ cannot excite these ground states).
- (c) **Control parameters** (the above properties hold for a range of continuously tunable external parameters).
- (d) **Parameter topology** (there exist closed parameter paths that cannot be continuously contracted).
- (e) **Local invariance** (continuously deformed parameter paths induce the same transformation on ground states).

This means that, in principle, the possibilities in which anyonic quantum states could arise in the laboratory are far more general than what has been explored to date.

Concretely, a key example of alternative parameters for ground states of a quantum material are points in their reciprocal *momentum space*: This is the space of (quasi-)momenta, hence of wave-vectors for plane quasi-particle waves going through the crystalline material.




We have observed before that candidate anyon-like solitons localized (not in position space but) in momentum space are plausible both theoretically [129] as well as experimentally [157][143][72] and may have been hiding in plain sight: as band nodes of (interacting) topological semimetals.

Indeed, momentum space naturally features key properties that are typically assumed for anyon braid gates but remain elusive in position space:

(i) toroidal base topology is routinely assumed [150][152][88] in order to achieve the required ground-state degeneracy, but is quite unrealistic in position space, even more so when meant to be punctured by defect anyons — while the momentum space of a crystal is automatically a torus (the *Brillouin torus*).

(ii) stable defect points need special engineering in position space but arise automatically in momentum space in the guise of *band nodes* of topological semi-metals [129, Fig. 6]

(iii) defect point movement in a *controlled* way is necessary for braid gates but remains elusive in position space, while band nodes in momentum space have already shown to be movable in a variety of systems, by tuning of external parameters (e.g., strain).



The geometric engineering of anyons discussed here goes towards providing also fundamental theoretical underpinning of the possibility of more “exotic” anyon realizations than have traditionally been envisioned.

6 Digest for Algebraic Topologists

We are concerned with **algebraic-topological phenomena** arising when **magnetic flux** penetrates a semi-conducting surface Σ^2 .


The “gauge group” of the electromagnetic field is $G \equiv U(1)$ and *ordinarily* such flux is classified by maps to $BU(1) \simeq CP^\infty$.

Precisely, when quantum-effects are being resolved, then:

Theorem [131] (Yang-Mills flux quantum observables):

For ordinary gauge fields on a spacetime $\simeq \mathbb{R}^{1,1} \times \Sigma^2$ the **quantum observables of field flux** through Σ^2

form the group-convolution C^* -algebra $\mathbb{C}[C^\infty(\Sigma^2, G \ltimes (\mathfrak{g}/\Lambda))]$ for $\Lambda \subset \mathfrak{g}$ an Ad-invariant lattice. **quantum flux observables**



G Lie group (“gauge group”)
 \mathfrak{g} its Lie algebra

Commercial-value quantum computing will require **robust** quantum observables, insensitive to local fluctuations, only depending on **topological sectors** of field configurations.

$\mathbb{C}[C^\infty(\Sigma^2, G \ltimes (\mathfrak{g}/\Lambda))]$ $\xleftarrow{[-]^*}$ $\mathbb{C}[\pi_0 C^\infty(\Sigma^2, G \ltimes (\mathfrak{g}/\Lambda))]$
 all quantum flux observables **robust topological observables**

$C^\infty(-, -)$ manifold of smooth functions
 $(-) \ltimes (-)$ semidirect product via adjoint
 $\mathbb{C}[-]$ group convolution C^* -algebra
 $\pi_0(-)$ path-connected components

Proposition [131] (topological sector observables):

The topological flux quantum observables form the homology Pontrjagin algebra of maps from space to classifying space.

(shown now assuming $\Lambda = 0$, for simplicity):

topological flux quantum observables

$\mathbb{C}[\pi_0 C^\infty(\Sigma^2, G)] \simeq \mathbb{C}[\pi_0 \text{Maps}(\Sigma^2, G)]$
 $\simeq \mathbb{C}[\pi_1 \text{Maps}(\Sigma^2, BG)] \simeq H_0(\text{Maps}^*((\mathbb{R}^1 \times \Sigma^2)_{\cup\{\infty\}}, BG); \mathbb{C})$
 group algebra of fundamental group of maps to classifying space **homology Pontrjagin algebra of soliton moduli space**

soliton on X = topological field configuration that vanishes at the ends of X
 classified by *pointed* map
 $\Rightarrow X_{\cup\{\infty\}} \rightarrow BG$
 from one-point compactification

Example: $\mathbb{C}[\pi_0 \text{Maps}(\Sigma_g^2, U(1))] \simeq \mathbb{C}[H^1(\Sigma_g^2; \mathbb{Z})] \simeq \mathbb{C}[\mathbb{Z}^{2g}]$

Σ_g^2 orientable surface of genus= g

Effective flux of “fractional quantum Hall systems” (FQH).

But, at very low temperature, experiment suggests

instead of \mathbb{Z}^{2g} its 2nd **integer Heisenberg extension** $\widehat{\mathbb{Z}^{2g}}$ being the observables of an “**effective Chern-Simons field**”, where the center $\mathbb{Z} \hookrightarrow \widehat{\mathbb{Z}^{2g}}$ observes an **anyon braiding phase**.

Question: Is there classifying space \mathcal{A} for this effective CS field?


Answer: Yes! The 2-sphere $S^2 \simeq CP^1 \hookrightarrow CP^\infty \simeq BU(1)$

Theorem [61][84]: The cofiber presentation of the surface

$$S^1 \xrightarrow{\Pi_i[a_i, b_i]} \bigvee_g (S_a^1 \vee S_b^1) \longrightarrow \Sigma_g^2 \longrightarrow S^2$$

induces short exact sequence exhibiting the Heisenberg extension:

$$1 \rightarrow \underbrace{\pi_1 \text{Maps}(S^2, S^2)}_{\mathbb{Z}} \rightarrow \underbrace{\pi_1 \text{Maps}(\Sigma_g^2, S^2)}_{\widehat{\mathbb{Z}^{2g}}} \rightarrow \underbrace{\pi_1 \text{Maps}^*(\bigvee_{2g} S^1, S^2)}_{\mathbb{Z}^{2g}} \rightarrow 1$$



$$\widehat{\mathbb{Z}^{2g}} := \left\{ \begin{array}{l} (\vec{a}, \vec{b}, n) \in \mathbb{Z}^g \times \mathbb{Z}^g \times \mathbb{Z} \\ (\vec{a}, \vec{b}, n) \cdot (\vec{a}', \vec{b}', n') = \\ (\vec{a} + \vec{a}', \vec{b} + \vec{b}', n + n' + \vec{a} \cdot \vec{b}' - \vec{a}' \cdot \vec{b}) \end{array} \right\}$$

twice the unit central extension

Question: Can we identify the center \mathbb{Z} as arising from braiding?

Answer: Yes!


Theorem [132]: $\text{Maps}^*(S^2, S^2)$ is configurations of charged strings

such that $\Omega \text{Maps}^*(S^2, S^2)$ is framed links subject to cobordism, $\pi_1 \text{Maps}^*(S^2, S^2)$ generated from framed unknot with 1 braiding

$$\Omega \text{Maps}^*(S^2, S^2) \xrightarrow{[-]} \pi_3(S^2) \simeq \mathbb{Z}$$

$$L \longmapsto \#L \quad \text{is CS observable (“Wilson loop”)}$$

framed link linking + framing number



$$\# \left(\begin{array}{c} \nearrow \\ \searrow \end{array} \right) = +1, \quad \# \left(\begin{array}{c} \searrow \\ \nearrow \end{array} \right) = -1,$$


Ergo: Remarkably, topological quantum observables of effective flux in quantum Hall systems is algebro-topologically described by $\left\{ \begin{array}{l} \text{replacing the classifying space } BU(1) \simeq CP^\infty \\ \text{with its 2-skeleton } S^2 \simeq CP^1 \end{array} \right.$

Question 1: Is there a deeper rationale for such replacement? **Answer:** Yes [130][135]: *Hypothesis H.*

Question 2: Does this new model make novel predictions? **Answer:** Yes – *defect anyons* in FQH-systems:


With the classifying space identified for known situations, we find its implications for previously inaccessible cases:

Namely generalize now to n -punctured surfaces $\Sigma_{g,n}^2$, reflecting n defect points in the semiconductor where the magnetic field is expelled (type-I superconducting spots).



Proposition.
The observables are, in this generality:

$$\begin{aligned} \text{Obs}_0 &\simeq \mathbb{C} \left[\pi_1 \text{Maps}^* \left((\Sigma_{g,n}^2)_{\cup \{\infty\}}, S^2 \right) \right] \\ &\simeq \mathbb{C} \left[\pi_1 \text{Maps}^* \left(\Sigma_g^2 \vee \bigvee_{n-1} S^1, S^2 \right) \right] \\ &\simeq \mathbb{C} \left[\pi_1 \text{Maps}^* \left(\Sigma_g^2, S^2 \right) \times \mathbb{Z}^{n-1} \right] \\ &\underset{g=0}{\simeq} \mathbb{C} \left[\mathbb{Z}^n \right] \end{aligned}$$



Topology changes due to defects! (cf. [65, p 11]) Different to but not unlike the *genetz*-proposal [8].


subject to the diffeomorph. action by:

$$1 \rightarrow \text{Br}_n(\Sigma_g^2) \xleftrightarrow{\text{surface braid group}} \pi_0 \text{Homeos}_{\text{or}}^* \left((\Sigma_{g,n}^2)_{\cup \{\infty\}} \right) \xrightarrow{\text{mapping class group of punctured surface}} \text{MCG}(\Sigma_g^2) \rightarrow 1$$


mapping class group of plain surface

Therefore the equivariant quantum states (jargon: “generally covariant”) on $\Sigma_{0,n}^2$ are representations of the wreath product of solitonic and defect phases:

$$\mathbb{Z} \wr \text{Br}_n(\Sigma_0^2) = \overset{\text{solitonic anyons}}{\mathbb{Z}^n} \times \overset{\text{defect anyons}}{\text{Br}_n(\Sigma_0^2)} \rightarrow \mathbb{Z}^n \rtimes \text{Sym}_n$$



Such braid representations for defects have not previously been derived for FQH systems – but are just what is needed for the grand goal of *topological quantum gates*: programmable unitary transformations of quantum systems, insensitive to continuous deformations (hence to noise!)




Concretely, the worldvolume domains with n defects retract onto those with a single defect, whence so do the corresponding quantum observables and quantum states:

$$\mathcal{H}_{\Sigma_{0,n}^2} \simeq \mathbb{C}^n \simeq \mathbb{C} \oplus \left[\begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array} \right] \in \text{Sym}_n \text{Rep}$$

This yields what are known as *controlled qdit-rotation gates*,

the workhorse of quantum algorithms & the bottleneck for noise-protection, here topologically protected as cyclic defect braidings:

$$\text{e.g.: } \left[\begin{array}{|c|c|} \hline & \\ \hline \end{array} \right] \simeq \left\{ (213) \mapsto \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, (231) \mapsto \begin{bmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{bmatrix} \text{ where } \alpha = 4\pi/3 \right\}$$



Conclusion & Outlook:
With non-linear flux-quantization laws taken into account in physics, substantial Algebraic Topology reveals previously unrecognized phenomena potentially visible in experiment and relevant for quantum technology, potentially a much more fruitful commercial AlgTop-application than TDA!

Vistas. With this map from AlgTop to quantum effects established, there is opportunity to make AlgTop research inform quantum technology. Concretely:

Open problems in low-dim AlgTop. **Relevance for quantum system.**

describe the $\text{Br}_n(\Sigma_{g>0}^2)$ -action on $\pi_1 \text{Maps}^*((\Sigma_{g>0,n}^2) \cup \{\infty\}, S^2)$	novel exotic topological quantum gates
higher degree homology $H_{\bullet>0}(\Omega \text{Maps}^*((\Sigma_{g,n}^2) \cup \{\infty\}, S^2); \mathbb{C})$	“higher order” effects in topological phases
generalize to allow $b > 0$ boundary components $\Sigma_{g,n,b}^2$	small $b > 0$ is experi- mentally most accessible case


For the remainder – let’s shift gears:

We have seen that:

topological quantum states \mathcal{H}_Σ of solitonic field fluxes with classifying space \mathcal{A} on spacetime domain $\mathbb{R}^{1,1} \times \Sigma$	form representations of π_1 of the soliton moduli space $\text{Fields}_\Sigma := \text{Maps}^*(\Sigma, \mathcal{A}) // \text{Aut}^*(\Sigma)$
--	--


This is remarkable
because

such representations are equiv.
vector bundles \mathcal{H}_Σ on Fields_Σ
with *flat connections* ∇
a.k.a.: *local systems* on moduli



with the homotopy type of Fields_Σ
understood as an ∞ -groupoid,
(physics newspeak: generalized symmetry)
flat vector bundles are equivalently
functors $\vdash \mathcal{H}_\Sigma$ to the groupoid $\text{Mod}_{\mathbb{C}}$

which is the special case
of ∞ -*local systems* [128]:
chain complex-bundles
with flat ∞ -connection



which are equivalently Fields_Σ -
parameterized module spectra
for the E_∞ -ring HC
hence $HC[\Omega \text{Fields}_\Sigma]$ -modules

detecting higher structure
in the moduli space:

here HC denotes the
homotopy complex numbers:
the EM-ring spectrum of \mathbb{C}

and $HC[\Omega \text{Fields}_\Sigma]$ is the
homotopy Pontrjagin algebra
whose π_\bullet is Obs_\bullet .

These objects form the **tangent ∞ -topos** $T\text{Grpd}_\infty$ (over HC), which is [127][128]:

- (i) the arena of parameterized stable homotopy theory,
- (ii.) categorial semantics of a novel quantum programming language

Remarkably, this provides an AlgTop angle on
an ill-understood but central physics aspect:

*What exactly is quantum measurement
of anyonic topological order?*

Fact. [127]

- Given quantum states $\mathcal{H} \in \text{Mod}_{\mathbb{C}}^{\text{Fields}}$,
- a *quantum measurement basis* is
 - a choice of space W (of “possible worlds”)
 - a map $W \xrightarrow{i} \text{Fields}$ whose
 - base change is *ambidextrous*:

$$\begin{array}{ccc}
 \text{Mod}_{\mathbb{C}}^W & \xrightarrow{i_! \simeq i_*} & \text{Mod}_{\mathbb{C}}^{\text{Fields}} \\
 & \perp \top & \\
 & \xleftarrow{i^*} &
 \end{array}$$

- a $V \in \text{Mod}_{\mathbb{C}}^W$ which (co)induces $\mathcal{H} \simeq i_* V$,
- the *measurement \mathcal{E} collapse operation* is is
the counit $i^* \mathcal{H} \simeq i^* i_* V \xrightarrow{\text{ret}_V^i} V$.

Example.

Focusing on $\text{Fields} := * // \pi_0 \text{Homeo}(\Sigma_{g,n,b}^2)$,
then such measurement bases are given by
finite index subgroups of $\pi_0 \text{Homeo}(\Sigma_{g,n,b}^2)$.

There is a rich theory of these,
potentially of direct relevance for
realizing topological quantum computing...

More on these
quantum-information theoretic
aspects in [138]

A Background on Homotopy Theory

We collect some notions used in the main text to establish notation and give basic pointers to the literature.

Homotopy theory (cf. [141]). For $f_0, f_1 : X \rightarrow Y$ a pair of continuous maps between (topological) spaces a *homotopy* $\eta : f_0 \Rightarrow f_1$ is a continuous deformation between them: a continuous map $\eta : [0, 1] \times X \rightarrow Y$ such that

$$\begin{aligned} \eta(0, x) &= f_0(x), \\ \eta(1, x) &= f_1(x), \end{aligned} \quad \text{denoted} \quad X \begin{array}{c} \xrightarrow{f_0} \\ \Downarrow \eta \\ \xrightarrow{f_1} \end{array} Y.$$

For example, a square “homotopy- commutative diagram”

$$\begin{array}{ccc} \Sigma & \overset{b}{\dashrightarrow} & \mathcal{A} \\ \phi \downarrow & \overset{\eta}{\dashleftarrow} & \downarrow p \\ X & \overset{c}{\dashrightarrow} & \mathcal{B} \end{array} \quad \text{means that} \quad \begin{array}{l} \eta : [0, 1] \times \Sigma \rightarrow \mathcal{B} \\ \eta(0, s) = p(b(s)), \\ \eta(1, s) = c(\phi(s)). \end{array}$$

If one declares – and we do – to work in a “convenient” full sub-category of all topological spaces (such as that of *compactly generated* or of *Delta-generated* topological spaces, cf. [125, p 21, 131]) then the topological space $\text{Maps}(X, Y)$ of all continuous maps $X \rightarrow Y$ satisfies the adjointness relation

$$\{P \rightarrow \text{Maps}(X, Y)\} \simeq \{P \times X \rightarrow Y\}.$$

For $P \equiv [0, 1]$, this says that homotopies are equivalently paths in mapping spaces, and that homotopy-classes of maps are the mapping spaces’ path-connected components:

$$\pi_0 \text{Maps}(X, Y) \simeq \text{Maps}(X, Y)_{/\text{hmtpt}}.$$

Since homotopies are maps themselves, there are homotopies-between-homotopies and ever higher-homotopies.

Thereby, topological spaces constitute a model for **higher categorical symmetry** namely for higher groupoids. As such, they represent both cohomology as well as higher gauge fields in the topological sector. ⁷

Cohomology	cocycle	coboundary	higher coboundary	...
homotopy	$X \xrightarrow{f} \mathcal{B}$	$X \begin{array}{c} \xrightarrow{f} \\ \Downarrow \eta \\ \xrightarrow{f'} \end{array} \mathcal{B}$	$X \begin{array}{c} \xrightarrow{f} \\ \eta \left(\begin{array}{c} \Downarrow \\ \Downarrow \end{array} \right) \eta' \\ \xrightarrow{f'} \end{array} \mathcal{B}$...
Physics	field	gauge transf.	higher gauge transf.	...

In this vein, spaces are homotopy-equivalent $\mathcal{B} \simeq \mathcal{B}'$ if they are gauge equivalent namely if we have maps

$$\mathcal{B} \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} \mathcal{B}' \quad \text{with} \quad \begin{array}{l} g \circ f \Rightarrow \text{id}_{\mathcal{B}} \\ f \circ g \Rightarrow \text{id}_{\mathcal{B}'} \end{array}$$

For example $\mathbb{R}^n \simeq *$ in homotopy theory, reflecting the fact that there is no non-trivial topological sector for fields on \mathbb{R}^n .

For actually computing homotopy classes of maps — hence cohomology, hence gauge-equivalence classes of fields in the topological sector — tools from *model category theory* are indispensable, which largely say how to “absorb homotopies into spaces” (cf. [42, §1]).

For example, if $p : \mathcal{A} \rightarrow \mathcal{B}$ is a *Serre fibration*, such as a fiber bundle, and Σ is a *cell complex*, such as a manifold, then sections-up-to-homotopy of p pulled back to Σ are homotopy equivalent to plain sections:

$$\left\{ \begin{array}{ccc} \Sigma & \overset{b}{\dashrightarrow} & \mathcal{A} \\ \phi \downarrow & \overset{\eta}{\dashleftarrow} & \downarrow p \\ X & \xrightarrow{c} & \mathcal{B} \end{array} \right\}_{/\text{hmtpt}} \simeq \begin{array}{l} \Sigma \in \text{Cof} \\ p \in \text{Fib} \end{array} \left\{ \begin{array}{ccc} \Sigma & \overset{b}{\dashrightarrow} & \mathcal{A} \\ \phi \downarrow & \parallel & \downarrow p \\ X & \xrightarrow{c} & \mathcal{B} \end{array} \right\}_{/\text{hmtpt}}$$

Pointed homotopy theory (cf. [71, §3]). To reflect the condition that *solitonic fields are localized* in that they *vanish at infinity* we

⁷Beyond the topological sector, full higher gauge fields are still represented by maps $X \rightarrow \mathcal{B}$ etc., only that now \mathcal{B} is no longer just a topological space but a “smooth ∞ -stack”, cf. [35][42, pp 41].

- equip domain spaces X with a *point at infinity*, $\infty_X \in X$,
- equip classifying spaces \mathcal{B} with a *point representing zero*, $0_{\mathcal{B}} \in \mathcal{B}$,
- require maps $f : (X, \infty_X) \rightarrow (\mathcal{B}, 0_{\mathcal{B}})$ to respect these base points so that maps literally vanish at infinity

$$\begin{array}{ccc} X & \xrightarrow{c} & \mathcal{B} \\ \uparrow & & \uparrow \\ \{\infty_X\} & \longrightarrow & \{0_{\mathcal{B}}\}. \end{array}$$

For instance, to make fields on \mathbb{R}^n vanish at infinity, we adjoin its would-be “point at infinity” to it (jargon: “one-point compactification”) to obtain $\mathbb{R}_{\cup\{\infty\}}^n \simeq S^n$. On the other hand, if we want fields on some X without a vanishing condition, we may adjoin a *disjoint* point-at-infinity, then pointed maps $X_{\sqcup\{\infty\}} \rightarrow \mathcal{B}$ are ordinary $X \rightarrow \mathcal{B}$. For example,

$$\begin{array}{ccc} \text{based loop space} & \text{free loop space} & \text{maps out of contractible} \\ \text{Maps}^*(\mathbb{R}_{\cup\{\infty\}}^1, \mathcal{B}) = \Omega\mathcal{B}, & \text{Maps}^*(S_{\sqcup\{\infty\}}^1, X) =: \mathcal{L}\mathcal{B}, & \text{Maps}^*(\mathbb{R}_{\sqcup\{\infty\}}^1, \mathcal{B}) = \mathcal{B}. \end{array}$$

Given a pair of pointed spaces $(X, \infty_X), (Y, \infty_Y)$, in their product space $X \times Y$ any point should be regarded as being at infinity which is so with respect to either factor space; this yields the *smash product*:

$$X \wedge Y := \frac{X \times Y}{\{\infty_X\} \times Y \cup X \times \{\infty_Y\}}$$

to which the sub-space $\text{Maps}^*(-, -)$ of pointed maps is again adjoint:

$$\{P \xrightarrow{\text{pntd}} \text{Maps}^*(X, Y)\} \simeq \{P \wedge X \xrightarrow{\text{pntd}} Y\}.$$

For example,

$$S^n \wedge S^m \simeq \mathbb{R}_{\cup\{\infty\}}^n \wedge \mathbb{R}_{\cup\{\infty\}}^m \simeq (\mathbb{R}^n \times \mathbb{R}^m)_{\cup\{\infty\}} \simeq S^{n+m},$$

so that, for instance:

$$\text{Maps}^*(X \wedge S^1, \mathcal{B}) \simeq \text{Maps}^*(S^1, \text{Maps}^*(X, \mathcal{B})) =: \Omega \text{Maps}^*(X, \mathcal{B}).$$

The differential character map $\text{ch}_{\mathcal{A}}$, at the heart of flux-quantization in the generality of flux densities with non-linear Bianchi identities:

- takes maps into a classifying space \mathcal{A} (classifying **charges**),
- to maps into the moduli ∞ -stack of closed $\mathcal{L}\mathcal{A}$ -valued differential forms (classifying corresponding **flux densities**),
- thereby allowing **gauge potentials** to relate local flux densities to global charges.

$$\begin{array}{ccc} & X & \\ \text{charges} \swarrow & & \searrow \text{fluxes} \\ \mathcal{A} & & \Omega_{\text{dR}}^1(-; \mathcal{L}\mathcal{A})_{\text{clsd}} \\ \text{character} \searrow & \xrightarrow{\text{gauge potentials}} & \swarrow \eta^\natural \\ & \int \Omega_{\text{dR}}^1(-; \mathcal{L}\mathcal{A})_{\text{clsd}} & \end{array}$$

At a high level, this $\text{ch}_{\mathcal{A}}$ is readily described: It is the smooth differential-form model for the \mathbb{R} -**rationalization** of \mathcal{A} , followed by derived extension of scalars $\mathbb{Q} \rightarrow \mathbb{R}$ — as indicated in the following paragraphs.

However, under the hood, this construction makes use of a fair bit of model category-theoretic rational-homotopy theory which we do not have space nor inclination to review here (all details in [42]), whence the following should be ignored by readers without serious background in (rational) homotopy theory — or else taken as motivation to learn it! (Start at [42, §1].) Here is how it goes:

Fundamental theorem of homotopy theory. Regarding (classifying) spaces up to (weak) homotopy equivalence means equivalently to regard them as their ∞ -*groupoids* (Kan simplicial sets) $\text{Sing}(-)$ of points, paths, 2-paths, etc., in that there is a Quillen equivalence [42, Ex. 1.13]

$$\begin{array}{ccc} \text{TopSp}_{\text{Qu}} & \xleftarrow{\simeq_{\text{Qu}}} & \Delta\text{Set}_{\text{Qu}} \\ & \xrightarrow{\text{Sing}} & \end{array}$$

Fundamental theorem of dg-algebraic rational homotopy theory. Sending simplicial sets to their dgc-algebras of simplex-wise \mathbb{Q} -polynomial differential forms (“piecewise linear”, PL) is the left adjoint in a Quillen adjunction [42, Prop. 5.5]

$$\begin{array}{ccc} (\mathrm{dgcAlgs}^{\geq 0})_{\mathrm{proj}}^{\mathrm{op}} & \xleftarrow{\Omega_{\mathbb{P}\mathbb{Q}\mathrm{LdR}}^{\bullet}} & \Delta\mathrm{Sets}_{\mathbb{Q}\mathrm{u}} \\ & \xrightarrow[\mathrm{Hom}(-, \Omega_{\mathbb{P}\mathbb{Q}\mathrm{LdR}}(\Delta^{\bullet}))]{\perp_{\mathbb{Q}\mathrm{u}}} & \end{array}$$

whose derived adjunction-unit models rationalization of (connected, nilpotent, \mathbb{Q} -finite) homotopy types \mathcal{A} [42, Prop. 5.6].

$$\mathcal{A} \xrightarrow{\eta_{\mathcal{A}}^{\mathbb{Q}}} L^{\mathbb{Q}}\mathcal{A}$$

For \mathbb{R} -rational homotopy. The analogous Quillen adjunction with \mathbb{R} -polynomial forms

$$\begin{array}{ccc} (\mathrm{dgcAlgs}^{\geq 0})_{\mathrm{proj}}^{\mathrm{op}} & \xleftarrow{\Omega_{\mathbb{P}\mathbb{R}\mathrm{L}}^{\bullet}} & \Delta\mathrm{Sets}_{\mathbb{Q}\mathrm{u}} \\ & \xrightarrow[\mathrm{Hom}(-, \Omega_{\mathbb{P}\mathbb{R}\mathrm{LdR}}(\Delta^{\bullet}))]{\perp_{\mathbb{Q}\mathrm{u}}} & \end{array}$$

models rationalization followed by derived extension of scalars from \mathbb{Q} to \mathbb{R} (no longer a localization but still denoted like one) [42, Prop. 5.8].

$$\mathcal{A} \xrightarrow{\eta_{\mathcal{A}}^{\mathbb{Q}}} L^{\mathbb{Q}}\mathcal{A} \longrightarrow L^{\mathbb{R}}\mathcal{A}$$

Now with \mathbb{R} -coefficients, we may equivalently use simplex-wise *smooth* differential forms (*piecewise smooth*, PS)

$$\begin{array}{ccc} (\mathrm{dgcAlgs}^{\geq 0})_{\mathrm{proj}}^{\mathrm{op}} & \xleftarrow{\Omega_{\mathbb{P}\mathrm{SdR}}^{\bullet}} & \Delta\mathrm{Sets}_{\mathbb{Q}\mathrm{u}} \\ & \xrightarrow[\mathrm{Hom}(-, \Omega_{\mathbb{P}\mathrm{SdR}}(\Delta^{\bullet}))]{\perp_{\mathbb{Q}\mathrm{u}}} & \end{array}$$

In fact, we may equivalently use smooth differential forms on simplices times any \mathbb{R}^n [42, Prop. 5.10].

$$\begin{array}{ccc} (\mathrm{dgcAlgs}^{\geq 0})_{\mathrm{proj}}^{\mathrm{op}} & \xleftarrow{\Omega_{\mathbb{P}\mathrm{SdR}}^{\bullet}} & \Delta\mathrm{Sets}_{\mathbb{Q}\mathrm{u}} \\ & \xrightarrow[\mathrm{Hom}(-, \Omega_{\mathbb{P}\mathrm{SdR}}(\mathbb{R}^n \times \Delta^{\bullet}))]{\perp_{\mathbb{Q}\mathrm{u}}} & \end{array}$$

Taking values in deformations of flux densities. Via the minimal Sullivan model $\mathrm{CE}(\mathcal{L}\mathcal{A})$ of \mathcal{A} , this derived adjunction takes values in closed smooth $\mathcal{L}\mathcal{A}$ -valued differential forms [42, (9.9)]

$$\Omega_{\mathrm{dR}}^1(\mathbb{R}^n \times \Delta^{\bullet}, \mathcal{L}\mathcal{A})_{\mathrm{clsd}} := \mathrm{Hom}(\mathrm{CE}(\mathcal{L}\mathcal{A}), \Omega_{\mathrm{dR}}(\mathbb{R}^n \times \Delta^{\bullet}))$$

which is the value on \mathbb{R}^n of the homotopy-constant ∞ -stack that is the *shape* $\int(-)$ of the sheaf of closed forms [125, Prop. 3.3.48]

$$\int \Omega_{\mathrm{dR}}^1(-; \mathcal{L}\mathcal{A})_{\mathrm{clsd}} \in \mathrm{Sh}_{\infty}(\mathrm{CartSp}).$$

In total, regarding also $\mathcal{A} \in \mathrm{Sh}_{\infty}(\ast) \xrightarrow{\mathrm{Disc}} \mathrm{Sh}_{\infty}(\mathrm{CartSp})$, this establishes the *differential character* map as promised [42, Def. 9.2]


$$\mathcal{A} \xrightarrow{\mathrm{ch}_{\mathcal{A}}} \int \Omega_{\mathrm{dR}}^1(-; \mathcal{L}\mathcal{A})_{\mathrm{clsd}}$$

B Background on TED Cohomotopy

Gauge potentials in twistorial Cohomotopy — and the Green-Schwarz mechanism. Consider the Whitehead L_{∞} -algebra of the twistor fibration $\mathbb{C}P^3 \xrightarrow{t_{\mathbb{H}}} \mathbb{H}P^1 \simeq S^4$,

$$\mathrm{CE}(t_{S^4} \mathbb{C}P^3) = \mathbb{R}_{\mathrm{d}} \left[\begin{array}{c} f_2 \\ h_3 \\ g_4 \\ g_7 \end{array} \right] / \left(\begin{array}{l} \mathrm{d} f_2 = 0 \\ \mathrm{d} h_3 = g_4 + f_2 f_2 \\ \mathrm{d} g_4 = 0 \\ \mathrm{d} g_7 = \frac{1}{2} g_4 g_4 \end{array} \right),$$

and bigons parameterized like this:



Theorem ([52, p 23][53, §4.1]). Given a manifold U_i (generically: a coordinate chart):

(i) Closed $\mathbb{I}_{S^4}\mathbb{C}P^3$ -valued differential forms are in natural bijection with **flux densities** of this form:

$$\left\{ \begin{array}{c} U_i \\ \downarrow \\ (F_2, H_3, G_4, G_7) \\ \downarrow \\ \Omega_{\text{dR}}^1(-; \mathbb{I}_{S^4}\mathbb{C}P^3)_{\text{clsd}} \end{array} \right\} \begin{array}{c} \xrightarrow[p_0]{i_0 \circ p_0 = \text{id}} \\ \xleftarrow[i_0]{p_0 \circ i_0 = \text{id}} \end{array} \left\{ \begin{array}{l} F_2 \in \Omega_{\text{dR}}^2(U_i) \\ H_3 \in \Omega_{\text{dR}}^3(U_i) \\ G_4 \in \Omega_{\text{dR}}^4(U_i) \\ G_7 \in \Omega_{\text{dR}}^7(U_i) \end{array} \middle| \begin{array}{l} dF_2 = 0 \\ dH_3 = G_4 + F_2 F_2 \\ dG_4 = 0 \\ dG_7 = \frac{1}{2} G_4 G_4 \end{array} \right\}$$

(ii) Given one of these, its set of coboundaries (null-concordances) naturally retracts onto the set of **gauge potentials** of this form:

$$\left\{ \begin{array}{c} U_i \xrightarrow{\quad} * \\ \downarrow \\ (F_2, H_3, G_4, G_7) \xrightarrow{\text{red dashed}} (\widehat{F}_2, \widehat{H}_3, \widehat{G}_4, \widehat{G}_7) \\ \downarrow \eta^f \\ \Omega_{\text{dR}}^1(-; \mathbb{I}_{S^4}\mathbb{C}P^3)_{\text{clsd}} \xrightarrow{\eta^f} \int \Omega_{\text{dR}}^1(-; \mathbb{I}_{S^4}\mathbb{C}P^3)_{\text{clsd}} \end{array} \right\} \begin{array}{c} \xrightarrow[p_1]{p_1 \circ i_1 = \text{id}} \\ \xleftarrow[i_1]{} \end{array} \left\{ \begin{array}{l} A_1 \in \Omega_{\text{dR}}^1(U_i) \\ B_2 \in \Omega_{\text{dR}}^2(U_i) \\ C_3 \in \Omega_{\text{dR}}^3(U_i) \\ C_6 \in \Omega_{\text{dR}}^6(U_i) \end{array} \middle| \begin{array}{l} dA_1 = F_2 \\ dB_2 = H_3 - C_3 - A_1 F_2 \\ dC_3 = G_4 \\ dC_6 = G_7 - \frac{1}{2} C_3 G_4 \end{array} \right\}$$

$$\left(\begin{array}{l} \widehat{F}_2 := t F_2 + dt A_1 \\ \widehat{H}_3 := t H_3 + dt B_2 + (t^2 - t) A_1 F_2 \\ \widehat{G}_4 := t G_4 + dt C_3 \\ \widehat{G}_7 := t^2 G_7 + 2tdt C_6 \end{array} \right) \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \left(\begin{array}{l} A_1 := \int_{[0,1]} \widehat{F}_2 \\ B_2 := \int_{[0,1]} \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right) \\ C_3 := \int_{[0,1]} \widehat{G}_4 \\ C_6 := \int_{[0,1]} \left(\widehat{G}_7 - \frac{1}{2} \left(\int_{[0,-]} \widehat{G}_4 \right) \widehat{G}_4 \right) \end{array} \right)$$

(iii) Given a pair of these, the set of higher coboundaries (2nd-order concordances) between them naturally retracts onto the set of **gauge transformations** of this form:

$$\left\{ \begin{array}{c} (\widehat{F}_2, \widehat{H}_3, \widehat{G}_4, \widehat{G}_7) \\ \downarrow \text{red dashed} \\ (\widehat{F}_2, \widehat{H}_3, \widehat{G}_4, \widehat{G}_7) \\ \downarrow \\ (F_2, H_3, G_4, G_7) \\ \downarrow \text{red dashed} \\ (\widehat{F}'_2, \widehat{H}'_3, \widehat{G}'_4, \widehat{G}'_7) \end{array} \right\} \begin{array}{c} \xrightarrow[p_2]{p_2 \circ i_2 = \text{id}} \\ \xleftarrow[i_2]{} \end{array} \left\{ \begin{array}{l} \alpha_0 \in \Omega_{\text{dR}}^0(U_i) \\ \beta_1 \in \Omega_{\text{dR}}^1(U_i) \\ \gamma_2 \in \Omega_{\text{dR}}^2(U_i) \\ \gamma_5 \in \Omega_{\text{dR}}^5(U_i) \end{array} \middle| \begin{array}{l} d\alpha_0 = A'_1 - A_1 \\ d\beta_1 = B'_2 - B_2 + \gamma_2 + \alpha_0 F_2 \\ d\gamma_2 = C'_3 - C_3 \\ d\gamma_5 = C'_6 - C_6 - \frac{1}{2} C'_3 C_3 \end{array} \right\}$$

$$\left(\begin{array}{l} \widehat{\widehat{F}}_2 := t F_2 + dt A_1 + s dt (A'_1 - A_1) - ds dt \alpha_0 \\ \widehat{\widehat{H}}_3 := t H_3 + dt B_2 + s dt (B'_2 - B_2) - ds dt \beta_1 \\ \quad + (t^2 - t) A_1 F_2 + (t^2 - t) s (A'_1 - A_1) F_2 \\ \quad + (t^2 - t) ds \alpha_0 F_2 \\ \widehat{\widehat{G}}_4 := t G_4 + dt C_3 + s dt (C'_3 - C_3) - ds dt \gamma_2 \\ \widehat{\widehat{G}}_7 := t^2 G_7 + 2tdt C_6 + 2stdt (C'_6 - C_6) \\ \quad - 2ds t dt (\gamma_5 + \frac{1}{2} \gamma_2 C_3) \end{array} \right) \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \left(\begin{array}{l} \alpha_0 := \int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{\widehat{F}}_2 \\ \beta_1 := \int_{s \in [0,1]} \int_{t \in [0,1]} \left(\widehat{\widehat{H}}_3 - \left(\int_{t' \in [0,-]} \widehat{\widehat{F}}_2 \right) \widehat{\widehat{F}}_2 \right) \\ \gamma_2 := \int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{\widehat{G}}_4 \\ \gamma_5 := \int_{s \in [0,1]} \int_{t \in [0,1]} \left(\widehat{\widehat{G}}_7 - \frac{1}{2} \left(\int_{t' \in [0,-]} \widehat{\widehat{G}}_4 \right) \widehat{\widehat{G}}_4 \right) \\ \quad - \frac{1}{2} \gamma_2 C_3 \end{array} \right)$$

Notice the expression for flux density subject to an (abelian) Green-Schwarz mechanism:

$$H_3 = d B_2 + A_1 F_2 + C_3 .$$

Proof. With the blue terms discarded, this is the statement of [52, p 23][53, §4.1]. We compile the full argument. To see that p_1 is well-defined:

- for C_3, C_6 this is [52, (70)],
- for A_1 it works just as for C_3 ,
- for B_2 we compute, in generalization of [53, below (138)], like this:

$$\begin{aligned}
d B_2 &\equiv d \int_{[0,1]} \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right) \\
&= \underbrace{\iota_1^* \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right)}_{H_3 - A_1 F_2} - \underbrace{\iota_0^* \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right)}_{=0} - \int_{[0,1]} \underbrace{d \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right)}_{\widehat{G}_4} \\
&= H_3 - A_1 F_2 - C_3.
\end{aligned}$$

To see that i_1 is well-defined:

- for $\widehat{G}_4, \widehat{G}_7$ this is [52, (72)],
- for \widehat{F}_2 it works just as for \widehat{G}_4 ,
- for \widehat{H}_3 we compute, in generalization of [53, further below (138)], as follows:

$$\left. \begin{aligned}
&d(tH_3 + dt B_2 + (t^2 - t)A_1 F_2) \\
&= dt H_3 + tG_4 + tF_2 F_2 \\
&\quad - dt H_3 + dt C_3 + dt A_1 F_2 \\
&\quad + d((t^2 - t)A_1 F_2)
\end{aligned} \right\} \text{hence indeed: } d \widehat{H}_3 = \underbrace{tG_4 + dt C_3}_{\widehat{G}_4} + \underbrace{(tF_2 + dt A_1)}_{\widehat{F}_2} \underbrace{(tF_2 + dt A_1)}_{\widehat{F}_2} \overset{=d(t^2 A_1 F_2)}{=}$$

Moreover, it is immediate from inspection that $\iota_1^* \widehat{H}_3 = H_3$ and $\iota_0^* \widehat{H}_3 = 0$.

To see that $p_1 \circ i_1 = \text{id}$:

- for C_3, C_6 this is [52, below (72)],
- for A_1 this works just as for C_3 ,
- for B_2 we immediately compute:

$$\int_{[0,1]} \left(\widehat{H}_3 - \left(\int_{[0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right) = \underbrace{\int_{[0,1]} dt B_2}_{B_2} - \int_{[0,1]} \underbrace{tA_1 dt A_1}_{=0} = B_2.$$

To see that p_2 is well-defined:

- for $\widehat{G}_4, \widehat{G}_7$ this is [52, (74-5)],
- for \widehat{F}_2 this works just as for \widehat{F}_2 ,
- for \widehat{H}_3 we compute, in generalization of [53, below (140)], as follows:

$$\begin{aligned}
d\beta_1 &\equiv d \int_{s \in [0,1]} \int_{t \in [0,1]} \left(\widehat{H}_3 - \left(\int_{t' \in [0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right) \\
&= \iota_{s=1}^* \int_{t \in [0,1]} \left(\widehat{H}_3 - \dots \right) - \iota_{s=0}^* \int_{t \in [0,1]} \left(\widehat{H}_3 - \dots \right) - \int_{s \in [0,1]} d \int_{t \in [0,1]} \left(\widehat{H}_3 - \dots \right) \\
&= \int_{t \in [0,1]} \iota_{s=1}^* \left(\widehat{H}_3 - \dots \right) - \int_{t \in [0,1]} \iota_{s=0}^* \left(\widehat{H}_3 - \dots \right) - \int_{s \in [0,1]} \iota_{t=1}^* \left(\widehat{H}_3 - \dots \right) + \int_{s \in [0,1]} \int_{t \in [0,1]} d \left(\widehat{H}_3 - \dots \right) \\
&= \int_{t \in [0,1]} \left(\widehat{H}'_3 - \dots \right) - \int_{t \in [0,1]} \left(\widehat{H}_3 - \dots \right) + \left(\int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{F}_2 \right) F_2 + \int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{G}_4 \\
&= B'_2 - B_2 + \alpha_0 F_2 + \gamma_2.
\end{aligned}$$

To see that i_2 is well-defined:

- for γ_2, γ_5 this is [52, (76)],
- for α_0 this works just as for γ_2 ,

– for β_1 we compute as follows:

$$\begin{aligned}
d(t H_3 + dt B_2 + s dt (B'_2 - B_2) - ds dt \beta_1) &= \overbrace{t G_4 + dt C_3 + s dt (C'_3 - C_3) - ds dt \gamma_2}^{\widehat{G}_4} \\
&\quad + t F_2 F_2 + dt A_1 F_2 + s dt (A'_1 - A_1) F_2 - ds dt \alpha_0 F_2 \\
d \left(\begin{array}{l} (t^2 - t) A_1 F_2 + (t^2 - t) s (A'_1 - A_1) F_2 \\ + (t^2 - t) ds \alpha_0 F_2 \end{array} \right) &= \overbrace{t^2 F_2 F_2 + 2t dt A_1 F_2 + 2t dt s (A'_1 - A_1) F_2 + 2t dt ds \alpha_0 F_2}^{\widehat{F}_2 \widehat{F}_2} \\
&\quad - t F_2 F_2 - dt A_1 F_2 - dt s (A'_1 - A_1) F_2 - dt ds \alpha_0 F_2 \\
\hline
d \widehat{H}_3 &= \widehat{G}_4 + \widehat{F}_2 \widehat{F}_2.
\end{aligned}$$

Moreover, it is immediate from inspection that $\iota_{s=0}^* \widehat{H}_3 = \widehat{H}_3$, $\iota_{s=1}^* \widehat{H}_3 = \widehat{H}'_3$ and $\iota_{t=0}^* = 0$, $\iota_{t=1}^* = H_3$.

To see that $p_2 \circ i_2 = \text{id}$, we directly compute, first

$$\begin{aligned}
\int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{G}_4 &= \int_{s \in [0,1]} \int_{t \in [0,1]} (-ds dt \gamma_2) = \gamma_2 \\
\int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{F}_2 &= \int_{s \in [0,1]} \int_{t \in [0,1]} (-ds dt \alpha_0) = \alpha_0
\end{aligned}$$

then

$$\begin{aligned}
&\int_{s \in [0,1]} \int_{t \in [0,1]} \left(\widehat{G}_7 - \frac{1}{2} \left(\int_{t' \in [0,t]} \widehat{G}_4 \right) \widehat{G}_4 \right) - \frac{1}{2} \gamma_2 C_3 \\
&= \int_{s \in [0,1]} \int_{t \in [0,1]} \widehat{G}_7 - \frac{1}{2} \int_{s \in [0,1]} \int_{t \in [0,1]} (t C_3 + s t (C'_3 - C_2) + t ds \gamma_2) (t G_4 + dt C_3 + s dt (C'_3 - C_3) - ds dt \gamma_2) \\
&\quad - \frac{1}{2} \gamma_2 C_3 \\
&= (\gamma_5 + \frac{1}{2} \gamma_2 C_3) \underbrace{- \frac{1}{2} C_3 \gamma_2 - \frac{1}{4} (C'_3 - C_3) \gamma_2 + \frac{1}{2} \gamma_2 C_3 + \frac{1}{4} \gamma_2 (C'_3 - C_3)}_0 - \frac{1}{2} \gamma_2 C_3 \\
&= \gamma_5
\end{aligned}$$

and analogously

$$\begin{aligned}
&\int_{s \in [0,1]} \int_{t \in [0,1]} \left(\int_{t' \in [0,-]} \widehat{F}_2 \right) \widehat{F}_2 \\
&= \int_{s \in [0,1]} \int_{t \in [0,1]} (t A_1 + s t (A'_1 - A_1) + t ds \alpha_0) (t F_2 + dt A_1 + s dt (A'_1 - A_1) - ds dt \alpha_0) \\
&= \frac{1}{2} A_1 \alpha_0 + \frac{1}{4} (A'_1 - A_1) \alpha_0 - \frac{1}{2} \alpha_0 A_1 - \frac{1}{4} \alpha_0 (A'_1 - A_1) \\
&= 0
\end{aligned}$$

so that also

$$\int_{s \in [0,1]} \int_{t \in [0,1]} \left(\widehat{H}_3 - \left(\int_{t' \in [0,-]} \widehat{F}_2 \right) \widehat{F}_2 \right) = \int_{s \in [0,1]} \int_{t \in [0,1]} (-ds dt \beta_1) = \beta_1. \quad \square$$

Cocycles in differential 2-Cohomotopy and the abelian Chern-Simons invariant on the 3-Sphere.


Notice that the Bianchi identities encoded by 2-Cohomotopy are the characteristic property of the abelian Chern-Simons term:

$$\text{CE}(\mathfrak{LS}^2) \simeq \mathbb{R}_d \left[\begin{array}{l} f_2 \\ h_3 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d h_3 = f_2 f_2 \end{array} \right) \Rightarrow \Omega_{\text{dR}}^1(X; \mathfrak{LS}^2)_{\text{clsd}} \simeq \left\{ \begin{array}{l} F_2 \in \Omega_{\text{dR}}^2(X) \\ H_3 \in \Omega_{\text{dR}}^3(X) \end{array} \middle| \begin{array}{l} d F_2 = 0 \\ d H_3 = F_2 F_2 \end{array} \right\}.$$

We may bring this out more concretely:

Gauge-field configurations on \mathbb{R}^3 flux-quantized in 2-Cohomotopy and vanishing in a neighborhood of infinity are cocycles in differential 2-Cohomotopy on $\mathbb{R}_{\cup\{\infty\}}^3$, hence dashed homotopies as shown on the right [130,


§3.3].



Theorem. For each $[n] \in \pi^2(\mathbb{R}_{\cup\{\infty\}}^3) \simeq \mathbb{Z}$ this exists with $H_3 = 0$ and $[n] = \int_{\mathbb{R}^3} A_1 F_2$ the Chern-Simons invariant.

To see this, first consider:

Lemma. On a smooth manifold Σ , every cocycle α in rational 3-Cohomotopy is represented by a globally defined differential form H_3 ,



Proof of the Lemma. Since $\mathbb{I}S^3 \simeq \mathbb{I}B^3\mathbb{Q}$ this is just the degree=3 case of the statement that cocycles in de Rham hyper-cohomology have global representatives on smooth manifolds (using partitions of unity). \square

Proof of the Theorem. Stereographic projection provides a homeomorphism $\mathbb{R}_{\cup\{\infty\}}^3 \xrightarrow{\sim} S^3$ which is smooth away from the point at infinity, which we may slightly deform to a smooth degree=1 map that is constant on a neighborhood of infinity. Since $\pi^2(S^3) \simeq \pi_2(S^3) \simeq \mathbb{Z}$ we may find a smooth map $n : S^3 \rightarrow S^2$, with compact support away from the base point, so that $\mathbb{R}_{\cup\{\infty\}}^3 \rightarrow S^3 \xrightarrow{n} S^2$ represents the charge $[n]$.

Now the 2-cohomotopical character map for charges on S^3 , shown in black, factors as shown in blue (by naturality of rationalization), which furthermore factors as shown in orange (by the above Lemma).

$$\begin{array}{ccccccc}
 \mathbb{R}_{\cup\{\infty\}}^3 & \rightarrow & S^3 & \xrightarrow{\eta^f} & \int S^3 & \xrightarrow{n} & \int S^2 \\
 & & \downarrow n \cdot \text{dvol}_{S^3} & \swarrow & \downarrow \text{ch}_{S^3} & \swarrow & \downarrow \text{ch}_{S^2} \\
 \Omega_{\text{dR}}^1(-; \mathbb{I}S^3)_{\text{clsd}} & \xrightarrow{\eta^f} & \int \Omega_{\text{dR}}^1(-; \mathbb{I}S^3)_{\text{clsd}} & \xrightarrow{(tn)_*} & \int \Omega_{\text{dR}}^1(-; \mathbb{I}S^2)_{\text{clsd}} & &
 \end{array} \tag{11}$$

Hence, to get a differential cocycle as desired, it is sufficient to exhibit gauge potentials (A_1, B_2) encoding a concordance filling the diagram on the right

$$\begin{array}{ccccccc}
 \mathbb{R}_{\cup\{\infty\}}^3 & \rightarrow & S^3 & \xrightarrow{n} & S^2 & \xrightarrow{\eta^f} & \int S^2 \\
 & & \downarrow n \cdot \text{dvol}_{S^3} & \swarrow & \downarrow & \swarrow (11) & \downarrow \text{ch}_2 \\
 & & (F_2, H_3=0) & \xrightarrow{(A_1, B_2)} & \Omega_{\text{dR}}^1(-; \mathbb{I}S^3)_{\text{clsd}} & \xrightarrow{(tn)_* \circ \eta^f} & \int \Omega_{\text{dR}}^1(-; \mathbb{I}S^2)_{\text{clsd}} \\
 & & \downarrow & & \downarrow & & \downarrow \\
 \Omega_{\text{dR}}^1(-; \mathbb{I}S^2)_{\text{clsd}} & \xrightarrow{\eta^f} & \int \Omega_{\text{dR}}^1(-; \mathbb{I}S^2)_{\text{clsd}} & & & &
 \end{array}$$

However, since $H_{\text{dR}}^2(S^3) = 0$, and by the *Whitehead integral formula* (cf. [56, p 134][12, p 228][38, p 19]) there exists:

$$\begin{cases} A_1 \in \Omega_{\text{dR}}^1(S^3) \\ B_2 \in \Omega_{\text{dR}}^2(S^3) \end{cases} \text{ s.t. } \begin{cases} dA_1 = F_2 := n^* \text{dvol}_{S^2} \\ dB_2 = n \cdot \text{dvol}_{S^3} - A_1 F_2 \end{cases} \tag{12}$$

From this we get the the desired concordance:

$$(0, n \cdot \text{dvol}_{S^3}) \Rightarrow (F_2, 0) : \begin{cases} \widehat{F}_2 := t F_2 + dt A_1, \\ \widehat{H}_3 := (t-1)n \text{dvol}_{S^3} + dt B_2 + (t^2-t)A_1 F_2, \end{cases} \begin{cases} (\widehat{F}_2, \widehat{H}_3)|_{t=0} = (0, n \cdot \text{dvol}_{S^3}) \\ (\widehat{F}_2, \widehat{H}_3)|_{t=1} = (F_2, 0) \\ d\widehat{F}_2 = 0, d\widehat{H}_3 = \widehat{F}_2 \widehat{F}_2. \end{cases} \quad \square$$

Cartesian M5-Probes charged in Cohomotopy. The equations of motion for a(n orbifolded) cartesian M5-probe demand that the flux $H_3 = \text{const}$ [53, Ex. 3.14], and thus its solitonic vanishing-at-infinity implies $H_3 = 0$. The above theorem says that such solutions still support non-vanishing cohomotopical charge, in fact that the vanishing of H_3 forces the charge to be carried by the Chern-Simons invariant of the auxiliary gauge field \mathbf{A}_1 that is brought in by the cohomotopical flux quantization.

References

- [1] Aharony, O., Gubser, S., Maldacena, J., Ooguri, H., Oz, Y., *Large N Field Theories, String Theory and Gravity*, Phys. Rept. **323** (2000), 183-386, [doi:10.1016/S0370-1573(99)00083-6], [arXiv:hep-th/9905111].
- [2] An, Y.-S., Li L., Yang, F.-G., Yang, R.-Q., *Interior Structure and Complexity Growth Rate of Holographic Superconductor from M-Theory*, J. High Energ. Phys. **2022** 133 (2022), [doi:10.1007/JHEP08(2022)133], [arXiv:2205.02442].
- [3] Arovas, D. P., Schrieffer, R., Wilczek, F., *Fractional Statistics and the Quantum Hall Effect*, Phys. Rev. Lett. **53** (1984) 722, [doi:10.1103/PhysRevLett.53.722].
- [4] Arovas, D. P., Schrieffer, R., Wilczek, F., Zee, A., *Statistical mechanics of anyons*, Nucl. Phys. B **251** (1985), 117-126, [doi:10.1016/0550-3213(85)90252-4].
- [5] Bae, J.-B., Lee, S., *Emergent Supersymmetry on the Edges*, SciPost Phys. **11** 091 (2021), [arXiv:2105.02148], [doi:10.21468/SciPostPhys.11.5.091].
- [6] Bakulev, A. P., Shirkov, D., *Inevitability and Importance of Non-Perturbative Elements in Quantum Field Theory*, Proceedings of the 6th Mathematical Physics Meeting, Belgrade (2010), 27–54, [arXiv:1102.2380], [ISBN:978-86-82441-30-4].
- [7] Balachandran, A. P., Srivastava, A. M., *Chern-Simons Dynamics and the Quantum Hall Effect*, [arXiv:hep-th/9111006].
- [8] Barkeshli, M., Jian, C.-M., X.-L. Qi, *Twist defects and projective non-Abelian braiding statistics*, Phys. Rev. B **87** (2013) 045130 [doi:10.1103/PhysRevB.87.045130], [arXiv:1208.4834].
- [9] Beardsley, J., Nakamura, S., *Projective Geometries and Simple Pointed Matroids as \mathbb{F}_1 -modules*, [arXiv:2404.04730].
- [10] Beterov, I. I., *Progress and Prospects in the Field of Quantum Computing*, Optoelectron. Instrument. Proc. **60** (2024), 74–83, [doi:10.3103/S8756699024700043].
- [11] Bhattacharjee, M., Macpherson, D., Möller, R. G., Neumann, P. M., *Notes on Infinite Permutation Groups*, Lecture Notes in Mathematics **1698**, Springer (2006), 67-76, [doi:10.1007/BFb0092558].
- [12] Bott, R., Tu, L., *Differential Forms in Algebraic Topology*, Graduate Texts in Mathematics **82**, Springer (1982), [doi:10.1007/978-1-4757-3951-0].
- [13] Bourjaily, J. L., Espahbodi S., *Geometrically Engineerable Chiral Matter in M-Theory*, [arXiv:0804.1132].
- [14] Burton, S., Sati, H., Schreiber, U., *Lift of fractional D-brane charge to equivariant Cohomotopy theory*, J. Geometry and Physics **161** (2021) 104034 [doi:10.1016/j.geomphys.2020.104034], [arXiv:1909.12277].
- [15] Chakraborty, T., Pietiläinen, P., *The Quantum Hall Effects – Integral and Fractional*, Springer Series in Solid State Sciences (1995), [doi:10.1007/978-3-642-79319-6].
- [16] Chan, P. O., Teo, J. C. Y., Ryu, S., *Topological Phases on Non-orientable Surfaces: Twisting by Parity Symmetry*, New J. Phys. **18** (2016) 035005, [arXiv:1509.03920], [doi:10.1088/1367-2630/18/3/035005].
- [17] Chen, G. et al., *Quantum Computing Devices – Principles, Designs, and Analysis*, Routledge (2007), [ISBN:9780367390372].
- [18] Cho, G. Y., Gang, D., Kim, H.-C., *M-theoretic Genesis of Topological Phases*, J. High Energ. Phys. **2020** 115 (2020) [doi:10.1007/JHEP11(2020)115], [arXiv:2007.01532].
- [19] Chu, C., Lorscheid, O., Santhanam, R., *Sheaves and K-theory for \mathbb{F}_1 -schemes*, Adv. Math. **229** 4 (2012), 2239-2286, [arXiv:1010.2896], [doi:10.1016/j.aim.2011.12.023].
- [20] Clay Math Institute, *The Millennium Prize Problems*, [www.claymath.org/millennium-problems].
- [21] Cohen, F. R., *Introduction to configuration spaces and their applications*, in: *Braids*, Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore **19** (2009), 183-261, [doi:10.1142/9789814291415_0003].

- [22] DARPA, *Quantum Benchmarking Initiative* (2024), [www.darpa.mil/work-with-us/quantum-benchmarking-initiative].
- [23] Das Sarma, S., *Quantum computing has a hype problem*, MIT Tech Review (March 2022), [www.technologyreview.com/2022/03/28/1048355/quantum-computing-has-a-hype-problem/]
- [24] Das Sarma, S., *In search of Majorana*, Nature Physics **19** (2023), 165-170, [arXiv:2210.17365], [doi:10.1038/s41567-022-01900-9].
- [25] D’Auria, R., Fré, P., *Geometric Supergravity in $D = 11$ and its hidden supergroup*, Nucl. Phys. B **201** (1982), 101-140, [doi:10.1016/0550-3213(82)90376-5].
- [26] Donos, A., Gauntlett, J. P., Sonner, J., Withers, B., *Competing orders in M-theory: superfluids, stripes and metamagnetism*, J. High Energ. Phys. **2013** 108 (2013), [doi:10.1007/JHEP03(2013)108], [arXiv:1212.0871].
- [27] Donos, A., Gauntlett, J. P., Pantelidou, C., *Semi-local quantum criticality in string/M-theory*, J. High Energ. Phys. **2013** 103 (2013), [doi:10.1007/JHEP03(2013)103], [arXiv:1212.1462].
- [28] Duff, M., *The World in Eleven Dimensions: Supergravity, Supermembranes and M-theory*, IoP (1999), [ISBN:9780750306720].
- [29] Dul, F., *General Covariance from the Viewpoint of Stacks*, Lett Math Phys **113** (2023) 30, [arXiv:2112.15473], [doi:10.1007/s11005-023-01653-3].
- [30] Dyakonov, M. I., *Prospects for quantum computing: extremely doubtful*, Int. J. of Modern Physics: Conf. Series **33** (2014) 1460357, [arXiv:1401.3629], [doi:10.1142/S2010194514603573].
- [31] Ezratty O., *Where are we heading with NISQ?*, [arXiv:2305.09518].
- [32] Ezratty O., *Where are we heading with NISQ?*, blog post (2023), [www.oezratty.net/wordpress/2023/where-are-we-heading-with-nisq/].
- [33] Farb, B., Margalit, D., *A primer on mapping class groups*, Princeton University Press (2012), [doi:j.ctt7rkjw], [ISBN:9780691147949].
- [34] Ferraz, A., Gupta, K. S., Semenoff, G. W., Sodano, P. (eds), *Strongly Coupled Field Theories for Condensed Matter and Quantum Information Theory*, Springer Proceedings in Physics **239**, Springer (2020), [doi:10.1007/978-3-030-35473-2].
- [35] Fiorenza, D., Sati, H., Schreiber, U., *A higher stacky perspective on Chern-Simons theory*, in *Mathematical Aspects of Quantum Field Theories* Springer (2014), 153-211, [doi:10.1007/978-3-319-09949-1].
- [36] Fiorenza, D., Sati, H., Schreiber, U., *The WZW term of the M5-brane and differential cohomotopy*, J. Math. Phys. **56** (2015) 102301, [doi:10.1063/1.4932618], [arXiv:1506.07557].
- [37] Fiorenza, D., Sati, H., Schreiber, U., *T-Duality from super Lie n -algebra cocycles for super p -branes*, Adv. Theor. Math. Phys. **22** 5 (2018), [arXiv:1611.06536], [doi:10.4310/ATMP.2018.v22.n5.a3].
- [38] Fiorenza, D., Sati, H., Schreiber, U., *Twisted Cohomotopy implies M5 WZ term level quantization*, Commun. Math. Phys. **384** (2021), 403-432, [doi:10.1007/s00220-021-03951-0], [arXiv:1906.07417].
- [39] Fiorenza, D., Sati, H., Schreiber, U., *Twisted Cohomotopy implies M-theory anomaly cancellation on 8-manifolds*, Commun. Math. Phys. **377** (2020), 1961-2025, [doi:10.1007/s00220-020-03707-2], [arXiv:1904.10207].
- [40] Fiorenza, D., Sati, H., Schreiber, U. *Twisted cohomotopy implies twisted String structure on M5-branes* J. Math. Phys. **62** (2021) 042301, [arXiv:2002.11093], [doi:10.1063/5.0037786].
- [41] Fiorenza, D., Sati, H., Schreiber, U., *Twistorial Cohomotopy Implies Green-Schwarz anomaly cancellation*, Rev. Math. Phys. **34** 05 (2022) 2250013, [doi:10.1142/S0129055X22500131], [arXiv:2008.08544].
- [42] Fiorenza, D., Sati, H., Schreiber, U., *The Character map in Nonabelian Cohomology — Twisted, Differential and Generalized*, World Scientific, Singapore (2023), [doi:10.1142/13422], [arXiv:2009.11909].
- [43] Foss-Feig, M., Pagano, G., Potter, A. C., Yao, N. Y., *Progress in Trapped-Ion Quantum Simulation*, Ann. Rev. Condensed Matter Phys. (2024), [doi:10.1146/annurev-conmatphys-032822-045619], [arXiv:2409.02990].
- [44] Fowler, A. G., Hollenberg, L. C. L., *Scalability of Shor’s algorithm with a limited set of rotation gates*, Phys. Rev. A **70** (2007) 032329 [doi:10.1103/PhysRevA.103.032417].
- [45] Freedman, M., Hastings, M. B., Nayak, C., Qi, X.-L., Walker, K., Wang, Z., *Projective Ribbon Permutation Statistics: a Remnant of non-Abelian Braiding in Higher Dimensions*, Phys. Rev. B **83** 115132 (2011), [arXiv:10.1103/PhysRevB.83.115132], [arXiv:1005.0583].
- [46] Freedman, M., Kitaev, A., Larsen, M., Wang, Z., *Topological quantum computation*, Bull. Amer. Math. Soc. **40** (2003), 31-38, [arXiv:quant-ph/0101025], [doi:10.1090/S0273-0979-02-00964-3].

- [47] Gallier, J., Xu, D., *A Guide to the Classification Theorem for Compact Surfaces*, Springer (2013), [doi:10.1007/978-3-642-34364-3].
- [48] Gauntlett, J. P., Sonner, J., Wiseman, T., *Holographic superconductivity in M-Theory*, Phys. Rev. Lett. **103** (2009) 151601, [arXiv:0907.3796], [doi:10.1103/PhysRevLett.103.151601].
- [49] Gauntlett, J., Sonner, J., Wiseman, T., *Quantum Criticality and Holographic Superconductors in M-theory*, J. High Energy Phys. **2010** 60 (2010), [arXiv:0912.0512], [doi:10.1016/j.physletb.2009.12.017].
- [50] Gent, E., *Quantum Computing's Hard, Cold Reality Check*, IEEE Spectrum (Dec. 2023), [spectrum.ieee.org/quantum-computing-skeptics]
- [51] Gill, S. G., et al., *Quantum Computing: Vision and Challenges*, [arXiv:2403.02240].
- [52] Giotopoulos G., Sati, H., Schreiber, U., *Flux Quantization on 11d Superspace*, J. High Energy Phys. **2024** (2024) 82, [doi:10.1007/JHEP07(2024)082], [arXiv:2403.16456].
- [53] Giotopoulos, G., Sati, H., Schreiber, U., *Flux Quantization on M5-Branes* J. High Energy Phys. **2024** 140 (2024), [doi:10.1007/JHEP10(2024)140], [arXiv:2406.11304].
- [54] Giotopoulos, G., Sati, H., Schreiber, U., *Super L_∞ T-Duality and M-theory*, [arXiv:2411.10260].
- [55] Grady, D., Sati, H., *Differential cohomotopy versus differential cohomology for M-theory and differential lifts of Postnikov towers*, J. Geom. Phys. **165** (2021) 104203, [doi:10.1016/j.geomphys.2021.104203], [arXiv:2001.07640].
- [56] Griffiths, P., Morgan, J., *Rational Homotopy Theory and Differential Forms*, Progress in Mathematics **16**, Birkhäuser (1981, 2013), [doi:10.1007/978-1-4614-8468-4].
- [57] Gromov, A., Martinec, E. J., Ryu, S., *Collective excitations at filling factor 5/2: The view from superspace*, Phys. Rev. Lett. **125** (2020) 077601, [doi:10.1103/PhysRevLett.125.077601], [arXiv:1909.06384].
- [58] Grumblin, E., Horowitz, M. (eds.), *Quantum Computing: Progress and Prospects*, The National Academies Press (2019), [doi:10.17226/25196], [ISBN:9780309479691].
- [59] Gubser, S. S., Pufu, S. S., Rocha, F. D., *Quantum critical superconductors in string theory and M-theory*, Phys. Lett. B **683** (2010), 201-204, [arXiv:0908.0011], [doi:10.1016/j.physletb.2009.12.017].
- [60] Hannabuss, K. C., *T-duality and the bulk-boundary correspondence*, J. Geom. Phys. **124** (2018), 421-435, [doi:10.1016/j.geomphys.2017.11.016], [arXiv:1704.00278].
- [61] Hansen, V. L., *On the Space of Maps of a Closed Surface into the 2-Sphere*, Math. Scand. **35** (1974), 149-158, [doi:10.7146/math.scand.a-11542], [jstor:24490694].
- [62] Hartle, J. B., Taylor, J. R., *Quantum Mechanics of Paraparticles*, Phys. Rev. **178** (1969) 2043 [doi:10.1103/PhysRev.178.2043].
- [63] Hartnoll, S., Lucas, A., Sachdev, S., *Holographic quantum matter*, MIT Press (2018), [arXiv:1612.07324], [ISBN:9780262348010].
- [64] Hasebe, K., *Unification of Laughlin and Moore-Read states in SUSY quantum Hall effect*, Phys. Lett. A **372** 9 (2008), 1516-1520, [doi:10.1016/j.physleta.2007.09.071].
- [65] Hatcher, A., *Algebraic Topology*, Cambridge University Press (2002), [ISBN:9780521795401].
- [66] Herzog, C. P., Kovtun, P., Sachdev, S., Thanh, D. S., *Quantum critical transport, duality, and M-theory*, Phys. Rev. D **75** (2007) 085020, [arXiv:hep-th/0701036], [doi:10.1103/PhysRevD.75.085020].
- [67] Hoefler, T., Haener, T., Troyer, M., *Disentangling Hype from Practicality: On Realistically Achieving Quantum Advantage*, Commun. ACM **66** 5 (2023), 82-87, [doi:10.1145/3571725], [arXiv:2307.00523].
- [68] Huerta, J., Sati, H., Schreiber, U., *Real ADE-equivariant (co)homotopy and Super M-branes*, Communications in Mathematical Physics **371** (2019) 425 [doi:10.1007/s00220-019-03442-3], [arXiv:1805.05987].
- [69] Iqbal, M., Tantivasadakarn, N., Verresen, R. et al., *Non-Abelian topological order and anyons on a trapped-ion processor*, Nature **626** (2024), 505-511, [doi:10.1038/s41586-023-06934-4].
- [70] Jacak, J. E., *Topological approach to electron correlations at fractional quantum Hall effect*, Ann. Phys. **430** (2021) 168493, [doi:10.1016/j.aop.2021.168493].
- [71] James, I. M., *General Topology and Homotopy Theory*, Springer (1984), [doi:10.1007/978-1-4613-8283-6].
- [72] Jiang, B., Bouhon, A., Lin, Z.-K., Zhou, X., Hou, B., Li, F., Slager, R.-J., Jiang, J.-H., *Experimental observation of non-Abelian topological acoustic semimetals and their phase transitions*, Nature Physics **17** (2021), 1239-1246, [doi:10.1038/s41567-021-01340-x], [arXiv:2104.13397].
- [73] Jordan, S. P., *Quantum Computation Beyond the Circuit Model*, PhD thesis, MIT (2010) [arXiv:0809.2307]
- [74] Jordan, S. P., *Permutational Quantum Computing*, Quantum Information and Computation **10** (2010) 470 [arXiv:0906.2508], [doi:10.26421/QIC10.5-6-7].

- [75] Kak, S., *Prospects for Quantum Computing*, talk at CIFAR Nanotechnology program meeting, Halifax (November 2008), [arXiv:0902.4884].
- [76] Kallel, S., *Configuration Spaces and the Topology of Curves in Projective Space*, in: *Topology, Geometry, and Algebra: Interactions and new directions*, Contemporary Mathematics **279**, AMS (2001), 151–175, [doi:10.1090/conm/279].
- [77] Kallel, S., *Configuration spaces of points: A user's guide*, Encyclopedia of Mathematical Physics 2nd ed. **4** (2025), [doi:10.1016/B978-0-323-95703-8.00211-1], [arXiv:2407.11092].
- [78] Katz, S., Klemm, A., Vafa, C., *Geometric Engineering of Quantum Field Theories*, Nucl. Phys. B **497** (1997), 173-195, [doi:10.1016/S0550-3213(97)00282-4], [arXiv:hep-th/9609239].
- [79] Kitaev, A., *Unpaired Majorana fermions in quantum wires*, Physics-Uspekhi **44** 10S (2001), 131-136, [arXiv:cond-mat/0010440], [doi:10.1070/1063-7869/44/10S/S29].
- [80] Kitaev, A., *Fault-tolerant quantum computation by anyons*, Ann. Phys. **303** (2003), 2-30, [doi:10.1016/S0003-4916(02)00018-0], [arXiv:quant-ph/9707021].
- [81] Kitaev, A., *Anyons in an exactly solved model and beyond*, Ann. Phys. **321** 1 (2006), 2-111, [doi:10.1016/j.aop.2005.10.005].
- [82] Ko, K. H., Smolinsky, L., *The framed braid group and 3-manifolds*, Proc. Amer. Math. Soc. **115** (1992) 541-551 [doi:10.1090/S0002-9939-1992-1126197-1].
- [83] Kosinski, A., *Differential manifolds*, Academic Press (1993), [ISBN:978-0-12-421850-5].
- [84] Larmore, L. L., Thomas, E., *On the Fundamental Group of a Space of Sections*, Math. Scand. **47** 2 (1980), 232-246, [jstor:24491393].
- [85] Lau J. W. Z., Lim, K. H., Shrotriya H., Kwek, L. C., *NISQ computing: where are we and where do we go?*, AAPPS Bull. **32** 27 (2022), [doi:10.1007/s43673-022-00058-z].
- [86] Lee, S. T., Packer, J. A., *The Cohomology of the Integer Heisenberg Groups*, J. Algebra **184** 1 (1996), 230-250, [doi:10.1006/jabr.1996.0258].
- [87] Lidar, D. A., Brun, T. A. (eds.), *Quantum Error Correction*, Cambridge University Press (2013), [ISBN:9780521897877], [doi:10.1017/CB09781139034807].
- [88] Liu, S., *Anyon quantum dimensions from an arbitrary ground state wave function*, Nature Commun. **15** (2024) 5134, [doi:10.1038/s41467-024-47856-7], [arXiv:2304.13235].
- [89] Liu, Y., Zhao, T., Xiang, T., *Resolving Geometric Excitations of Fractional Quantum Hall States*, Phys. Rev. B **110** 195137 (2024), [arXiv:2406.11195], [doi:10.1103/PhysRevB.110.195137].
- [90] Lloyd, S., *Quantum computation with abelian anyons*, Quantum Information Processing **1** 1/2 (2002), 13-18, [doi:10.1023/A:1019649101654], [arXiv:quant-ph/0004010].
- [91] Lobb, A., *A feeling for Khovanov homology*, Notices AMS **71** 5 (2024), [doi:10.1090/noti2928].
- [92] Lutchyn, R. M. Sau, J. D., Das Sarma, S., *Majorana Fermions and a Topological Phase Transition in Semiconductor-Superconductor Heterostructures*, Phys. Rev. Lett. **105** (2010) 077001, [doi:10.1103/PhysRevLett.105.077001].
- [93] Manoliu, M., *Abelian Chern-Simons theory*, J. Math. Phys. **39** (1998), 170-206, [arXiv:dg-ga/9610001], [doi:10.1063/1.532333].
- [94] Massuyeau, G., *Lectures on Mapping Class Groups, Braid Groups and Formality* (2021), [massuyea.perso.math.cnrs.fr/notes/formality.pdf].
- [95] Mathai, V., Thiang, G. C., *T-Duality of Topological Insulators*, J. Phys. A: Math. Theor. **48** (2015) 42FT02, [arXiv:1503.01206], [doi:10.1088/1751-8113/48/42/42FT02].
- [96] Mathai, V., Thiang, G. C., *T-Duality Simplifies Bulk-Boundary Correspondence*, Commun. Math. Phys. **345** (2016), 675–701, [arXiv:1505.05250], [doi:10.1007/s00220-016-2619-6].
- [97] Morava, J., *A homotopy-theoretic context for CKM/Birkhoff renormalization* [arXiv:2307.10148].
- [98] Morava, J., *Some very low-dimensional algebraic topology* [arXiv:2411.15885].
- [99] Myers, D. J., Sati, H., Schreiber, U., *Topological Quantum Gates in Homotopy Type Theory*, Comm. Math. Phys. **405** 172 (2024), [arXiv:2303.02382], [10.1007/s00220-024-05020-8].
- [100] Naito, S., Osada, K., Fukui, T., *Fierz Identities and Invariance of Eleven-dimensional Supergravity Action*, Phys. Rev. D **34** (1986), 536-552, [doi:10.1103/PhysRevD.34.536]; Erratum Ibid. **35** (1986), 1536-1536, [doi:10.1103/PhysRevD.35.1536].
- [101] Nakamura, J., et al., *Direct observation of anyonic braiding statistics*, Nat. Phys. **16** (2020), 931–936, [doi:10.1038/s41567-020-1019-1].

- [102] Nielsen, M. A., Chuang, I. L., *Quantum computation and quantum information*, Cambridge University Press (2000), [doi:10.1017/CB09780511976667].
- [103] nLab, *Yang-Mills mass gap*, [ncatlab.org/nlab/show/Yang-Mills+mass+gap]
- [104] nLab, *cell structure of projective spaces*, [ncatlab.org/nlab/show/cell+structure+of+projective+spaces].
- [105] Okuyama, S., *The space of intervals in a Euclidean space*, *Algebr. Geom. Topol.* **5** (2005), 1555-1572, [arXiv:math/0511645], [doi:10.2140/agt.2005.5.1555].
- [106] Ohtsuki, T., *Quantum Invariants – A Study of Knots, 3-Manifolds, and Their Sets*, World Scientific (2001), [doi:10.2140/agt.2005.5.1555].
- [107] Pachos, J. K., *Quantum computation with abelian anyons on the honeycomb lattice*, *Int. J. Quantum Information* **4** 6 (2006), 947-954, [doi:10.1142/S0219749906002328], [arXiv:quant-ph/0511273].
- [108] Pachos, J. K., *Introduction to Topological Quantum Computation*, Cambridge University Press (2012), [doi:10.1017/CB09780511792908].
- [109] Papić, Z., Balram, A. C., *Fractional quantum Hall effect in semiconductor systems*, *Encyclopedia of Condensed Matter Physics* 2nd ed. **1** (2024), 285-307, [doi:10.1016/B978-0-323-90800-9.00007-X], [arXiv:2205.03421].
- [110] Polchinski, J., *Dualities of Fields and Strings*, *Studies in History and Philosophy of Modern Physics* **59** (C) (2017), 6-20, [arXiv:1412.5704].
- [111] Polychronakos, A. P., *Path Integrals and Parastatistics*, *Nucl. Phys. B* **474** (1996) 529-539 [arXiv:hep-th/9603179], [doi:10.1016/0550-3213(96)00277-5].
- [112] Pontrjagin, L., *Classification of continuous maps of a complex into a sphere, Communication I*, *Doklady Akademii Nauk SSSR* **19** 3 (1938), 147-149.
- [113] Prange, R. E., Girvin, S. M. (eds.), *The Quantum Hall Effect*, *Graduate Texts in Contemporary Physics*, Springer (1986, 1990), [doi:10.1007/978-1-4612-3350-3].
- [114] Preskill, J., *Quantum Computing in the NISQ era and beyond*, *Quantum* **2** 79 (2018), [arXiv:1801.00862], [doi:10.22331/q-2018-08-06-79].
- [115] Preskill, J., *Crossing the Quantum Chasm: From NISQ to Fault Tolerance*, talk at Q2B 2023, Silicon Valley (2023), [ncatlab.org/nlab/files/Preskill-Crossing.pdf].
- [116] Preskill, J., *Beyond NISQ: The Megaquop Machine*, talk at Q2B 2024 Silicon Valley (Dec. 2024), [www.preskill.caltech.edu/talks/Preskill-Q2B-2024.pdf].
- [117] Pu, S., Balram, A. C., Fremling, M., Gromov, A., Papić, Z., *Signatures of Supersymmetry in the $\nu = 5/2$ Fractional Quantum Hall Effect*, *Phys. Rev. Lett.* **130** (2023) 176501, [doi:10.1103/PhysRevLett.130.176501], [arXiv:2301.04169].
- [118] Rigolin, G., Ortiz, G., *The Adiabatic Theorem for Quantum Systems with Spectral Degeneracy*, *Phys. Rev. A* **85** 062111 (2012), [arXiv:1111.5333], [doi:10.1103/PhysRevA.85.062111].
- [119] Rowell, E. C., *An Invitation to the Mathematics of Topological Quantum Computation*, *J. Phys.: Conf. Ser.* **698** (2016) 012012, [doi:10.1088/1742-6596/698/1/012012], [arXiv:1601.05288].
- [120] Rowell, E. C., *Braids, Motions and Topological Quantum Computing*, [arXiv:2208.11762].
- [121] Sati, S., Schreiber, U., *Equivariant Cohomotopy implies orientifold tadpole cancellation*, *J. Geom. Phys.* **156** (2020), 103775, [doi:10.1016/j.geomphys.2020.103775], [arXiv:1909.12277].
- [122] Sati, H., Schreiber, U., *Proper Orbifold Cohomology*, [arXiv:2008.01101].
- [123] Sati, H., Schreiber, U. *Twisted Cohomotopy implies M5-brane anomaly cancellation*, *Lett. Math. Phys.* **111** (2021) 120, [doi:10.1007/s11005-021-01452-8].
- [124] Sati, H., Schreiber, U., *M/F-Theory as Mf-Theory*, *Rev. Math. Phys.* **35** 10 (2023), [arXiv:2103.01877], [doi:10.1142/S0129055X23500289].
- [125] Sati, H., Schreiber, U., *Equivariant principal ∞ -bundles*, [arXiv:2112.13654].
- [126] Sati, H., Schreiber, U., *Anyonic Defect Branes and Conformal Blocks in Twisted Equivariant Differential K-Theory*, *Rev. Math. Phys.* **35** 06 (2023) 2350009, [doi:10.1142/S0129055X23500095], [arXiv:2203.11838].
- [127] Sati, H., Schreiber, U., *The Quantum Monadology* [arXiv:2310.15735].
- [128] Sati, H., Schreiber, U., *Entanglement of Sections* [arXiv:2309.07245].
- [129] Sati, H., Schreiber, U., *Anyonic topological order in TED K-theory*, *Rev. Math. Phys.* (2023) **35** 03 (2023) 2350001, [arXiv:2206.13563], [doi:10.1142/S0129055X23500010].
- [130] Sati, H., Schreiber, U., *Flux quantization*, *Encyclopedia of Mathematical Physics* (2nd ed.) **4** (2025), 281-324, [doi:10.1016/B978-0-323-95703-8.00078-1], [arXiv:2402.18473].

- [131] Sati, H., Schreiber, U., *Quantum Observables of Quantized Fluxes*, Ann. Henri Poincaré (2024), [arXiv:2312.13037], [doi:10.1007/s00023-024-01517-z].
- [132] Sati, H., Schreiber, U., *Abelian Anyons on Flux-Quantized M5-Branes*, [arXiv:2408.11896].
- [133] Sati, H., Schreiber, U., *Topological QBits in Flux-Quantized Supergravity*, in: *Quantum Gravity and Computation*, Routledge (2025), [arXiv:2411.00628].
- [134] Sati, H., Schreiber, U., *The character map in Equivariant Twistorial Cohomotopy*, in: *Applied Algebraic Topology*, Beijing J. Pure Appl. Math. (2025), [arXiv:2011.06533].
- [135] Sati, H., Schreiber, U., *Anyons on M5-Probes of Seifert 3-Orbifolds via Flux-Quantization*, Letters in Mathematical Physics (2025, in print) [arXiv:2411.16852].
- [136] Sati, H., Valera, S., *Topological Quantum Computing*, Encyclopedia of Mathematical Physics 2nd ed **4** (2025), 325-345, [doi:10.1016/B978-0-323-95703-8.00262-7].
- [137] Sau S., *A Roadmap for a Scalable Topological Quantum Computer*, Physics **10** 68 (2017), [physics.aps.org/articles/v10/68].
- [138] Schreiber, U., *Quantum Language via Linear Homotopy Types*, ICMAT lecture notes (2025) [ncatlab.org/schreiber/show/Quantum+Language+via+Linear+Homotopy+Types].
- [139] Segal, G., *Configuration-spaces and iterated loop-spaces*, Invent. Math. **21** (1973), 213-221, [doi:10.1007/BF01390197].
- [140] Störmer, H. L., *Nobel Lecture: The fractional quantum Hall effect*, Rev. Mod. Phys. **71** (1999) 875, [doi:10.1103/RevModPhys.71.875].
- [141] Strom, J., *Modern classical homotopy theory*, Graduate Studies in Mathematics **127**, American Mathematical Society (2011), [ams:gsm/127].
- [142] Teo, J. C. Y., Kane, C. L. *Majorana Fermions and Non-Abelian Statistics in Three Dimensions*, Phys. Rev. Lett. **104** (2010) 046401, [doi:10.1103/PhysRevLett.104.046401], [arXiv:0909.4741].
- [143] Tiwari, A., Bzdušek, T., *Non-Abelian topology of nodal-line rings in PT-symmetric systems*, Phys. Rev. B **101** (2020) 195130, [doi:10.1103/PhysRevB.101.195130].
- [144] Tong, D., *The Quantum Hall Effect*, lecture notes (2016), [arXiv:1606.06687], [www.damtp.cam.ac.uk/user/tong/qhe/qhe.pdf].
- [145] von Klitzing, K., *The quantized Hall effect*, Rev. Mod. Phys. **58** 519 (1986), [doi:10.1103/RevModPhys.58.519].
- [146] Waintal, X., *The Quantum House Of Cards*, PNAS **121** 1 (2024) e2313269120, [arXiv:2312.17570], [doi:10.1073/pnas.2313269120].
- [147] Waldorf, K., *Geometric T-duality: Buscher rules in general topology*, Ann. Henri Poincaré **25** (2024), 1285–1358, [arXiv:2207.11799], [doi:10.1007/s00023-023-01295-0].
- [148] Wang, Y., Hu, Z., et al., *Qudits and high-dimensional quantum computing*, Front. Phys. **8** 479 (2020) [arXiv:2008.00959], [doi:10.3389/fphy.2020.589504].
- [149] Wang, Z., Hazzard, K. R. A., *Particle exchange statistics beyond fermions and bosons*, Nature **637** (2025) 314-318 [arXiv:2308.05203], [doi:10.1038/s41586-024-08262-7].
- [150] Wen, X.-G., *Vacuum Degeneracy of Chiral Spin State in Compactified Spaces*, Phys. Rev. B **40** 7387 (1989), [doi:10.1103/PhysRevB.40.7387].
- [151] Wen, X.-G., *Topological orders and Edge excitations in FQH states*, Adv. Phys. **44** 5 (1995) 405, [arXiv:cond-mat/9506066], [doi:10.1080/00018739500101566].
- [152] Wen, X.-G., Dagoatto, E., Fradkin, E., *Anyons on a torus*, Phys. Rev. B **42** (1990) 6110, [doi:10.1103/PhysRevB.42.6110].
- [153] Williams, L., *Configuration Spaces for the Working Undergraduate*, Rose-Hulman Undergrad. Math. J. **21** 1 (2020) 8, [arXiv:1911.11186], [rhumj:vol21/iss1/8].
- [154] Witten, E., *Three Lectures On Topological Phases Of Matter*, La Rivista del Nuovo Cimento **39** (2016), 313-370, [arXiv:1510.07698], [doi:10.1393/ncr/i2016-10125-3].
- [155] Wootton, J. R., *Dissecting Topological Quantum Computation*, PhD thesis, Leeds (2010), [etheses:1163].
- [156] Wootton, J. R., Pachos, J. K., *Universal Quantum Computation with Abelian Anyon Models*, Electron. Notes Theor. Comput. Sci. **270** 2 (2011), 209-218, [doi:10.1016/j.entcs.2011.01.032], [arXiv:0904.4373].
- [157] Wu, Q., Soluyanov, A. A., Bzdušek, T., *Non-Abelian band topology in noninteracting metals*, Science **365** (2019) 1273-1277 [doi:10.1126/science.aau8740], [arXiv:1808.07469].
- [158] Ye, C. et al., *Quantum Fourier Transform and Phase Estimation in Qudit System*, Commun. Theor. Phys. **55** 790 (2011) [doi:10.1088/0253-6102/55/5/11].

- [159] Zaanen, J., Liu, Y., Sun, Y.-W., Schalm, K., *Holographic Duality in Condensed Matter Physics*, Cambridge University Press (2015), [doi:10.1017/CB09781139942492].
- [160] Zeng, B., Chen, X., Zhou, D.-L., Wen, X.-G., *Quantum Information Meets Quantum Matter – From Quantum Entanglement to Topological Phases of Many-Body Systems*, Quantum Science and Technology (QST), Springer (2019), [arXiv:1508.02595], [doi:10.1007/978-1-4939-9084-9].
- [161] Zhang, S. C. *The Chern-Simons-Landau-Ginzburg theory of the fractional quantum Hall effect*, Int. J. Mod. Phys. B **06** 01 (1992), 25-58, [doi:10.1142/S0217979292000037].