## Identifying Anyonic Topological Order in Fractional Quantum Anomalous Hall Systems

Hisham Sati<sup>\*</sup> and Urs Schreiber<sup>†</sup>

Center for Quantum and Topological Systems, Division of Science, New York University, Abu Dhabi, UAE

(Dated: July 3, 2025)

Recently observed fractional quantum anomalous Hall materials (FQAH) are candidates for topological quantum hardware, but their required anyon states are elusive. We point out dependence on monodromy of the *fragile* band topology in 2-cohomotopy. An algebro-topological theorem of Larmore & Thomas (1980) then identifies FQAH anyons over momentum space. Admissible braiding phases are 2C-th roots of unity, for C the Chern number. This lays the foundation for understanding symmetry-protected topological order in FQAH systems, reducing the problem to computations in equivariant cohomotopy.

## MOTIVATION AND INTRODUCTION

The holy grail of quantum materials research is arguably (cf. [1]) the understanding and manipulation of:

- (i) crystalline topological phases of matter where electron Bloch states in gapped valence bands span topologically non-trivial vector bundles over Brillouin tori Î<sup>d</sup> of crystal momenta (cf. [2][3, §2]),
- (ii) exhibiting topological order where the system's ground state is degenerate and gapped, whence effectively (far above the lattice scale) that of a topological field theory (cf. [4][5]),
- (iii) hosting anyons whence homotopy classes of adiabatic deformations of the material, notably the *braiding* of soliton worldlines, act on these ground states as unitary Berry phases (cf. [6, §3][3, §3]).



FIG. 1. In an FQH system at unit filling fraction 1/K, each electron in the effectively 2-dimensional material is "bound" to K quanta of transverse magnetic flux, as an effective result of strong interaction. On this backdrop, each surplus flux quantum is like the lack of 1/Kth of an electron, and as such called a *quasi-hole*. Just these quasi-hole surplus flux quanta are the (abelian) anyons in FQH systems (cf. [7]), in that the quantum state of the system picks up a braiding phase  $\zeta \in \mathbb{C}^{\times}$  (a root of unity) when a pair of these swap positions.

Among candidates, fractional quantum Hall systems (FQH, cf. [7] & Fig. 1) stand out in that their (abelian) anyons are actually being observed in recent years, by independent groups and across different platforms (starting with [8], recent pointers in [9]). This makes FQH materials a candidate hardware for much anticipated topological quantum computers [4], plausibly necessary for future quantum computing of utility value.

While the large external magnetic fields required for typical FQH systems may hinder their practicality as hardware, it has rather recently been confirmed for various materials [10][11][12][13] that there also exist "anomalous" forms of FQH materials (FQAH), where the role of the external magnetic field is instead played by magnetic properties intrinsic to crystalline topological phases of matter called *fractional Chern insulators* (FCI, cf. [14][15] and Fig. 2).



FIG. 2. Fractional Chern insulators (FCI) exhibiting a fractional quantum anomalous Hall effect (FQAH) are typically realized [16, §II.A] by gapping Dirac cones of 2D 2-band systems.

Concretely, the hallmark properties of anomalous Hall systems are all due to the existence of non-vanishing *Berry curvature*  $\Omega(\vec{k})$  over the crystal's Brillouin torus of Bloch momenta,  $\vec{k} \in \hat{T}^2$  (cf. [17][18][19, §III.D][20, §5.1]), a measure for the electron state's dependence on their crystal momenta, which here takes the role otherwise played by the magnetic flux density  $F(\vec{x})$  in the

<sup>\*</sup> hsati@nyu.edu

<sup>&</sup>lt;sup>†</sup> us13@nyu.edu

ordinary space of positions  $\vec{x}$  inside FQH systems.

This analogy is quite strong; we suggest to think of it as a *duality* (cf. Fig. 3): The *Lorentz force* law which underlies the ordinary Hall effect and the *anomalous velocity law* which underlies the anomalous Hall effect are mirror images of each other under exchanging position space with momentum space and magnetic flux density with Berry curvature (an observation that goes back to [21] [22][23, §III.B], review in [24][17, (69)][19, §III.A][25][26, (12-13)]), see Fig. 3. This gives a good understanding of the anomalous Hall effect as a Hall effect in momentum space.



FIG. 3. There is a duality between the FQH effect and its anomalous FQAH version, under which the ordinary space of positions inside the 2D material is exchanged for the "reciprocal" space of crystal momenta, while the external magnetic flux density is exchanged for the Berry curvature of the Bloch bands over this momentum space.

This duality is particularly remarkable for our purpose, because the full definition of flux in position space is

- (i.) [27] known to require choices of *flux-quantization laws* which exhibit the total flux – hence dually: the Chern number (8) – as a "rationalized" shadow of maps to a classifying space, cf. (8), while
- (ii.) [28] exotic (below: *fragile*) flux-quantization laws were recently understood to control the existence and nature of anyonic solitons in FQH systems.

Just this identification of anyonic solitons in the dual FQAH systems had remained wide open but needs addressing for these systems to serve as topological quantum hardware.

We now present a resolution by:

- (i.) pointing out that the *fragile* band topology of (fractional) Chern insulators (while often conflated with the *stable* band topology) influences the adiabatic Berry holonomy (11) acting on the ground states,
- (ii.) computing the effect, verifying that it exhibits anyonic topological order over the Brillouin torus (Thm. 1 below).

We will invoke basics of algebraic topology [29]; for exposition in our context see [2][28, §A.3][30, §A][31, §A.2].

## METHODS AND RESULTS

First to note that generic fractional Chern insulators are (cf. [16, §II.A.1]) 2D 2-band systems whose Bloch Hamiltonian over the Brillouin torus  $\hat{T}^2$  of crystal momenta may be expanded in Pauli matrices  $\{\sigma_i\}_{i=1}^3$  as:

$$H_{\text{Blch}}: \ \widehat{T}^2 \longrightarrow \text{Mat}_{2\times 2}(\mathbb{C})$$
$$H_{\text{Blch}}(\vec{k}) = h_0(\vec{k}) + \underbrace{\sum_{i=1}^3 h_i(\vec{k}) \sigma_i}_{H(\vec{k})}, \quad h_{(-)}(-) \in \mathbb{R}, \quad (1)$$

where the system is *gapped* in that the difference of the two eigenvalues is non-vanishing:

$$|H(\vec{k})| := \sqrt{\sum_{i=1}^{3} (h_i(\vec{k}))^2} > 0,$$
 (2)

whence the valence bundle  $\mathcal{V}$  of lower eigenspaces (cf. Fig. 2) is the kernel bundle over  $\widehat{T}^2$  of the projectivized operators:

$$\mathcal{V} \simeq \ker(\mathrm{id} + H/|H|),$$
 (3)

which shows that the valence bundle is exactly encoded in a map  $\vec{h}/|H|$  to the 2-sphere (cf. [2, (8.3-4)]):

$$\widehat{T^2} \xrightarrow{\vec{h}/|H|} S^2 \xrightarrow{\sigma} \operatorname{Mat}_{2\times 2}(\mathbb{C}). \quad (4)$$

More concretely, the lower eigenspace bundle of (what we may identify as the "universal normalized 2-band Bloch Hamiltonian", denoted)  $\sigma$  in (4) is the *tautological* complex line bundle on the 2-sphere, the one with unit Chern class class (this is [32, Lem. 4.5.12]):

$$\ker(\mathrm{id} + \sigma) = 1 \in \mathbb{Z} \simeq H^2(S^2; \mathbb{Z}), \qquad (5)$$

whence it is the pullback of the universal complex line bundle on infinite projective space along the canonical inclusions (cf. [33, §3.8]):

$$\iota: S^2 \simeq \mathbb{C}P^1 \hookrightarrow \mathbb{C}P^2 \longleftrightarrow \mathbb{C}P^\infty := \bigcup_{N \to \infty} \mathbb{C}P^N, \qquad (6)$$

of complex projective *Grassmannian* moduli spaces of complex lines (fibers of the valence bundle) inside a fixed  $\mathbb{C}^{1+N}$  (the fiber of a trivial ambient Bloch bundle).

The infinite projective space  $\mathbb{C}P^{\infty}$  (6) is the *classi-fying space* for (complex line bundles and) integral 2cohomology, meaning that the homotopy classes of maps into it, hence the connected components  $\pi_0$  of the mapping space Map(-, -), are (cf. [34, Ex. 2.1]):

$$\pi_0 \operatorname{Map}(-, \mathbb{C}P^{\infty}) \simeq H^2(-; \mathbb{Z}), \qquad (7)$$

and the corresponding class of the valence bundle (3), via (4) and (6), is its (first and only) Chern number C:

$$\left[\widehat{T}^2 \xrightarrow{\stackrel{\widetilde{h}}{|H|}} S^2 \hookrightarrow \mathbb{C}P^{\infty}\right] = C \in \mathbb{Z} \simeq H^2(\widehat{T}^2; \mathbb{Z}), \quad (8)$$

characterizing the *topological phase* of the Chern insulator.

Now we come to the crucial subtlety not previously fully appreciated:

1. The valence bundle arises in (3) as a sub-bundle of the trivial rank = 2 bundle and as such is classified by the Grassmannian  $\mathbb{C}P^1 \simeq S^2$ , reflecting equivalence of topological phases under paths of deformations of the system small enough for the valence bundle not to mix with higher conduction bands. We note here that also  $\mathbb{C}P^1 \simeq S^1$  classifies a cohomology theory, namely an exotic "nonabelian" cohomology theory called 2-cohomotopy  $\pi^2$  [35, §VII][34, Ex. 2.7]:

$$\pi_0 \operatorname{Map}(-; S^2) =: \pi^2(-).$$
 (9)

Therefore, in the language of [36] (following [37], cf. [38]) 2-cohomotopy classifies the *fragile* band topology of generic FCIs: fragile, since its classification may in principle be broken by more drastic deformations.

2. But the class in (8) is that classified by  $\mathbb{C}P^{\infty}$  (6) and thus reflects equivalence under paths of exactly such more drastic deformations that may involve an arbitrary number of higher conduction bands, preserving only the *stable* band topology.

3. This situation is subtle and prone to be underappreciated because, in the present case of 2D materials, it makes no difference for the topological charge: The classical Hopf degree theorem (cf. [39, (35)]) says that the *n*-cohomotopy of closed *n*-manifolds coincides with their integral *n*-cohomology:

$$\begin{array}{cccc}
\pi^{2}(\widehat{T}^{2}) & & \xrightarrow{\sim} & H^{2}(\widehat{T}^{2}; \mathbb{Z}) \\
\parallel & & \parallel \\
\pi_{0} \operatorname{Map}(\widehat{T}^{2}, S^{2}) & \xrightarrow{\iota_{*}} & \operatorname{Map}(\widehat{T}^{2}, \mathbb{C}P^{\infty}),
\end{array}$$
(10)

whence authors may and do routinely address the Chern number (8) as the *winding number* of the map  $\vec{h}/|H|$  (4), thereby tacitly conflating the fragile band topology in (what we identified as)  $\pi^2$  with the stable band topology in  $H^2(-;\mathbb{Z})$  — which is harmless in the study of ordinary Chern insulators.

4. But as we now turn attention to the fractional situation of Chern insulators exhibiting an FQAH effect, we highlight that the whole point of the expected topological order is (cf. [6, §3][3]) the possibility that — while the external parameters of a system may adiabatically move through a closed path  $\gamma$  of deformations, here: returning the deformation class [ $\mathcal{V}$ ] of the valence bundle back to itself — the Hilbert space  $\mathcal{H}_C$  of anyonic quantum ground states may be transformed by a non-trivial unitary  $U_{\gamma}$  sensitive to (the homotopy class  $[\gamma]$  of) the actual path (cf. [40, §II.A.2][6, §3]).

In other words, the Hilbert space  $\mathcal{H}_C$  of anyonic ground states observable under deformations that allow excursions (of the valence bundle  $\mathcal{V}$  of Chern number C) into the first N Bloch bands above the gap must be a unitary representation  $\rho$  of the *fundamental group*  $\pi_1$  of maps to the relevant Grassmannian  $\mathbb{C}P^N$ :

Thereby, topological order is sensitive not just to the charge sector in  $\pi_0$  of the mapping space, but also to  $\pi_1$  of its connected components, and for that the distinction between fragile and stable band topology makes all the difference, in contrast to (10) — this is our main result to be communicated here:

**Theorem 1.** For fragile valence band topology, N = 1 (6), the Hilbert spaces in (11) are modules over this noncommutative algebra of topological observables:

$$Obs_{C} := \mathbb{C}\left[\pi_{1}\left(\operatorname{Map}\left(\widehat{T}^{2}, S^{2}\right), C\right)\right]$$
(12)  
$$\simeq \left\langle \widehat{W}_{\left[1\right]}, \widehat{W}_{\left[1\right]}, \widehat{\zeta} \mid \widehat{W}_{\left[1\right]}, \widehat{W}_{\left[1\right]}, \widehat{W}_{\left[1\right]} = \widehat{\zeta}^{2} \widehat{W}_{\left[1\right]}, \widehat{W}_{\left[1\right]}, \widehat{\zeta}^{2C} = 1\right\rangle,$$

where in each superselection sector (meaning: on irreducible representations) with  $C \neq 0$  the operator  $\hat{\zeta}$  acts by multiplication with a 2C-th root of unity:

$$\widehat{\zeta} |\psi\rangle = e^{\pi i \frac{p}{C}} |\psi\rangle, \quad p \in \left\{0, 1, \cdots, |C| - 1\right\}.$$
(13)

On the other hand, for all more stable band topologies,  $N \ge 2$  (6), the central generator  $\hat{\zeta}$  disappears and the algebra becomes commutative with its two generators identified with a basis of torus 1-cycles:

$$\mathbb{C}\left[\pi_1\left(\operatorname{Map}(\widehat{T}^2, \mathbb{C}P^{N>1}), C\right)\right] \simeq \mathbb{C}\left[H^1(\widehat{T}^2; \mathbb{Z})\right] \quad (14)$$

$$\simeq \left\langle \widehat{W}_{[0]}^{1}, \widehat{W}_{[0]}^{0} \middle| \widehat{W}_{[0]}^{1} \widehat{W}_{[0]}^{0} = \widehat{W}_{[0]}^{0} \widehat{W}_{[0]}^{1} \right\rangle.$$

Proof. With the identification of (11) it is a question in pure algebraic topology to compute the fundamental groups of these mapping spaces. For (14) this is elementary in the extreme case  $N = \infty$  (cf. [28, (15)]), using that  $\mathbb{C}P^{\infty} \simeq K(\mathbb{Z}, 2)$  is an Eilenberg-MacLane space. The computation for (12) is much more intricate: It turns out to be a theorem of Larmore & Thomas (1980) [41, Thm. 1] (also [42, Prop. 1.5]), following Hansen (1974) [43, Thm. 1] who obtained the result except for fixing the factor "2" in the exponent of  $\hat{\zeta}^{2C} = 1$ . The remaining case (14) for finite N is due to Kallel (2001) [42, Lem. 7.5]. That these purely algebro-topological results pertain to anyonic topological order of quantum Hall systems we recently observed in [28, §3.3-4].

With this, the statement (13) is a direct consequence of Schur's lemma, cf. [28, (201)]. (In the remaining case C = 0, a variant of (13) may be proven by invoking the modular equivariance expected of an effective topological field theory of anyons, see [28, §3.4].)

But now we observe that the algebra (12) is exactly the characteristic algebra of observables for anyons on a torus (here: on the Brillouin torus of crystal momenta) with braiding phase  $\zeta$ : This is due to [44, (4.9)][45, (4.14)], reviewed in [46, (4.21)][47, (5.28)], cf. Fig. 4.



FIG. 4. The relation in the algebra (12) says that an adiabatic parameter loop around a basis of torus cycles induces transformation of the topological states by multiplication with the square of a complex phase  $\zeta$  to be identified [28, Prop. 3.21] with an anyon braiding phase (cf. Fig. 1).

In particular, from this observable algebra (12) a variant of the Stone-von Neumann theorem theorem implies [28, §3.4] that the superselection sectors of  $\mathcal{H}_C$  have dimension  $\propto \operatorname{ord}(\zeta)$ , thus exhibiting the ground state degeneracy characteristic of topological order.

## CONCLUSION AND OUTLOOK

Highlighting that topological order is sensitive not just to the connected components  $\pi_0$  of the moduli space of band topologies, but crucially to its fundamental groups  $\pi_1$  (where the commonly conflated *fragile* and *stable* band topologies of Chern insulators may differ substantially) we have invoked mathematical results from algebraic topology to derive (predict) how, when the fragile topology of FQAH systems can be resolved in *cohomotopy*  $\pi^2$ , then topological order becomes detectable with braiding phases of anyons localized over the reciprocal space of crystal momenta. Curiously, this makes FQAH anyons be momentum-space duals of similarly cohomotopically classified anyons in FQH systems [28]. But realistic (fractional) Chern insulators of interested are typically subject to *crystalline symmetry*, forcing the Bloch Hamiltonians (1) to transform under certain crystal point group symmetries  $g \in G \subset \hat{T}^2$  as (cf. [4, §4.1])

$$H_{\text{Blch}}(g \cdot \vec{k}) = U_g(\vec{k}) \circ H_{\text{Blch}}(\vec{k}) \circ U_{g^{-1}}(\vec{k}), \qquad (15)$$

for compatible unitary operators

$$\begin{array}{rcl} \widehat{T}^2 \times G \longrightarrow \operatorname{SU}(2) \\ \\ (\vec{k},g) & \longmapsto & U_g(\vec{k}) \,. \end{array}$$

In view of (4) this means equivalently that the map characterizing the valence bundle is actually *G*-equivariant

$$\begin{array}{ccc} G \subset \operatorname{Pin}(2) & G \subset \operatorname{Spin}(3) \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\$$

We may observe that if and when the relevant deformations are all constrained to preserve this symmetry — whence one speaks of the topological phases being symmetry-protected (cf. [48][3, §2.3]) — the fragile topological moduli space of the system shrinks to the subspace Map $(-, -)^G \subset Map(-, -)$  of G-equivariant maps (16), whose connected components  $C_G$  in G-equivariant cohomotopy [32, Ex. 4.5.8][49, §6][50]

$$\pi_0 \left( \operatorname{Map}(\widehat{T}^2, S^2)^G \right) =: \pi_G^2(\widehat{T}^2)$$
(17)

measure the fragile symmetry-protected band topology.

These fragile symmetry-protected topological phases have not been considered before. We highlight that there is still a stabilization map (cf. [32, Prop. 4.5.17, Rem. 4.5.18) which coarsens and hence compares them to the classes in *G*-equivariant *K*-theory that did find much attention in this context (cf. [51]).

What is more, in view of Thm. 1 we immediately obtain a precise formula for the algebra of topological Berry phases to be expected in symmetry-protected FQAH systems in topological phases  $C_G$ :

$$GObs_{C_G} := \mathbb{C}\left[\pi_1\left(\operatorname{Map}\left(\widehat{T}^2, S^3\right)^G, C_G\right)\right].$$
(18)

This predicts anyonic topologically ordered FQAH phases whenever the G-symmetry (15) is such that (18) is non-abelian,

While there do not seem to be results available computing these equivariant cohomotopy algebras (18) for non-trivial *G*-actions, in generalization of Thm. 1, it is already noteworthy that hereby the problem of describing *G*-protected topological order in FQAH systems is reduced to a straightforward question in pure algebraic topology which may be handed to specialists for further investigation (cf. the end of [30, §6]). We thank Adrien Bouhon, Sadok Kallel, and Robert-Jan Slager for useful discussions.

This research was supported by *Tamkeen* under the NYU Abu Dhabi Research Institute grant CG008.

- K. L. Wang, Y. Wu, C. Eckberg, G. Yin, and Q. Pan, Topological quantum materials, MRS Bulletin 45, 373 (2020).
- [2] A. S. Sergeev, Topological insulators and geometry of vector bundles, SciPost Physics Lecture Notes 10.21468/scipostphyslectnotes.67 (2023), arXiv:2011.05004 [cond-mat.mes-hall].
- [3] H. Sati and U. Schreiber, Anyonic topological order in twisted equivariant differential (TED) K-theory, Reviews in Mathematical Physics 35, 2350001 (2022), arXiv:2206.13563 [hep-th].
- [4] T. D. Stanescu, Introduction to Topological Quantum Matter & Quantum Computation (CRC Press, 2020).
- [5] S. H. Simon, *Topological Quantum* (Oxford University Press, 2023).
- [6] D. J. Myers, H. Sati, and U. Schreiber, Topological Quantum Gates in Homotopy Type Theory, Communications in Mathematical Physics 405, 1 (2024), arXiv:2303.02382 [hep-th].
- [7] H. L. Stormer, Nobel Lecture: The fractional quantum Hall effect, Rev. Mod. Phys. 71, 875 (1999).
- [8] J. Nakamura, S. Liang, G. C. Gardner, and M. J. Manfra, Direct observation of anyonic braiding statistics, Nature Physics 16, 931 (2020), arXiv:2006.14115 [cond-mat.meshall].
- [9] A. Veillon, C. Piquard, P. Glidic, Y. Sato, A. Aassime, A. Cavanna, Y. Jin, U. Gennser, A. Anthore, and F. Pierre, Observation of the scaling dimension of fractional quantum Hall anyons, Nature 632, 517 (2024), arXiv:2401.18044 [cond-mat.mes-hall].
- [10] J. Cai, E. Anderson, C. Wang, X. Zhang, X. Liu, W. Holtzmann, Y. Zhang, F. Fan, T. Taniguchi, K. Watanabe, Y. Ran, T. Cao, L. Fu, D. Xiao, W. Yao, and X. Xu, Signatures of fractional quantum anomalous Hall states in twisted MoTe2, Nature **622**, 63 (2023).
- [11] Y. Zeng, Z. Xia, K. Kang, J. Zhu, P. Knüppel, C. Vaswani, K. Watanabe, T. Taniguchi, K. F. Mak, and J. Shan, Thermodynamic evidence of fractional Chern insulator in moiré MoTe2, Nature **622**, 69 (2023).
- [12] H. Park, J. Cai, E. Anderson, Y. Zhang, J. Zhu, X. Liu, C. Wang, W. Holtzmann, C. Hu, Z. Liu, T. Taniguchi, K. Watanabe, J.-H. Chu, T. Cao, L. Fu, W. Yao, C.-Z. Chang, D. Cobden, D. Xiao, and X. Xu, Observation of fractionally quantized anomalous Hall effect, Nature **622**, 74 (2023), arXiv:2308.02657 [cond-mat.mes-hall].
- [13] Z. Lu, T. Han, Y. Yao, A. P. Reddy, J. Yang, J. Seo, K. Watanabe, T. Taniguchi, L. Fu, and L. Ju, Fractional quantum anomalous Hall effect in multilayer graphene, Nature 626, 759 (2024).
- [14] N. Regnault and B. A. Bernevig, Fractional Chern Insulator, Physical Review X 1, 021014 (2011), arXiv:1105.4867 [cond-mat.str-el].
- [15] T. Neupert, C. Chamon, T. Iadecola, L. H. Santos, and C. Mudry, Fractional (Chern and topological) insulators, Physica Scripta **T164**, 014005 (2015), arXiv:1410.5828

[cond-mat.str-el].

- [16] C.-Z. Chang, C.-X. Liu, and A. H. MacDonald, Colloquium: Quantum anomalous Hall effect, Reviews of Modern Physics 95, 011002 (2023), arXiv:2202.13902 [condmat.mes-hall].
- [17] N. A. Sinitsyn, Semiclassical theories of the anomalous Hall effect, Journal of Physics: Condensed Matter 20, 023201 (2007), arXiv:0712.0183 [cond-mat.mes-hall].
- [18] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, Anomalous Hall effect, Reviews of Modern Physics 82, 1539 (2010), arXiv:0904.4154 [cond-mat.meshall].
- [19] D. Xiao, M.-C. Chang, and Q. Niu, Berry phase effects on electronic properties, Reviews of Modern Physics 82, 1959 (2010), arXiv:0907.2021 [cond-mat.mes-hall].
- [20] D. Vanderbilt, Berry Phases in Electronic Structure Theory: Electric Polarization, Orbital Magnetization and Topological Insulators (Cambridge University Press, 2018).
- [21] M.-C. Chang and Q. Niu, Berry Phase, Hyperorbits, and the Hofstadter Spectrum, Physical Review Letters 75, 1348 (1995), arXiv:cond-mat/9505021 [cond-mat].
- [22] M.-C. Chang and Q. Niu, Berry phase, hyperorbits, and the Hofstadter spectrum: Semiclassical dynamics in magnetic Bloch bands, Physical Review B 53, 7010 (1996).
- [23] G. Sundaram and Q. Niu, Wave-packet dynamics in slowly perturbed crystals: Gradient corrections and Berry-phase effects, Physical Review B 59, 14915 (1999), arXiv:cond-mat/9908003 [cond-mat].
- [24] D. Xiao, J. Shi, and Q. Niu, Berry Phase Correction to Electron Density of States in Solids, Physical Review Letters 95, 137204 (2005), arXiv:cond-mat/0502340 [condmat.mes-hall].
- [25] Y. D. Chong, Berry's phase and the anomalous velocity of Bloch wavepackets, Physical Review B 81, 052303 (2010).
- [26] M. A. Stephanov and Y. Yin, Chiral Kinetic Theory, Physical Review Letters 109, 162001 (2012), arXiv:1207.0747 [hep-ph].
- [27] H. Sati and U. Schreiber, Flux Quantization, in *Encyclopedia of Mathematical Physics* (Elsevier, 2025) pp. 281–324, arXiv:2312.12517 [hep-th].
- [28] H. Sati and U. Schreiber, Fractional Quantum Hall Anyons via the Algebraic Topology of Exotic Flux Quanta (2025), arXiv:2505.22144 [cond-mat.mes-hall].
- [29] A. Hatcher, Algebraic Topology (Cambridge University Press, 2002).
- [30] H. Sati and U. Schreiber, Engineering of Anyons on M5-Probes via Flux Quantization, SciPost Physics Lecture Notes (2025), arXiv:2501.17927 [hep-th].
- [31] H. Sati and U. Schreiber, Quantum Observables of Quantized Fluxes, Annales Henri Poincaré, 1 (2024), arXiv:2306.01214 [hep-th].
- [32] H. Sati and U. Schreiber, Equivariant Principal ∞-Bundles (Cambridge University Press, 2025) in press, ncatlab.org/schreiber/show/EPBund, arXiv:2112.13654 [math.AT].
- [33] H. Sati and U. Schreiber, M/F-theory as Mf-theory, Reviews in Mathematical Physics 35, 23500289 (2023), arXiv:2103.01877 [hep-th].
- [34] D. Fiorenza, H. Sati, and U. Schreiber, *The Char-acter Map in Non-abelian Cohomology: Twisted, Dif-ferential, and Generalized* (World Scientific, 2023) arXiv:2009.11909 [math.AT].

- [35] S.-T. Hu, *Homotopy Theory*, Pure and Applied Mathematics, Vol. 8 (Academic Press, New York and London, 1959).
- [36] A. Bouhon, T. Bzdušek, and R.-J. Slager, Geometric approach to fragile topology beyond symmetry indicators, Physical Review B 102, 115135 (2020), arXiv:2005.02044 [cond-mat.str-el].
- [37] H. C. Po, H. Watanabe, and A. Vishwanath, Fragile Topology and Wannier Obstructions, Physical Review Letters 121, 126402 (2018), arXiv:1709.06551 [condmat.str-el].
- [38] A. Bouhon, Non-Abelian and Euler multi-gap topologies in crystalline materials (2023), talk at: Quantum Information and Quantum Matter, CQTS @ NYU Abu Dhabi, ncatlab.org/nlab/files/Bouhon-MultiGapTopolgies.pdf.
- [39] D. Fiorenza, H. Sati, and U. Schreiber, Twisted Cohomotopy Implies M-Theory Anomaly Cancellation on 8-Manifolds, Communications in Mathematical Physics 377, 1961 (2020), arXiv:1904.10207 [hep-th].
- [40] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Non-Abelian anyons and topological quantum computation, Reviews of Modern Physics 80, 1083 (2008), arXiv:0707.1889 [cond-mat.str-el].
- [41] L. L. Larmore and E. Thomas, On the fundamental group of a space of sections, Mathematica Scandinavica 47, 232 (1980).
- [42] S. Kallel, Configuration spaces and the topology of curves in projective space, in *Topology, Geometry, and Algebra: Interactions and New Directions*, Contemporary Mathematics, Vol. 279, edited by M. Ando, G. Carlsson, R. L. Cohen, D. M. Davis, J. R. Harper, and W. S. Wilson (American Mathematical Society, 2001) pp. 151–175,

arXiv:math-ph/0003010 [math.ph].

- [43] V. L. Hansen, On the Space of Maps of a Closed Surface into the 2-Sphere., Mathematica Scandinavica 35, 149 (1974).
- [44] X. G. Wen and Q. Niu, Ground-state degeneracy of the fractional quantum Hall states in the presence of a random potential and on high-genus Riemann surfaces, Physical Review B 41, 9377 (1990).
- [45] R. Iengo, Anyon quantum mechanics and Chern-Simons theory, Physics Reports 213, 179 (1992).
- [46] E. Fradkin, Field Theories of Condensed Matter Physics (Cambridge University Press, 2013).
- [47] D. Tong, Lectures on the Quantum Hall Effect (2016), www.damtp.cam.ac.uk/user/tong/qhe.html.
- [48] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, Reviews of Modern Physics 88, 035005 (2016), arXiv:1505.03535 [cond-mat.mes-hall].
- [49] H. Sati and U. Schreiber, Geometric Orbifold Cohomology (CRC Press, 2026) (in press), arXiv:2008.01101 [math.DG].
- [50] H. Sati and U. Schreiber, Equivariant Cohomotopy implies orientifold tadpole cancellation, Journal of Geometry and Physics 156, 103775 (2020), arXiv:1909.12277 [hep-th].
- [51] L. Stehouwer, J. de Boer, J. Kruthoff, and H. Posthuma, Classification of crystalline topological insulators through *K*-theory, Advances in Theoretical and Mathematical Physics 25, 723 (2021), arXiv:1811.02592 [hep-th].