

Introduction to

Higher Gauge Theory

via Differential Nonabelian Cohomology
in Cohesive Homotopy Theory

HISHAM SATI^{1,2} AND URS SCHREIBER¹
hsati@nyu.edu us13@nyu.edu

¹Mathematics Program and Center for Quantum and Topological Systems (CQTS),
New York University Abu Dhabi, UAE

²The Courant Institute for Mathematical Sciences,
New York University, New York, USA

June 19, 2026

Abstract

This is a streamlined introduction to the global (infrared) completion of Maxwell-type higher gauge fields (as in the higher gauge sectors of higher dimensional supergravity and its brane probes) by electromagnetic flux quantization in differential nonabelian cohomology, using cohesive homotopy theory. Applications include D/NS brane charge in (unstable) K-theory, M-brane charge in unstable Cohomotopy and geometric engineering of topological quantum order on probe M5-branes.

keywords: higher gauge theory, flux quantization, nonabelian cohomology, differential cohomology, cohesive homotopy theory, supergravity, branes

MSC codes: Primary: 53C08, 81T13, 18N60, 55P62, 55N20; Secondary: 83E50, 81T30, 81V70

funding: by *Tamkeen UAE* under the *Abu Dhabi Research Institute Grant CG008*

Contents

1. Introduction	3
2. Cohesive Homotopy Theory	5
2.1. Geometry	6
2.1.1. Smooth Sets	6
2.1.2. Concreteness	8
2.2. Homotopy	11
2.2.1. Gauged Sets	11
2.2.2. Coherence	13
2.3. Geometric Homotopy	14
2.3.1. Smooth Gauged Sets	14
2.3.2. Cohesion	15
3. Higher Maxwell-type Equations of Motion	17
3.1. Flux Densities	17
3.1.1. Differential Form Moduli	17
3.1.2. Flux Density Moduli	18
3.2. Equations of Motion	19
3.2.1. Higher Maxwell-type EoM	19
3.2.2. Characteristic L_∞ -Algebra	21
4. Global Phase Space of Higher Gauge Fields	23
4.1. Flux Quantization	23
4.1.1. Rational Homotopy	24
4.1.2. Nonabelian Character	27
4.2. Global Completion	29
4.2.1. The Phase Space	30
4.2.2. The Observables	34
5. Enhancements	36
5.1. Sources and Branes	36
5.2. CS/BF-Like Fields	38
6. Examples and Applications	39
6.1. 11D/10D Supergravity	39
6.2. M5-Brane Probes	42
A. Basic Background	43
A.1. Higher Lie Algebras	43
A.2. Classifying Spaces	45
References	48

1. Introduction

The Problem. It is an open secret (cf. [Sch14]) that traditional methods of mathematical physics, sophisticated as they may be (cf. [HT92]), are insufficient for capturing the global (topological, deep infrared) structure of higher gauge theories, on which, however, the nature of the topological monopole/brane charges notably depends:

Already for ordinary gauge theory, the standard assumption (already more “advanced” than what most textbooks get to, cf. [GMS09, §2]) that field configurations are sections of an ordinary fiber bundle over spacetime, the *field bundle*, means to restrict attention to a single topological sector (to be contrasted with the global situation, cf. [BSS18]). For higher gauge fields (cf. [Alf25]) the assumption of an ordinary field bundle tacitly restricts to just the *trivial* topological sector.

Similarly, the usual Lagrangian densities for higher gauge fields, being functions of globally defined gauge potential forms, tacitly assume that the given theory is restricted to the trivial topological sector or to a single chart of spacetime — because only there are gauge potentials globally defined.

Flux Quantization. The global definition of higher gauge fields is given by *flux quantization* [SS25d], but this has traditionally been relegated, at best, to an afterthought when faced with inconsistencies emerging in a naive approach. For instance:

- (i) The action functional of the electron is found to be inconsistent (*anomalous*) unless the ambient Maxwell field is flux-quantized according to [Dir31] (cf. [Alv85; Bry93, §7.1; Fra11, §16.4e]).
- (ii) The action functional of the M2-brane is anomalous unless the ambient C-field is shifted flux-quantized according to [Wit97b; Wit97a].
- (iii) The global definition of the action functional of the M5-brane similarly requires a flux quantization condition on the ambient C-field [FSS21a].

In the first case, the mathematical structure of the flux quantization has become classical; in the second case, there are proposals [HS05, §2.7] and [FSS20, §3.4]. The last case appears to have been addressed explicitly only in [FSS21a] (cf. [Sat10]).

In general, the problem of global completion of (higher gauge sectors of) supergravity theories (cf. [nLa26b]) by flux quantization is only recently finding some attention, cf. also [LS22a; LS22b; GSS24a; GST25]. And yet, this is paramount for many questions: It may be premature to speculate about the UV-completion of 11D Supergravity (working title: “M-theory” [Duf99]) before investigating its IR-completion by proper flux quantization.

Cohesive Approach. To settle such questions in examples, one needs a formalism that sets the stage in generality. In particular one needs to have basic notions, such as a generalization of the notion of *phase space*, to the case of higher gauge fields with possibly nontrivial topological sectors.

The aim of this note is to lay out steps in this direction, following [SS24b; SS25d] but further streamlining and expanding on some details.

Our methods are necessarily *higher geometric* (cf. [CGX11; FSS15a; Sch16; Jur+19; FSS19; Alf25]), hence of *geometric homotopy theory* (cohesive ∞ -topos theory, [SS26e; SS26d; Sch13], exposition in [Sch25]), which we briefly recall in § 2. In lowest degree this means generalizing smooth manifolds to *smooth sets* [KS26, Def. 2.1] (for discussion in field theory see [GS25a; GS25b], further exposition is in [IM25]) while in general degree this concerns generalizing *Lie groupoids* (cf. [BH25; MM03]) to *smooth ∞ -groupoids* (smooth ∞ -stacks, for concise definitions see [FSS23, §1], more exposition is in [FSS15a]).

The Solution. Using these methods, the problem of global (topological, infrared) completion of higher gauge fields turns out to have an elegant solution; in outline:

- (i) The smooth moduli set of higher flux densities solving their Maxwell-type equations of motion is identified with that of closed (meaning: flat, Maurer-Cartan) differential forms,

$$\{\text{On-shell flux densities}\} \simeq \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \quad (1)$$

on a Cauchy surface X^d with coefficients in a characteristic L_∞ -algebra \mathfrak{a} encoding the duality symmetric Bianchi identities and the electromagnetic higher Gauss laws (Thm. 3.3).

- (ii) The admissible electromagnetic flux quantization laws are given by classifying spaces \mathcal{A} whose real Whitehead-bracket L_∞ -algebra reproduces the Gauss laws (§ 4.1.1):

$$\mathfrak{L}\mathcal{A} \simeq \mathfrak{a}. \quad (2)$$

- (iii) A *nonabelian character map* (generalizing the Chern character on K-theory)

$$\mathbf{Map}(X^d; \mathcal{A}) \xrightarrow{\text{ch}^{\mathcal{A}}} \int \Omega_{\text{dR}}^1(X^d; \mathfrak{a}) \quad (3)$$

sends (monopole/brane) *charges* and their higher global symmetries in the nonabelian $\Omega\mathcal{A}$ -cohomology of X^d to the flux densities and their deformations sourced by these charges (§ 4.1.2).

- (iv) The on-shell higher gauge field configurations globally completed by flux quantization in $\Omega\mathcal{A}$ -cohomology are triples of (a) on-shell fluxes \vec{B} , (b) compatible charges χ , (c) gauge potentials \vec{A} witnessing that these fluxes are sourced by these charges. That is, we have a homotopy cone of smooth ∞ -groupoids of this form (§ 4.2.1):

$$\begin{array}{ccc} & \xrightarrow{\text{charges}} & \mathbf{Map}(X^d, \mathcal{A}) \\ & \swarrow \vec{A} \text{ potentials} & \downarrow \text{ch}^{\mathcal{A}} \\ * & \searrow \vec{B} \text{ fluxes} & \\ & \xrightarrow{\eta^f} & \int \Omega_{\text{cl}}^1(X^d; \mathfrak{L}\mathcal{A}), \end{array} \quad (4)$$

whereby the completed phase space stack is the homotopy fiber product

$$\mathbf{Phs}(X^d; \mathcal{A}) = \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \times_{\int \Omega_{\text{cl}}^1(X^d; \mathfrak{a})} \mathbf{Map}(X^d, \mathcal{A}). \quad (5)$$

- (v) For linear Gauss laws this reproduces familiar global completions by flux quantization such as in ordinary differential cohomology (line bundles, bundle gerbes, etc.) and in differential K-theory (though there is large room for alternative choices not traditionally

appreciated); but it crucially generalizes also to the non-linear electric Gauss laws that are famously seen (but traditionally ignored) in higher-dimensional supergravity, whose global infrared completions thereby become accessible (§ 6.1).

- (vi) A twisted relative generalization of this procedure applies also to higher gauge fields on probe branes, such as notably to the self-dual flux on M5-branes (§ 6.2.)

In particular, there exists an admissible completion (“Hypothesis H”) of 11D SuGra with M5-probes at A_n -singularities, which serves to geometrically engineer the topological quantum order of fractional quantum Hall systems in fine detail, making experimentally accessible predictions, such as concerning nonabelian anyons attached to super-semiconductor heterostructures.

Comparison to the Literature. The higher gauge theory via differential nonabelian cohomology that we present (§ 4.2.1) is *different* (cf. Rem. 3.4 and Fig. 2) from that via *higher principal connections* ([Sch05, §11-12; BS07; FSS12; SSS12; SW13; Bor+25; Alf25, §2; RSW26; Wal26; Bun+26]):

The basis of our formulation is exactly the electromagnetic Bianchi identities / Gauss laws that characterize higher Maxwell-type physical equations of motion (§ 3.2) as they appear notably in type I/II/M supergravity (§ 6). Just choosing flux quantization laws compatible with these (§ 4.1) already determines the full higher gauge potential/field structure globally (§ 4.2), without need or justification for a further principality constraint. For instance, differential K-theory is not a higher principal connection theory, but is a special case of differential nonabelian cohomology (Ex. 4.2).

While our discussion thus pertains in particular to the realm of “M-theory” (see [Duf99; Sat10; Jur+19; FSS19]) — geometrically engineering topological quantum materials (§ 6.2) in a realistic $N = 1$ alternative to traditional large- N holographic condensed matter theory ([Her+07; GSW09; GPR10; GSW10; Don+13; DGP13], cf. [Bag19, §4.3]) — at no point do we rely on string folklore. All of the following constructions have definitions and all claims have proofs.

Acknowledgements.. We thank the organizers of the meetings *Higher Differential Geometry* (WIKO Greifswald, May 2026) and *6th Mini Symposium on Physics and Geometry* (RBI Zagreb, May 2026) where parts of these lectures were first presented.


2. Cohesive Homotopy Theory

The completed phase spaces of higher gauge fields in § 4 combine the differential geometry of flux densities with the homotopy theory of (“topological”) charge sectors, cf. Fig. 1.

Such unification of geometry with homotopy theory happens in *geometric homotopy theory* (in ∞ -*topoi* of simplicial sheaves [Bro73; Jar87; TV05; Lur09]) and for *differential geometry* specifically in *cohesive homotopy theory* (in *cohesive* ∞ -*topoi* [Sch13; SS26e; SS26d], lecture notes in [Sch18]).

We aim to provide an exposition of the basics, but we assume the reader is familiar with the basic concepts and facts of ordinary category theory, as presented in [Awo10; Sch18]. For further background reading cf. [Ric20].

Figure 1: The geometric homotopy theory of ∞ -topoi is the conceptual unification of geometry with homotopy theory, hence the theory of *higher geometry*. In our context, *fluxes* are (differential) geometric, *charges* are homotopical, and their unification happens through the construction of the *phase space* (§ 4.2.1) in the ambient cohesive ∞ -topos (§ 2.3).



2.1. Geometry

Consider the ordinary category of smooth manifolds with smooth maps between them (cf. [Tu11; Lee12]), and its full subcategory of Cartesian spaces (\mathbb{R}^n s for $n \in \mathbb{N}$):

$$\text{CartSp} \hookrightarrow \text{SmthMfd}. \quad (6)$$

We will denote by $*$:= \mathbb{R}^0 the *point manifold*.

One reason why nonperturbative quantum field theory appears technically hard is that spaces of field configurations that ought to be objects in differential geometry tend to fall outside the realm of available textbook constructions. For example, for X, Y smooth manifolds of positive dimension, with X non-compact (such as $X \equiv \text{spacetime}$ and $Y \equiv \mathbb{C}$ for scalar fields), the would-be smooth mapping space

$$\mathbf{Map}(X, Y) \in ?? \quad (7)$$

fails to be even an infinite-dimensional manifold.

And yet, it is perfectly clear what should count as a smooth map from a manifold like \mathbb{R}^n into this mapping space (a smooth *probe* of the mapping space), namely this should correspond to a smooth map of ordinary smooth manifolds to Y from the product of X with \mathbb{R}^n :

$$\begin{aligned} & \{ \mathbb{R}^n \longrightarrow \mathbf{Map}(X, Y) \} \\ & \simeq \{ \mathbb{R}^n \times X \longrightarrow Y \}. \end{aligned} \quad (8)$$

But since knowing all such *probes* of a generalized smooth space is going to be as good as knowing the space “itself”, we are to make this the definition next.

2.1.1. Smooth Sets

A *smooth set* \mathbf{X} is (equivalently encoded by) a presheaf $\text{Plt}(-, \mathbf{X})$ of sets on the category of Cartesian spaces (6), hence a functor

$$\begin{aligned} \text{CartSp}^{\text{op}} & \longrightarrow \text{Set} \\ \mathbb{R}^n & \longmapsto \text{Plt}(\mathbb{R}^n, \mathbf{X}) \end{aligned} \quad (9)$$

which we think of as assigning:

- to \mathbb{R}^n (for $n \in \mathbb{N}$) the set of smooth maps from this \mathbb{R}^n into \mathbf{X} (the *plots* of \mathbf{X} by the *probe* \mathbb{R}^n), and

- to smooth maps $\mathbb{R}^{n'} \xrightarrow{\phi} \mathbb{R}^n$ their precomposition operation $\phi^* : \text{Plt}(\mathbb{R}^n, \mathbf{X}) \rightarrow \text{Plt}(\mathbb{R}^{n'}, \mathbf{X})$ with such plots, defined thereby.

In this vein of characterizing spaces via the system of their *plots* by *probe spaces*, a *smooth map* $\mathbf{X} \xrightarrow{f} \mathbf{Y}$ between such smooth sets must, by postcomposition, be a rule that sends \mathbb{R}^n -plots ϕ of \mathbf{X} to \mathbb{R}^n -plots $f \circ \phi$ of \mathbf{Y} , compatible with the precomposition operation of plots, hence must be a natural transformation $f_* : \text{Plt}(-, \mathbf{X}) \Rightarrow \text{Plt}(-, \mathbf{Y})$ between the functors of plots (9).

So far, this appears to say that the category of smooth sets should be the category of presheaves over CartSp . However, some of this information ought in fact to be redundant: For probing the (smooth) structure of a smooth set \mathbf{X} it ought to already be sufficient to know just the restriction of these \mathbb{R}^n -plots to arbitrarily small open balls $\mathbb{D}_{r>0}^n \simeq \mathbb{R}^n \subset \mathbb{R}^n$ around any point of \mathbb{R}^n (which without restriction we may fix to be the origin), hence to know the *germs* of plots, to be denoted

$$\text{Plt}(\mathbb{G}^n, \mathbf{X}) := \bigcap_{r \in \mathbb{R}_{>0}} \text{Plt}(\mathbb{D}_r^n, \mathbf{X}). \quad (10)$$

Any natural transformation f_* between presheaves of plots restricts to a transformation of such germs of plots, and if that restriction is a natural *isomorphism*, then one says that f_* is a *local isomorphism*. We denote the class of local isomorphisms by

$$W_1 \subset \text{Mor}(\text{Func}(\text{CartSp}^{\text{op}}, \text{Set})). \quad (11)$$

Hence we should regard local isomorphisms as *actual* isomorphisms between (systems of plots of) smooth sets, a process known as *localization* $L^{W_1}(-)$ at the class of local isomorphisms. Doing so changes their category to the *sheaf topos* over CartSp , which we thereby recognize as the correct *category of smooth sets* (cf. [GS25a; GS25b; IM25]):

$$\text{SmothSet} := L^{W_1} \text{Func}(\text{CartSp}^{\text{op}}, \text{Set}) \simeq \text{Sh}(\text{CartSp}). \quad (12)$$

True to this effective identification of smooth sets \mathbf{X} with their systems of sets of plots, we will often abbreviate the latter as

$$\mathbf{X}(\mathbb{R}^n) := \text{Plt}(\mathbb{R}^n, \mathbf{X}). \quad (13)$$

For example, an ordinary smooth manifold becomes a smooth set by taking its probes to be the ordinary smooth functions $C^\infty(-, -)$ between smooth manifolds, a fully faithful construction known, in generality, as the *Yoneda embedding*:

$$\begin{array}{ccc} \text{SmothMfd} & \hookrightarrow & \text{SmothSet} \\ X & \mapsto & C^\infty(-, X). \end{array} \quad (14)$$

Interestingly, this raises a potential for inconsistency: While we started out *declaring* the sets $\text{Plt}(\mathbb{R}^n, \mathbf{X})$ of smooth maps from \mathbb{R}^n to a smooth set \mathbf{X} defined thereby, we now see, with (14), that there are also the actual sets of such smooth maps $\mathbb{R}^n \rightarrow \mathbf{X}$ between smooth sets. Remarkably, it is the full *Yoneda lemma* which ensures that these naturally coincide, after the fact:

$$\text{Plt}(\mathbb{R}^n, \mathbf{X}) \simeq \text{Hom}(\mathbb{R}^n, \mathbf{X}), \quad (15)$$

where $\text{Hom}(-, -)$ on the right denotes the hom-sets of SmothSet .

However, smooth sets may be considerably more general than smooth manifolds. For example, *moduli spaces* $\Omega_{\text{dR}}^n(*)$ of differential n -forms ($n \in \mathbb{N}$) exist as smooth sets (more

on this example in § 3.1.1):

$$\begin{aligned} \Omega_{\mathrm{dR}}^n(*) &\in \mathrm{SmthSet} && \text{smooth moduli set of differential } n\text{-forms} \\ \mathrm{Plt}(\mathbb{R}^n, \Omega_{\mathrm{dR}}^n) &:= \Omega_{\mathrm{dR}}^n(\mathbb{R}^n) && \text{ordinary set of smooth differential } n\text{-forms on } \mathbb{R}^n, \end{aligned} \quad (16)$$

where precomposition of plots by smooth functions is given by pullback of differential forms.

Moreover, it is straightforward to construct further smooth sets from given ones, following the logic of their plots. For instance, given a pair $\mathbf{X}, \mathbf{Y} \in \mathrm{SmthSet}$, their Cartesian product $\mathbf{X} \times \mathbf{Y} \in \mathrm{SmthSet}$ must – as plots – have pairs of plots into the factors, which is already the definition:

$$\mathrm{Plt}(\mathbb{R}^n, \mathbf{X} \times \mathbf{Y}) := \mathrm{Plt}(\mathbb{R}^n, \mathbf{X}) \times \mathrm{Plt}(\mathbb{R}^n, \mathbf{Y}). \quad (17)$$

Finally, in the category of smooth sets, the motivating problem (7) is naturally resolved: For any pair $\mathbf{X}, \mathbf{Y} \in \mathrm{SmthSet}$ we have that the set of smooth maps between them becomes itself a smooth set, with plots

$$\begin{aligned} \mathrm{Map}(\mathbf{X}, \mathbf{Y}) &\in \mathrm{SmthSet} \\ \mathrm{Plt}(\mathbb{R}^n, \mathrm{Map}(\mathbf{X}, \mathbf{Y})) &:= \mathrm{SmthSet}(\mathbb{R}^n \times \mathbf{X}, \mathbf{Y}). \end{aligned} \quad (18)$$

Standard results imply that this evident definition already implies natural isomorphisms

$$\mathrm{Map}(\mathbf{X} \times \mathbf{Y}, \mathbf{Z}) \simeq \mathrm{Map}(\mathbf{X}, \mathrm{Map}(\mathbf{Y}, \mathbf{Z})). \quad (19)$$

This exhibits the category $\mathrm{SmthSet}$ as being *Cartesian closed*, one of the key properties of what is called a *convenient category of spaces* (going back to [Ste67], cf. [BH11]). In fact, $\mathrm{SmthSet}$ is “particularly convenient” in that it is [Sch13] a *cohesive topos* in the sense of [Law07]. We begin describing this in § 2.1.2 and will exhibit the full structure in § 2.3.2.

2.1.2. Concreteness

We discuss the *concrete* objects among the smooth sets from § 2.1.1, identified with *diffeological spaces* (for which cf. [Igl13]).

Traditionally, mathematical notions of *space* tend to be conceived as sets of points equipped with further structure (topology, smooth structure, etc.). But smooth sets \mathbf{X} (12) exist whose underlying set of points, $\mathrm{Plt}(\mathbb{R}^0, \mathbf{X})$, does not support all their higher-dimensional plots.

For example, with $n \geq 1$ the smooth moduli set $\Omega_{\mathrm{dR}}^n(*)$ (16) has a single point (the zero n -form on \mathbb{R}^0) but an infinite set of higher k -dimensional plots!

But we may neatly characterize those smooth sets that *are* underlying sets of points with extra structure, as the *concrete* smooth sets, commonly known as *diffeological spaces*. This proceeds as follows.

The Flat Modality. First, observe that the assignment of underlying sets of points is a functor

$$\begin{aligned} \mathrm{SmthSet} &\xrightarrow{\mathrm{Pnt}} \mathrm{Set} \\ \mathbf{X} &\longmapsto \mathrm{Plt}(*, \mathbf{X}), \end{aligned} \quad (20)$$

which has a fully faithful left adjoint:

$$\begin{array}{ccc} & \longleftarrow \mathrm{Dsc} \longrightarrow & \\ & | & \\ \mathrm{SmthSet} & \xrightarrow{\mathrm{Pnt}} & \mathrm{Set}. \end{array} \quad (21)$$

Namely, $\text{Dsc}(S)$ is the *discrete smooth structure* on $S \in \text{Set}$, whose only smooth plots are constant maps:

$$\text{Plt}(\mathbb{R}^n, \text{Dsc}(S)) = S. \quad (22)$$

The composite functor

$$\flat : \text{SmthSet} \xrightarrow{\text{Pnt}} \text{Set} \xleftarrow{\text{Dsc}} \text{SmthSet} \quad (23)$$

sends a smooth set to its underlying set of points, regarded as a discrete smooth set:

$$\text{Plt}(\mathbb{R}^n, \mathbf{X}) \simeq \text{Hom}(*, \mathbf{X}). \quad (24)$$

By the adjunction property, this comes with an evident natural transformation:

$$\begin{array}{ccc} \flat \mathbf{X} & \xrightarrow{\epsilon_{\mathbf{X}}^{\flat}} & \mathbf{X} \\ \text{Hom}(*, \mathbf{X}) & \longrightarrow & \text{Hom}(\mathbb{R}^n, \mathbf{X}) \\ (* \xrightarrow{x} \mathbf{X}) & \longmapsto & (\mathbb{R}^n \rightarrow * \xrightarrow{x} \mathbf{X}), \end{array} \quad (25)$$

which has the universal property that it uniquely factors smooth maps out of any $\flat \mathbf{Y}$:

$$\begin{array}{ccc} & \overbrace{\quad \quad \quad}^{\forall} & \\ \flat \mathbf{Y} & \xrightarrow{\exists!} \flat \mathbf{X} & \xrightarrow{\epsilon_{\mathbf{X}}^{\flat}} \mathbf{X}. \end{array} \quad (26)$$

In words, this just says that a map to \mathbf{X} from a discrete set of points factors through the underlying set of points of \mathbf{X} .

The Sharp Modality. But the functor Pnt , furthermore, has a fully faithful right adjoint:

$$\begin{array}{ccc} & \longleftarrow \text{Dsc} \longrightarrow & \\ \text{SmthSet} & \xrightarrow{\text{Pnt}} \text{Set} & \\ & \longleftarrow \text{Cht} \longrightarrow & \end{array} \quad (27)$$

Here $\text{Cht}(S)$ is the *chaotic smooth structure* on $S \in \text{Set}$, for which *all* maps from the discrete underlying set of a probe space count as smooth plots:

$$\text{Plt}(\mathbb{R}^n, \text{Cht}(S)) = \text{Hom}(\flat \mathbb{R}^n, \text{Dsc}(S)). \quad (28)$$

This chaotic smooth structure is typically not of interest in itself, but it serves to characterize concrete smooth sets, as follows.

The composite functor

$$\sharp : \text{SmthSet} \xrightarrow{\text{Pnt}} \text{Set} \xleftarrow{\text{Cht}} \text{SmthSet} \quad (29)$$

sends a smooth set to the chaotic smooth structure on its underlying set of points,

$$\text{Plt}(\mathbb{R}^n, \sharp \mathbf{X}) \simeq \text{Hom}(\flat \mathbb{R}^n, \mathbf{X}). \quad (30)$$

By the adjunction property, this receives a natural transformation from the identity, given by restricting \mathbb{R}^n -plots to all points in \mathbb{R}^n :

$$\begin{array}{ccc} \mathbf{X} & \xrightarrow{\eta_{\mathbf{X}}^{\sharp}} & \sharp \mathbf{X} \\ \text{Hom}(\mathbb{R}^n, \mathbf{X}) & \xrightarrow{(\epsilon_{\mathbb{R}^n}^{\flat})^*} & \text{Hom}(\flat \mathbb{R}^n, \mathbf{X}) \\ (\mathbb{R}^n \xrightarrow{\phi} \mathbf{X}) & \longmapsto & (\flat \mathbb{R}^n \xrightarrow{\epsilon_{\mathbb{R}^n}^{\flat}} \mathbb{R}^n \xrightarrow{\phi} \mathbf{X}), \end{array} \quad (31)$$

which has the universal property that it uniquely factors smooth maps into any $\sharp\mathbf{Y}$:

$$\mathbf{X} \xrightarrow{\eta_{\mathbf{X}}^{\sharp}} \sharp\mathbf{X} \dashrightarrow^{\exists!} \sharp\mathbf{Y} \quad \begin{array}{c} \lrcorner \\ \vee \\ \lrcorner \end{array} \quad (32)$$

Concretification. With this, we may now say that $\mathbf{X} \in \text{SmthSet}$ is *concrete* iff $\eta_{\mathbf{X}}^{\sharp}$ (31) is a monomorphism:

$$\mathbf{X} \text{ concrete} \quad \Leftrightarrow \quad \mathbf{X} \xrightarrow{\eta_{\mathbf{X}}^{\sharp}} \sharp\mathbf{X}, \quad (33)$$

because this means that the smooth probe maps $\mathbb{R}^n \rightarrow \mathbf{X}$ form a subset of the maps of underlying points from $\flat\mathbb{R}^n$ to $\flat\mathbf{X}$. Such concrete smooth sets are equivalently known as *diffeological spaces* (for which cf. [Igl13]):

$$\{\mathbf{X} \in \text{SmthSet} \mid \mathbf{X} \text{ concrete}\} \simeq \text{diffeological spaces}. \quad (34)$$

Better yet, every smooth set \mathbf{X} has a universal approximation by a concrete one, its *concretification* $\sharp_1\mathbf{X}$, given by the epi/mono factorization (cf. [Bor94, §4.4]) of $\eta_{\mathbf{X}}^{\sharp}$ through its image:

$$\mathbf{X} \xrightarrow{\eta_{\mathbf{X}}^{\sharp_1}} \sharp_1\mathbf{X} \hookrightarrow \sharp\mathbf{X}. \quad \begin{array}{c} \lrcorner \\ \vee \\ \lrcorner \end{array} \quad (35)$$

In words, the plots of $\sharp_1\mathbf{X}$ are those of \mathbf{X} after forgetting all information that disappears when restricting to points of probe spaces.

A key example of concretification is that of smooth moduli sets of differential forms, see (90).

Noticing that $\text{Pnt}(\sharp_1\mathbf{X}) \simeq \text{Pnt}(\mathbf{X})$, naturally, we have natural isomorphisms

$$\begin{aligned} \sharp\sharp_1\mathbf{X} &\simeq \sharp\mathbf{X}, \\ \flat\sharp_1\mathbf{X} &\simeq \flat\mathbf{X}. \end{aligned} \quad (36)$$

In variation of (32), a smooth map to a concrete smooth set (33) factors uniquely through the concretification (35):¹

$$\mathbf{X} \xrightarrow{\eta_{\mathbf{X}}^{\sharp_1}} \sharp_1\mathbf{X} \dashrightarrow^{\exists!} \sharp_1\mathbf{Y} \quad \begin{array}{c} \lrcorner \\ \vee \\ \lrcorner \end{array} \quad (37)$$

We highlight that the points of a concretification are the original points:

$$\flat\sharp_1\mathbf{X} \simeq \flat\mathbf{X} \quad (38)$$

This is clear on plots. An abstract way to see it is that the image factorization in a topos is the colimit of a limit (the coequalizer of the kernel pair) and both are preserved by the left and right adjoint functor \flat .

¹The universal property (37) follows from the orthogonal epi/mono factorization in SmthSet , which implies a unique dashed lift in the following diagram:

$$\begin{array}{ccc} \mathbf{X} & \xrightarrow{f} & \sharp_1\mathbf{Y} \\ \downarrow \eta_{\mathbf{X}}^{\sharp_1} & & \parallel \\ \sharp_1\mathbf{X} & \dashrightarrow & \sharp_1\mathbf{Y} \\ \downarrow \lrcorner & & \downarrow \lrcorner \\ \sharp\mathbf{X} & \xrightarrow{\sharp f} & \sharp\mathbf{Y}. \end{array}$$

Cohesion. Finally, there is a further left adjoint functor to the triple (27), which may be understood as taking smooth sets to their sets of *connected components*. The resulting adjoint quadruple [Sch13] exhibits SmthSet as a *cohesive topos* in the sense of [Law07]:

$$\begin{array}{ccc} \times & \xrightarrow{\quad} & \\ & \downarrow & \\ & \leftarrow \text{Dsc} \rightarrow & \\ \text{SmthSet} & \xrightarrow{\quad \text{Pnt} \rightarrow} & \text{Set} \\ & \downarrow & \\ & \leftarrow \text{Cht} \rightarrow & \end{array} \quad (39)$$

But this further left adjoint functor comes into its own only after generalization from the cohesive 1-topos of smooth sets to the cohesive ∞ -topos of smooth ∞ -groupoids. This is what we turn to next in § 2.2.

2.2. Homotopy

2.2.1. Gauged Sets

Where the term *group* is a historically evolved shorthand for *group of symmetries* of a single object, the term *groupoid* is to be understood as the *consistent collection of symmetry transformations* between possibly several objects. In a context specifically of *gauge symmetries* we have, as a slogan, that *groupoids are sets with gauge equivalences* and *higher groupoids are sets with higher gauge-of-gauge equivalences*. Or for short: *Groupoids are gauged sets*.

For example, where the collection of scalar field configurations on Minkowski spacetime $\mathbb{R}^{1,d}$ forms the plain set

$$\{ \Phi \mid \Phi \in C^\infty(\mathbb{R}^{1,d}, \mathbb{C}) \}, \quad (40)$$

the field configurations of a gauge field with gauge group G (and Lie algebra \mathfrak{g}) form the groupoid which we may schematically but suggestively denote as

$$\left\{ A \xrightarrow{g} A' = g(d + A)g^{-1} \mid \begin{array}{l} A, A' \in \Omega_{\text{dR}}^1(\mathbb{R}^{1,d}; \mathfrak{g}) \\ g \in C^\infty(\mathbb{R}^{1,d}, G) \end{array} \right\}, \quad (41)$$

while the configurations of the B-field form the 2-groupoid, which in this schematic but suggestive vein we may denote as:

$$\left\{ \begin{array}{ccc} B' = B + da & & \\ \begin{array}{ccc} a \nearrow & \Downarrow \lambda & \searrow a' \\ B & \xrightarrow{a+a'+\lambda d\lambda^{-1}} & B'' = B' + da' \end{array} & \left| \begin{array}{l} B, B', B'' \in \Omega_{\text{dR}}^2(\mathbb{R}^{1,d}) \\ a, a' \in \Omega_{\text{dR}}^1(\mathbb{R}^{1,d}) \\ \lambda \in C^\infty(\mathbb{R}^{1,d}, \text{U}(1)) \end{array} \right. & \end{array} \right\}. \quad (42)$$

And so on.

To make this precise, it is expedient to stick to the strategy of defining generalized spaces by the sets of *plots* by basic *probe spaces* which they admit. In the present case, as the above examples indicate, the elementary *n-fold gauge transformation* should be thought of as the *n-dimensional generalization of the cellular triangle*, known as the cellular *n-simplex* Δ^n . The cellular maps between these cellular *n-simplices* (mapping vertices to vertices, edges to edges, etc.) form a category accordingly known as the *simplex category* Δ .

Hence, a higher gauged set \mathcal{X} should be characterized by the system of plots $\text{Plt}(\Delta^n, \mathcal{X})$ of the shape of the standard simplices, forming a functor

$$\begin{array}{ccc} \Delta^{\text{op}} & \longrightarrow & \text{Set} \\ \Delta^n & \longmapsto & \text{Plt}(\Delta^n, \mathcal{X}). \end{array} \quad (43)$$

This suggests that (the systems of plots of) higher gauged sets form the functor category

$$\mathbb{S}\text{Set} := \text{Func}(\Delta^{\text{op}}, \text{Set}) \quad (44)$$

known as the category of *simplicial sets* (cf. [Fri12]).

But in more detail, we should still require that n -fold gauge transformations may consistently be composed and have inverses, up to higher gauge transformations. To this end, we recognize the boundary $\partial\Delta^n$ of the n -simplex with its k th $(n-1)$ -face $d_k(\Delta^n)$ removed, denoted

$$\Lambda_k^n := \partial\Delta^n - d_k\Delta^n \in \mathbb{S}\text{Set}, \quad (45)$$

as an abstract situation of n composable $(n-1)$ -transformations, some of them inverted. Then the requirement of composites and inverses in a simplicial set \mathcal{X} means that every image $\Lambda_k^n \rightarrow \mathcal{X}$ of this situation in \mathcal{X} may be completed to a full n -simplex $\Delta^n \rightarrow \mathcal{X}$, hence that \mathcal{X} admits all dashed lifts in diagrams of the following form:

$$\mathcal{X} \text{ Kan complex} \quad \Leftrightarrow \quad \forall_{\substack{n \geq 1, \\ k \leq n}} \quad \begin{array}{ccc} \Lambda_k^n & \xrightarrow{\forall} & \mathcal{X} \\ \downarrow & \nearrow \exists & \\ \Delta^n & & \end{array} \quad (46)$$

We write

$$\text{KanSSet} \subset \mathbb{S}\text{Set} \quad (47)$$

for the full subcategory of simplicial sets on those that satisfy this *Kan fibrancy condition* (46).

By an ∞ -category we may understand a category whose hom-sets are promoted to (hence which is *enriched in*) such Kan simplicial sets. In such an ∞ -category we have a notion of morphisms being *homotopy equivalences* if they have inverses up to gauge transformation.

There is a fairly evident notion of *homotopy groups* of such *Kan fibrant simplicial sets*. Let

$$W_{\text{wh}} \subset \text{Mor}(\text{KanSSet}) \quad (48)$$

be the class of maps that induce isomorphisms on all these homotopy groups: the *simplicial weak homotopy equivalences*. Then there is an ∞ -category which is the universal solution to turning weak homotopy equivalences into actual homotopy equivalences, denoted

$$\text{Grpd}_\infty := L^{W_{\text{wh}}} \text{KanSSet}. \quad (49)$$

This is the ∞ -category of ∞ -groupoids, making precise the idea of the *gauged category of gauge sets*.

Ordinary (ungauged) sets are faithfully included here as the *0-truncated* ∞ -groupoids ($\text{Set} \simeq \text{Grpd}_0$), with the inclusion having a left adjoint (*0-truncation*):

$$\text{Grpd}_\infty \begin{array}{c} \xrightarrow{\tau_0} \\ \perp \\ \longleftarrow \end{array} \text{Set}. \quad (50)$$

By adjunction, the composite ∞ -functor

$$[-]_0 : \text{Grpd}_\infty \xrightarrow{\tau_0} \text{Set} \hookrightarrow \text{Grpd}_\infty \quad (51)$$

comes with a natural transformation

$$\begin{array}{ccc} \text{gauged set} & \mathcal{X} & \xrightarrow{\eta_{\mathcal{X}}^{[-]_0}} & [\mathcal{X}]_0 & \text{set of gauge equivalence classes} \end{array} \quad (52)$$

with the universal property that it uniquely factors maps to 0-truncated ∞ -groupoids:

$$\mathcal{X} \xrightarrow{\eta_{\mathcal{X}}^{[-1]_0}} [\mathcal{X}]_0 \overset{\exists!}{\dashrightarrow} [\mathcal{Y}]_0.$$
(53)

As indicated in (52), with $\mathcal{X} \in \text{Grpd}_{\infty}$ thought of as a gauge set, its 0-truncation $[\mathcal{X}]_0 \in \text{Set}$ is its set of *gauge equivalence classes*.

2.2.2. Coherence

The grand insight of ∞ -category theory is that it proceeds essentially along the lines of ordinary category theory (a point originally made in [Joy08]), with equalities generalized to homotopies and uniqueness generalized to contractible ∞ -groupoids of choices.

For instance:

- A *Cartesian/pullback square* over a pair of coincident maps of (smooth) sets

$$\begin{array}{ccc} & & \mathbf{Y} \\ & & \downarrow \\ \mathbf{X} & \longrightarrow & \mathbf{B} \end{array}$$
(54)

is a commuting square (meaning that the diagonal composite maps are *equal*)

$$\begin{array}{ccc} \mathbf{X} \times_{\mathbf{B}} \mathbf{Y} & \longrightarrow & \mathbf{Y} \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{X} & \longrightarrow & \mathbf{B}, \end{array}$$
(55)

which is universal with this property, in that for any other such square there is a *unique* dashed comparison map

$$\begin{array}{ccc} \mathbf{Q} & & \\ \dashrightarrow & & \searrow \\ \mathbf{X} \times_{\mathbf{B}} \mathbf{Y} & \longrightarrow & \mathbf{Y} \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{X} & \longrightarrow & \mathbf{B} \end{array}$$
(56)

so that the two resulting triangles commute.

- A *homotopy Cartesian/pullback square* over a pair of coincident maps of (smooth, cf. below) ∞ -groupoids

$$\begin{array}{ccc} & & \mathbf{y} \\ & & \downarrow \\ \mathbf{x} & \longrightarrow & \mathbf{B} \end{array}$$
(57)

is a square that commutes up to a specified *homotopy* (often notationally suppressed!)

$$\begin{array}{ccc} \mathbf{x} \times_{\mathbf{B}} \mathbf{y} & \longrightarrow & \mathbf{y} \\ \downarrow & \lrcorner & \downarrow \\ \mathbf{x} & \longrightarrow & \mathbf{B}, \end{array}$$
(58)

which is universal with this property in that for any other such square the ∞ -groupoid

of compatible dashed comparison data

$$\begin{array}{ccc}
 \mathcal{Q} & & \\
 \swarrow \text{---} & & \searrow \\
 \mathcal{X} \times_{\mathcal{B}} \mathcal{Y} & \longrightarrow & \mathcal{Y} \\
 \downarrow & \lrcorner & \downarrow \\
 \mathcal{X} & \longrightarrow & \mathcal{B}
 \end{array}
 \tag{59}$$

is *contractible*.

Several ingenious ways have been found to handle such *homotopy coherence*. A generally useful one is *model category* theory, which is a toolbox to cleverly model such situations in (for our case) the ordinary category of simplicial presheaves $\text{Fun}(\text{CartSp}^{\text{op}}, \text{SSet})$ such that the homotopy coherent universal construction (say a homotopy pullback) is modeled by the corresponding ordinary universal construction (say an ordinary pullback). A concise introduction to these methods, aimed at our context, is in [FSS23, §1].

2.3. Geometric Homotopy

2.3.1. Smooth Gauged Sets

The unification, as advertised in Fig. 1, of the differential geometry of smooth sets (§ 2.1.1) with the homotopy theory of gauged sets (∞ -groupoids, § 2.2.1) is now immediate:

A *smooth gauged set* (*smooth ∞ -groupoid* or *smooth ∞ -stack*) should be a generalized space whose structure is detected by plotting out probe spaces which are formal products $\mathbb{R}^n \times \Delta^k$ of a Cartesian space (for probing the smooth structure) with a cellular simplex (for probing the higher gauge structure). Such formal products are the objects in the product site $\text{CartSp} \times \Delta$ and hence (the system of plots of) a smooth gauged set \mathcal{X} is encoded by a functor of the form

$$\begin{array}{ccc}
 (\text{CartSp} \times \Delta)^{\text{op}} & \longrightarrow & \text{Set} \\
 \mathbb{R}^n \times \Delta^k & \longmapsto & \text{Plt}(\mathbb{R}^n \times \Delta^k, \mathcal{X}),
 \end{array}
 \tag{60}$$

or equivalently of the form

$$\begin{array}{ccc}
 \text{CartSp}^{\text{op}} & \longrightarrow & \text{SSet} \\
 \mathbb{R}^n & \longmapsto & (\Delta^k \mapsto \text{Plt}(\mathbb{R}^n \times \Delta^k, \mathcal{X})).
 \end{array}
 \tag{61}$$

As such, this is known as a *simplicial presheaf* on the site CartSp . In more detail, the simplicial sets of plots of a smooth gauged set should be Kan complexes (46), hence of the form

$$\text{CartSp}^{\text{op}} \longrightarrow \text{KanSSet} \hookrightarrow \text{SSet}.
 \tag{62}$$

As before, there is redundancy in this description: For detecting the smooth structure, it ought to be sufficient to consider germs of probes, and for properly recognizing the gauged structure we need to regard weak homotopy equivalences of simplicial sets of plots as actual homotopy equivalences. Thus, denoting by

$$W_{\text{lwh}} \subset \text{Mor}(\text{Func}(\text{CartSp}^{\text{op}}, \text{KanSSet}))
 \tag{63}$$

the class of such *local* (germ-wise) *weak homotopy equivalences*, the ∞ -category of *smooth*

gauged sets is the simplicial localization

$$\mathrm{SmthGrpd}_\infty := L^{W_{\mathrm{wh}}} \mathrm{Func}(\mathrm{CartSp}^{\mathrm{op}}, \mathrm{KanSSet}). \quad (64)$$

As a fundamental example, for a surjective submersion $p : U \twoheadrightarrow X$ of smooth manifolds (such as an open cover), its Čech nerve is

$$\begin{aligned} \mathbf{u} &\in \mathrm{SmthGrpd}_\infty \\ \mathrm{Plt}(\mathbb{R}^n \times \Delta^k, \mathbf{u}) &:= C^\infty(\mathbb{R}^n, U^{\times_k X}) \end{aligned} \quad (65)$$

and the comparison map

$$\begin{array}{ccc} \mathbf{u} & \longrightarrow & X \\ C^\infty(\mathbb{R}^n, U^{\times_k X}) & \xrightarrow{p_*} & C^\infty(\mathbb{R}^n, X) \end{array}$$

is a local weak homotopy equivalence.

In evident generalization of (50), we have that smooth sets form a reflective sub- ∞ -category of smooth ∞ -groupoids, which we denote by the same symbols:

$$\mathrm{SmthGrpd}_\infty \begin{array}{c} \xrightarrow{\tau_0} \\ \xleftarrow{\perp} \end{array} \mathrm{SmthSet}. \quad (66)$$

with the corresponding 0-truncation modality, generalizing (51)

$$[-]_0 : \mathrm{SmthGrpd}_\infty \xrightarrow{\tau_0} \mathrm{SmthSet} \hookrightarrow \mathrm{SmthGrpd}_\infty. \quad (67)$$

This is just given by plotwise 0-truncation (“followed by sheafification”, which however we have absorbed in the localization (64)):

$$\mathrm{Plt}(\mathbb{R}^n, [\mathbf{X}]_0) \simeq [\mathrm{Plt}(\mathbb{R}^n, \mathbf{X})]_0. \quad (68)$$

This ∞ -category (64) is an ∞ -topos. In particular, this means that it is *Cartesian closed*, in generalization of (18) and (19), in that for all $\mathbf{X}, \mathbf{Y} \in \mathrm{SmthGrpd}_\infty$ we have the smooth ∞ -groupoid of smooth maps between them (the *mapping stack*)

$$\begin{aligned} \mathrm{Map}(\mathbf{X}, \mathbf{Y}) &\in \mathrm{SmthGrpd}_\infty \\ \mathrm{Plt}(\mathbb{R}^n, \mathrm{Map}(\mathbf{X}, \mathbf{Y})) &= \mathrm{Hom}(\mathbb{R}^n \times \mathbf{X}, \mathbf{Y}), \end{aligned} \quad (69)$$

where now $\mathrm{Hom}(-, -)$ denotes the plain hom ∞ -groupoid between smooth ∞ -groupoids.

But as an ∞ -topos, Grpd_∞ is special: it is *cohesive*. This is what we discuss next.

2.3.2. Cohesion

Above, we motivated the construction of $\mathrm{SmthGrpd}_\infty$ (64) by imagining probing these higher smooth spaces by plotting out little smooth balls inside them. The intuition is that these balls probe how the spatial constituents smoothly *hang together*, hence how there is a smooth “cohesion” among them. This is the *differential geometric* aspect of smooth gauged sets.

Remarkably, beyond motivating the definition, this cohesion is visible in the abstract properties of the ∞ -topos, namely in that it carries an adjoint quadruple of the form (39), but now promoted to ∞ -functors over ∞ -groupoids [Sch13, Prop. 4.4.8]:

$$\begin{array}{ccccc} \times & \xrightarrow{\mathrm{Shp}} & & & \\ & \downarrow & & & \\ & \xleftarrow{\mathrm{Dsc}} & & & \\ \mathrm{SmthGrpd}_\infty & \xrightarrow{\mathrm{Pnt}} & \mathrm{Grpd}_\infty & & \\ & \downarrow & & & \\ & \xleftarrow{\mathrm{Cht}} & & & \end{array} \quad (70)$$

Here, the bottom three functors are just as in (27) in § 2.1.2 and have the analogous interpretation: They witness the underlying (∞ -groupoids of) points and the discrete and chaotic smooth structures on these, generalized straightforwardly from smooth sets to smooth ∞ -groupoids.

The Shape Modality. More profound is the generalization embodied by the further left adjoint functor \int in (70), to be pronounced the *shape* operation. This may be understood as contracting all cohesive blobs of points (the \mathbb{R}^k) to points, while retaining their global intersection structure.

The corresponding composite functor we denote by

$$\int : \text{SmthGrpd}_\infty \xrightarrow{\text{Shp}} \text{Grpd}_\infty \xrightarrow{\text{Dsc}} \text{SmthGrpd}_\infty \quad (71)$$

where “ \int ” is (not the integral symbol but) the *esh* symbol from the phonetic alphabet: The pronunciation of “shape” is /ʃerp/.

We may think of $\int \mathfrak{X}$ as being the *smooth path ∞ -groupoid* of \mathfrak{X} , as follows:

Consider the cosimplicial object Δ of *extended smooth simplices*²

$$\begin{aligned} \Delta &\longrightarrow \text{SmthMfd} \subset \text{SmthGrpd}_\infty \\ \Delta[k] &\longmapsto \Delta^k := \{\vec{x} \in \mathbb{R}^{k+1} \mid \sum_i x^i = 1\} \simeq \mathbb{R}^k \end{aligned} \quad (72)$$

which on injections $\Delta[k] \hookrightarrow \Delta[k+1]$ is given by inserting a zero coordinate, and on surjections $\Delta[k+1] \twoheadrightarrow \Delta[k]$ is given by adding consecutive coordinates. Then [BBP24]:

$$\int \mathfrak{X} \simeq \varinjlim_k \mathbf{Map}(\Delta^k, \mathfrak{X}) \simeq \text{Dsc}(\varinjlim_k \mathfrak{X}(\Delta^k)). \quad (73)$$

For instance, when $\mathfrak{X} \equiv \mathbf{X}$ is 0-truncated (a smooth set), then the homotopy colimit in (73) is equivalently the *smooth singular simplicial set*

$$\mathbf{x} \in \text{SmthSet} \quad \vdash \quad \begin{cases} \text{Sing}(\mathbf{X}) \in \text{SSet} \rightarrow \text{Grpd}_\infty \\ \text{Plt}(\Delta[k], \text{Sing}(\mathbf{X})) = \text{Hom}(\Delta^k, \mathbf{X}) \end{cases} \quad (74)$$

in that

$$\mathbf{x} \in \text{SmthSet} \quad \Rightarrow \quad \int \mathbf{X} \simeq \text{Dsc} \text{Sing}(\mathbf{X}). \quad (75)$$

This way, one may generally understand $\int(-)$ as a generalization of the traditional $\text{Sing}(-)$ to smooth ∞ -groupoids (cf. [SS26d, pp. 155]). Moreover, by adjunction, there is a natural transformation (the *shape unit*)

$$\mathfrak{X} \xrightarrow{\eta_{\mathfrak{X}}^{\int}} \int \mathfrak{X} \quad (76)$$

which, under the equivalence (73), is the inclusion of constant paths. Under the shape operation, the shape unit is an equivalence:

$$\int \mathfrak{X} \xrightarrow[\sim]{\int \eta_{\mathfrak{X}}^{\int}} \int \int \mathfrak{X}. \quad (77)$$

Further important properties of the shape operation are:

²The usual bounded (smooth) simplices are as in (72) except for the additional condition $\forall_i : x^i > 0$, which makes them manifolds with boundary. The extended simplices (72) are a “collared” version of these ordinary simplices, which improves their technical properties without changing their homotopy theoretic content.

(i) **(Partial exactness)**. Shape preserves homotopy fiber products (58) at least:

(a) over geometrically discrete objects [SS26d, Prop. 4.3.8; Sch13, Thm. 3.8.19]:

$$\left. \begin{array}{l} \mathcal{X}, \mathcal{Y} \in \text{SmothGrpd}_\infty \\ \mathcal{B} \in \text{Dsc}(\text{Grpd}_\infty) \end{array} \right\} \Rightarrow \int(\mathcal{X} \times_{\mathcal{B}} \mathcal{Y}) \simeq (\int \mathcal{X}) \times_{\mathcal{B}} (\int \mathcal{Y}), \quad (78)$$

(b) of 0-truncated objects if one of the maps becomes a Kan fibration under the smooth singular simplicial complex Sing (74):³

$$\left. \begin{array}{l} \mathbf{X} \begin{array}{c} \xrightarrow{f} \\ \searrow \\ \mathbf{B} \end{array} \begin{array}{c} \xleftarrow{g} \\ \swarrow \\ \mathbf{Y} \end{array} \in \text{SmothSet} \\ \text{Sing}(f) \text{ is Kan fibration} \end{array} \right\} \Rightarrow \int(\mathbf{X} \times_{\mathbf{B}} \mathbf{Y}) \simeq (\int \mathbf{X}) \times_{\int \mathbf{B}} (\int \mathbf{Y}). \quad (79)$$

(ii) **(Smooth Oka principle)**. Shape preserves mappings out of smooth manifolds [BBP24, Prop. 1.4; SS26d, Thm. 4.3.54]:

$$\left. \begin{array}{l} X \in \text{SmothMfd} \\ \mathcal{Y} \in \text{SmothGrpd}_\infty \end{array} \right\} \Rightarrow \int \mathbf{Map}(X, \mathcal{Y}) \simeq \mathbf{Map}(X, \int \mathcal{Y}) \simeq \mathbf{Map}(\int X, \int \mathcal{Y}). \quad (80)$$

The Flat Modality. The understanding of the flat modality (23) on gauged smooth sets

$$\begin{array}{ccc} & \overbrace{\hspace{10em}}^{\flat} & \\ \text{SmothSet} & \xrightarrow{\text{Pnt}} \text{Set} & \xleftarrow{\text{Dsc}} \text{SmothSet} \\ & \underbrace{\hspace{10em}}_{\flat} & \end{array} \quad (81)$$

is more straightforward. We highlight the following immediate but important point: By (68), the flat modality commutes with the 0-truncation modality (67):

$$[\flat \mathcal{X}]_0 \simeq \flat[\mathcal{X}]_0. \quad (82)$$

3. Higher Maxwell-type Equations of Motion

We make precise the notion of higher Maxwell-type equations of motion (EoM) and how they are equivalently solved by closed differential forms with coefficients in a characteristic L_∞ -algebra.

3.1. Flux Densities

3.1.1. Differential Form Moduli

Universal de Rham Complex. The smooth moduli sets of differential n -forms $\Omega_{\text{dR}}^n(*) \in \text{SmothSet}$ (16) canonically carry the further structure of smooth \mathbb{R} -vector spaces (by plotwise addition and multiplication):

$$\Omega_{\text{dR}}^n(*) \in \text{Vec}_{\mathbb{R}}(\text{SmothSet}). \quad (83)$$

Moreover, equipped with the universal wedge products (given plot-wise by the ordinary wedge product):

$$\Omega_{\text{dR}}^{n_1}(*) \times \Omega_{\text{dR}}^{n_2}(*) \xrightarrow{(-) \wedge (-)} \Omega_{\text{dR}}^{n_1+n_2}(*) \quad (84)$$

³The implication (79) follows because Sing (74) evidently preserves 1-categorical fiber products and since (cf. [Lur09, Prop. A.2.4.4] with [GJ09, §II.8.6]) in SSet these present homotopy fiber products as soon as one of the maps is a Kan fibration.

and universal de Rham differential (again plotwise given by the ordinary de Rham differential),

$$\Omega_{\text{dR}}^n(*) \xrightarrow{\text{d}} \Omega_{\text{dR}}^{n+1}(*), \quad (85)$$

these smooth vector spaces form the *universal de Rham complex*:

$$\Omega_{\text{dR}}^\bullet(*) \in \text{dgcAlg}_{\mathbb{R}}(\text{SmthSet}). \quad (86)$$

De Rham moduli over a Smooth Set. More generally, given $\mathbf{X} \in \text{SmthSet}$, we say that its plain *de Rham complex* is

$$\begin{aligned} \Omega_{\text{dR}}^\bullet(\mathbf{X}) &:= \text{bMap}(\mathbf{X}, \Omega_{\text{dR}}^\bullet(*)) \\ &= \text{Hom}(\mathbf{X}, \Omega_{\text{dR}}^\bullet(*)) \in \text{dgcAlg}_{\mathbb{R}}(\text{Set}), \end{aligned} \quad (87)$$

and its *smooth de Rham complex* is

$$\Omega_{\text{dR}}^\bullet(\mathbf{X}) := \text{Map}(\mathbf{X}, \Omega_{\text{dR}}^\bullet(*)) \in \text{dgcAlg}_{\mathbb{R}}(\text{SmthSet}). \quad (88)$$

We note that, by the formula (18) for $\text{Map}(-, -)$, we have:

$$\begin{aligned} \text{Plt}(\mathbb{R}^n, \Omega_{\text{dR}}^\bullet(\mathbf{X})) &\equiv \text{Plt}(\mathbb{R}^n, \Omega_{\text{dR}}^\bullet(\mathbf{X})) \\ &\equiv \text{Plt}(\mathbb{R}^n, \text{Map}(\mathbf{X}, \Omega_{\text{dR}}^\bullet(*))) \\ &= \text{Hom}(\mathbb{R}^n \times \mathbf{X}, \Omega_{\text{dR}}^\bullet(*)) \\ &= \Omega_{\text{dR}}^\bullet(\mathbb{R}^n \times \mathbf{X}). \end{aligned} \quad (89)$$

Concrete Smooth de Rham complex. The smooth de Rham complex (88) is far from concrete (33). It follows from (89) that its *concretification* (35) has as \mathbb{R}^n -plots the vertical differential forms relative to \mathbb{R}^n , hence the smoothly \mathbb{R}^n -parameterized differential forms on \mathbf{X} ([Sch13, Prop. 1.266]):

$$\begin{aligned} \sharp_1 \Omega_{\text{dR}}^\bullet(\mathbf{X}) &\in \text{SmthSet} \\ \text{Plt}(\mathbb{R}^n, \sharp_1 \Omega_{\text{dR}}^\bullet(\mathbf{X})) &\simeq \Omega_{\text{vert}/\mathbb{R}^n}^\bullet(\mathbb{R}^n \times \mathbf{X}) \\ &\simeq C^\infty(\mathbb{R}^n, \Omega_{\text{dR}}^\bullet(\mathbf{X})). \end{aligned} \quad (90)$$

Compactly Supported de Rham complex. For $X \in \text{SmthMfd}$ write $\text{Cmp}(X)$ for the category whose objects are compact submanifolds with boundary and whose morphisms are inclusions. The smooth de Rham complex of differential forms on X which are *compactly supported* or *vanishing at infinity* is the colimit over such K of the moduli of differential forms whose pullback ι_K^* to the complement $X - K^\circ$ (of the interior of K) vanishes:

$$\begin{aligned} \Omega_{\text{dR},c}^\bullet(X) &:= \varinjlim_{K \in \text{Cmp}(X)} \Omega_{\text{dR}}^\bullet(X, X - K^\circ) \\ &:= \varinjlim_{K \in \text{Cmp}(X)} \text{fib}\left(\Omega_{\text{dR}}^\bullet(X) \xrightarrow{\iota_K^*} \Omega_{\text{dR}}^\bullet(X - K^\circ)\right). \end{aligned} \quad (91)$$

3.1.2. Flux Density Moduli

Consider a finite set I and an I -indexed set of degrees:

$$\text{deg}_{(-)} : I \longrightarrow \mathbb{N}. \quad (92)$$

Given $i \in I$, write

$$b^{(i)} \in \Omega_{\text{dR}}^{\text{deg}_i}(\Omega_{\text{dR}}^{\text{deg}_i}(*)) \hookrightarrow \Omega_{\text{dR}}^{\bullet}(\Omega_{\text{dR}}^{\text{deg}_i}(*)) \quad (93)$$

for the universal n -form, the one given by the identity map

$$b^{(i)} \equiv \text{id}_{\Omega_{\text{dR}}^{\text{deg}_i}(*)}. \quad (94)$$

Via pullback along the product projection maps

$$\prod_j \Omega_{\text{dR}}^{\text{deg}_j}(*). \xrightarrow{\text{pr}_i} \Omega_{\text{dR}}^{\text{deg}_i}(*), \quad (95)$$

these induce differential forms

$$\text{pr}_i^* b^{(i)} \in \prod_j \Omega_{\text{dR}}^{\text{deg}_j}(*). \quad (96)$$

on their joint product moduli set. This way, for $\mathbf{X} \in \text{SmthSet}$, a map

$$\mathbf{X} \xrightarrow{\vec{B}} \prod_i \Omega_{\text{dR}}^{\text{deg}_i}(*). \quad (97)$$

is equivalently an $i \in I$ -indexed set of deg_i -forms on \mathbf{X} :

$$\vec{B} = (B^{(i)} := \vec{B}^* \text{pr}_i^* b^{(i)})_{i \in I} \in \prod_i \Omega_{\text{dR}}^{\text{deg}_i}(\mathbf{X}). \quad (98)$$

In this manner,

$$\prod_j \Omega_{\text{dR}}^{\text{deg}_j}(*). \in \text{SmthSet} \quad (99)$$

is the *smooth moduli set* for *flux densities* in degrees deg_{\bullet} on \mathbf{X} .

Next, we impose equations of motion on such flux densities

3.2. Equations of Motion

3.2.1. Higher Maxwell-type EoM

Let $X^{1,d}$ be a Lorentzian spacetime manifold, hence in particular a smooth manifold equipped with a pseudo-Riemannian metric with respect to which we have the Hodge star operator

$$\star : \Omega_{\text{dR}}^{\bullet}(X^{1,d}) \xrightarrow{\sim} \Omega_{\text{dR}}^{1+d-\bullet}(X^{1,d}). \quad (100)$$

For sets I of flux densities in degrees deg_i (92)

$$\vec{F} := \{F^{(i)} \in \Omega_{\text{dR}}^{\text{deg}_i}(X)\}, \quad (101)$$

we say [SS24b, §2.1] that *higher Maxwell-type equations of motion* are systems of equations of the form

$$\forall_i : dF^{(i)} = P^{(i)}(\vec{F}), \quad (102a)$$

$$\forall_i : \star F^{(i)} = \mu^{(i)}(\vec{F}), \quad (102b)$$

where:

- (i) the $P^{(i)}$ are graded-symmetric polynomials in I variables of degrees deg_i ,
- (ii) μ is a linear automorphism on the linear span of I .

Here, the first equations (102a) are called the *higher Bianchi identities* and the second equations (102b) are the *higher duality relations*.

Basic examples are the following (where we indicate the degrees of differential forms in their subscripts):

4D Vacuum Maxwell (gauge sector of 4D SuGra)	$\begin{aligned} dF_2 &= 0 \\ dG_2 &= 0, \end{aligned}$	$\star F_2 = G_2, \quad (103a)$
---	---	---------------------------------

Chiral Flux in $D = 4k+2$	$dF_{2k+1} = 0,$	$\star F_{2k+1} = F_{2k+1}. \quad (103b)$
---------------------------	------------------	---

5D Maxwell-Chern-Simons (gauge sector of 5D SuGra)	$\begin{aligned} dF_2 &= 0 \\ dG_3 &= \frac{1}{2}F_2 \wedge F_2, \end{aligned}$	$\star F_2 = G_3, \quad (103c)$
---	---	---------------------------------

10D massive IIA NS/RR-Fluxes (gauge sector in massive 10D SuGra)	$\begin{aligned} dF_{2\bullet} &= H_3 \wedge F_{2\bullet-2} \\ dH_3 &= 0 \\ dH_7 &= \frac{1}{2}\sum_i (-1)^i F_{2i} \wedge F_{8-2i}, \end{aligned}$	$\begin{aligned} \star F_{2\bullet} &= F_{10-2\bullet} \\ \star H_3 &= H_7, \end{aligned} \quad (103d)$
---	---	---

10D actual IIA NS/RR-Fluxes (gauge sector of 10D IIA SuGra)	$\begin{aligned} dF_2 &= 0 \\ dF_4 &= H_3 \wedge F_2 \\ dF_6 &= H_3 \wedge F_4 \\ dH_3 &= 0 \\ dH_7 &= \frac{1}{2}F_4 \wedge F_4 - F_2 \wedge F_6, \end{aligned}$	$\begin{aligned} \star F_{2\bullet} &= F_{10-2\bullet} \\ \star H_3 &= H_7, \end{aligned} \quad (103e)$
--	---	---

11D Maxwell-Chern-Simons (gauge sector of 11D SuGra)	$\begin{aligned} dG_4 &= 0 \\ dG_7 &= \frac{1}{2}G_4 \wedge G_4, \end{aligned}$	$\star G_4 = G_7. \quad (103f)$
---	---	---------------------------------

Not an example are the equations of motion of nonabelian Yang-Mills theory with non-trivial structure constants (f_{jk}^i) , since their Bianchi identities

$$dF^i = \sum_{j,k} f_{jk}^i A^j \wedge F^k \quad (104)$$

involve *gauge potential* forms A^i that are not in general globally defined as required in (101).

Due to this restriction, we speak of *Maxwell-type* (hence: abelian Yang-Mills type) equations of motion. But beware that, even so, the nonlinear Bianchi identities that we admit, through the polynomials $P^{(i)}$ in (102a), make such Maxwell-type fluxes in general have quantization laws in *nonabelian cohomology* — this is the topic of § 4.

Also *not* immediately an example of (102) are Chern-Simons type equations of motion that require flux densities to vanish, $F^{(i)} = 0$. For such gauge fields, their entire non-trivial content is in the flux-quantized gauge potentials, which we come to in § 4. But this case is subsumed by a natural generalization which we discuss in § 5.2.

Initial Cauchy Data. Now consider the special case that spacetime is globally hyperbolic, $X^{1,d} \simeq \mathbb{R}^{1,0} \times X^d$, and consider the set of germs of solutions to such higher Maxwell-type EoM (102) around the spatial Cauchy surface

$$\{0\} \times X^d \xrightarrow{\iota} X^{1,d}, \quad (105)$$

hence:

$$\text{Sol}(X^d) := \left\{ \epsilon \in \mathbb{R}_+, \left(F^{(i)} \in \Omega_{\text{dR}}^{\text{deg}_i}((-\epsilon, +\epsilon) \times X^d) \right)_{i \in I} \left| \begin{array}{l} d\vec{F} = \vec{P}(\vec{F}) \\ \star \vec{F} = \vec{\mu}(\vec{F}) \end{array} \right. \right\} / \sim, \quad (106)$$

where $(\epsilon, \vec{F}) \sim (\epsilon', \vec{F}')$ iff there exists $0 < \epsilon'' < \epsilon, \epsilon'$ such that $\vec{F} = \vec{F}'$ after joint restriction to $(-\epsilon'', +\epsilon'') \times X^d$.

Observing that Hodge duality exchanges purely spatial forms with those having a temporal dt -factor, one finds [SS24b, Thm 2.2 & §A] that these germs of solutions are equivalently encoded in Cauchy data on X^d , which has the same form as (101) and (102) except that the Hodge duality relation disappears:

$$\text{Sol}(X^d) \xrightarrow[\sim]{\iota^*} \left\{ B^{(i)} \in \Omega_{\text{dR}}^{\text{deg}_i}(X^d) \mid d\vec{B} = \vec{P}(\vec{B}) \right\}. \quad (107)$$

The conditions on the right of (107) are the higher *Gauss laws*, being the spatial components of the higher Bianchi identities (102a), while the Hodge duality relation (102b) is absorbed into witnessing the existence of an inverse to (107).

3.2.2. Characteristic L_∞ -Algebra

We proceed to further identify this Cauchy data (107) with closed differential forms with coefficients in a characteristic L_∞ -algebra (§ 3.2.2). For this purpose we now require (in § 3.2.2) that the higher Maxwell-type EoM are suitably universal (which is the case for all the examples (103) of interest, but deserves a formalization).

Universal Exterior Differential Systems. Recall that on $X \in \text{SmthMfd}$, an *exterior differential system* (cf. [Bry+91; McK19]) is a dgc-ideal in the de Rham complex $\mathcal{G} \subset \Omega_{\text{dR}}^\bullet(X)$. Its *integral solutions* are submanifold embeddings along which this ideal pulls back to zero, forming the solution set

$$\text{Sol}(X, \mathcal{G}) := \left\{ Y \xrightarrow{\phi} X \mid \phi^*\mathcal{G} = 0 \right\}. \quad (108)$$

This is a way of encoding (solutions to) differential equations with variables in X .

Our goal is to formulate *universal* such differential equations which do not depend on the choice of domain X but only on an abstract set $i \in I$ of deg_i -forms (92) on any domain. But with the universal de Rham complex (§ 3.1.1) in hand, this is immediate: We take such *universal exterior differential systems* to be dgc-ideals \mathcal{G} in the de Rham complex (87) on the smooth moduli set (97) of deg_i -forms, $i \in I$:

$$\mathcal{G} \subset \Omega_{\text{dR}}^\bullet\left(\prod_i \Omega_{\text{dR}}^{\text{deg}_i}(*)\right), \quad (109)$$

and say that the solution set on $\mathbf{X} \in \text{SmthSet}$ is

$$\text{Sol}(\mathbf{X}, \mathcal{G}) := \left\{ \mathbf{X} \xrightarrow{\phi} \prod_i \Omega_{\text{dR}}^{\text{deg}_i}(*)\right\} \mid \phi^*\mathcal{G} = 0 \right\}. \quad (110)$$

Specifically, we say that:

- (i) *Higher Gauss laws* in degrees deg_i , $i \in I$ (92) are universal exterior differential systems (109) which are *algebraically* generated as

$$\mathcal{G} = \left\langle d \text{pr}_i^* b^{(i)} - P^{(i)}\left(\left(\text{pr}_j^* b^{(j)}\right)_{j \in I}\right) \right\rangle_{i \in I} \subset \Omega_{\text{dR}}^\bullet\left(\prod_i \Omega_{\text{dR}}^{\text{deg}_i}(*)\right), \quad (111)$$

for graded-symmetric (wedge product) polynomials $P^{(i)}$ in the set I of deg-graded variables, evaluated here on the universal differential forms (96).

(ii) Flux densities \vec{B} (97) on \mathbf{X} solve the Gauss law (111) if they are in the corresponding solution set (110), hence if

$$\vec{B}^*\mathcal{J} = 0 \in \Omega_{\text{dR}}^{\bullet}(\mathbf{X}), \quad (112)$$

which means equivalently that

$$\forall_i : d B^{(i)} = P^{(i)}(\{B^{(j)}\}_{j \in I}). \quad (113)$$

Characteristic L_{∞} -Algebra. For given degrees (92) such that all $\text{deg}_i \geq 1$, let

$$\wedge^1 \mathfrak{a}^{\vee} := \langle \beta^{(i)} \rangle \in \text{Vec}_{\mathbb{R}}^{\mathbb{N}} \quad (114)$$

denote the \mathbb{N} -graded vector space spanned by elements $\beta^{(i)}$ in degree deg_i , and denote by

$$\wedge^{\bullet} \mathfrak{a}^{\vee} := \text{Sym}(\wedge^1 \mathfrak{a}^{\vee}) \quad (115)$$

its graded-symmetric algebra, cf. (115).

Lemma 3.1. *Given a pregeometric higher Maxwell-type EoM (111), in degrees deg (92), a differential d (graded derivation of degree +1 that squares to 0) on $\wedge^{\bullet} \mathfrak{a}^{\vee}$ (115) is given on generators by the formulas*

$$\forall_i : d \beta^{(i)} = P^{(i)}(\{\beta^{(j)}\}_{j \in I}), \quad (116)$$

in that it follows that $d^2 = 0$.

Proof. By the graded derivation property of d , we are equivalently claiming that the expressions

$$d(P^{(i)}(\{\beta^{(j)}\}_{j \in I})) = \sum_{k \in I} P^{(k)} \frac{\partial}{\partial \beta^{(k)}} P^{(i)}(\{\beta^{(j)}\}_{j \in I}) \quad (117)$$

vanish for all $i \in I$ (where on the right we have the sum over graded partial derivatives). Since the $\beta^{(j)}$ are free generators, this is the case iff the polynomials on the right vanish identically as functions (for all possible arguments), hence iff

$$\forall_i \quad \sum_{k \in I} P^{(k)} \frac{\partial}{\partial \beta^{(k)}} P^{(i)} = 0. \quad (118)$$

This is to be shown.

Now, the fact that \mathcal{J} is a differential ideal (111) implies analogously that

$$\forall_i \quad d(P^{(i)}(\{\text{pr}_j^* b^{(j)}\}_{j \in I})) = \sum_{k \in I} P^{(k)} \frac{\partial}{\partial \text{pr}_k^* b^{(k)}} P^{(i)}(\{\text{pr}_j^* b^{(j)}\}_{j \in I}) = 0. \quad (119)$$

This being an expression in $\Omega_{\text{dR}}^{\text{deg}_i+1}(\prod_j \Omega_{\text{dR}}^{\text{deg}_j}(*))$, its vanishing implies that for all $U \in \text{SmthMfd}$ and all $(B^{(j)} \in \Omega_{\text{dR}}^{\text{deg}_j}(U))_{j \in I}$ we have

$$\sum_{k \in I} P^{(k)} \frac{\partial}{\partial B^{(k)}} P^{(i)}(\{B^{(j)}\}_{j \in I}) = 0. \quad (120)$$

In particular, we may take $U := \mathbb{R}^N$ with N so large that there exist constant differential forms $B_{\text{free}}^{(j)} \in \Omega_{\text{dR}}^{\text{deg}_j}(U)$, $j \in I$ (sums of wedge products of differentials of the canonical coordinate functions), with the property that all their nontrivial wedge monomials of total degree less than or equal to $\text{deg}_i + 1$ are non-vanishing except for reasons of graded-commutativity. Hence specialized to these, (120) implies (118), thereby the vanishing of (117) and thus the claim (116). \square

With the terminology recalled in § A.1, we have:

Definition 3.2. The dgc-algebra implied by Lem. 3.1, to be denoted

$$\text{CE}(\mathfrak{a}) := (\wedge^\bullet \mathfrak{a}^\vee, d) \in \text{dgcAlg}_{\mathbb{R}}(\text{Set}), \quad (121)$$

is the Chevalley-Eilenberg algebra of an L_∞ -algebra

$$\mathfrak{a} \in L_\infty \text{Alg}_{\mathbb{R}}(\text{Set}), \quad (122)$$

which we call the *characteristic L_∞ -algebra* of the given pregeometric higher Maxwell-type EoMs.

The grand conclusion of this section is thus:

Theorem 3.3 ([SS24b, §2.1]). *The solution set (106) of higher Maxwell-type equations (102) that are universal in the above sense (111), is isomorphic to the set of closed differential forms (250) on the Cauchy surface with coefficients in the characteristic L_∞ -algebra (122):*

$$\text{Sol} \xrightarrow[\text{(107)}]{\sim} \left\{ B^{(i)} \in \Omega_{\text{dR}}^{\text{deg}_i}(X^d) \mid d\vec{B} = \vec{P}(\vec{B}) \right\} \xrightarrow{\sim} \Omega_{\text{cl}}^1(X^d; \mathfrak{a}). \quad (123)$$

Remark 3.4 (Two notions of higher gauge theory). Beware (cf. Fig. 2) that the role of L_∞ -algebras here is different from the “higher principal connection” approach to higher gauge theory ([Sch05, §11-12; BS07; FSS12; SSS12; SW13; Bor+25; Alf25, §2; RSW26; Wal26; Bun+26]): There L_∞ -algebras serve as the coefficients of gauge potential forms which are generically not flat/closed — while here they serve as coefficients of the flux densities and their closure is no less than the electromagnetic Gauss law constraint putting the theory on shell.

Figure 2: There are two approaches to globally modeling “higher gauge fields”: Either as higher generalization of connections on principal/fiber bundles, or as nonabelian generalization of ordinary differential cohomology. Here we are concerned with the latter (from [FSS23, §9]).

	Principal	Cohomological
Specified L_∞ -algebra	coefficient of gauge potentials (connections)	coefficient of flux densities (curvatures)
Flatness / closure	non-generic	Gauss law / Bianchi identity
Examples	heterotic SuGra / B-field on T-folds	type I/II SuGra / 11D SuGra

But with the electromagnetic Gauss law constraint on initial Cauchy data of flux densities understood as an L_∞ -closure condition, by Thm. 3.3, we have opened the door to understanding the “quantization” (discretization) of flux densities, in generality. This is the topic of § 4.1.

4. Global Phase Space of Higher Gauge Fields

4.1. Flux Quantization

We have characterized (Thm. 3.3) the solution space of higher flux densities. This generalizes the historical situation of [Max65], when Maxwell’s equations for electromagnetic flux were

formulated. Later, [Dir31] noticed that the electromagnetic *gauge potential* exhibits a global “quantization” (discretization) of the admissible total flux, and our task now is to discuss the corresponding generalization of that *flux quantization* via *gauge potentials*.

4.1.1. Rational Homotopy

Smooth Moduli Set of On-Shell Fluxes. In Thm. 3.3 we identified the solution set of Maxwell-type equations around a Cauchy surface X^d as the set of closed \mathfrak{a} -valued differential forms for a characteristic L_∞ -algebra \mathfrak{a} :

$$\text{Sol} \simeq \Omega_{\text{cl}}^1(X^d; \mathfrak{a}). \quad (124)$$

This means that there is actually a *smooth moduli set* of solutions/closed forms, inside the smooth de Rham complex (88) from § 3.1.1:

$$\begin{aligned} \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) &\in \text{SmthSet} \\ \text{Plt}(\mathbb{R}^n, \Omega_{\text{cl}}^1(X^d; \mathfrak{a})) &:= \Omega_{\text{dR}}^1(\mathbb{R}^n \times X^d; \mathfrak{a}), \end{aligned} \quad (125)$$

hence, by (18), with:

$$\Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \simeq \mathbf{Map}(X^d, \Omega_{\text{dR}}^1(*)). \quad (126)$$

Shape of On-Shell Flux Moduli. The shape (71) of this smooth moduli set is, by (75), (80) and (126):

$$\begin{aligned} \int \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) &\simeq \int \mathbf{Map}(X^d, \Omega_{\text{cl}}^1(*, \mathfrak{a})) \\ &\simeq \mathbf{Map}(\int X^d, \int \Omega_{\text{cl}}^1(*, \mathfrak{a})) \\ &\simeq \text{Dsc Map}(\text{Sing } X^d, \text{Sing } \Omega_{\text{cl}}^1(*, \mathfrak{a})), \end{aligned} \quad (127)$$

with

$$\begin{aligned} \text{Sing}(\Omega_{\text{cl}}^1(*)) &\in \text{SSet} \rightarrow \text{Grpd}_\infty \\ \text{Plt}(\Delta[k], \text{Sing}(\Omega_{\text{cl}}^1(*))) &= \Omega_{\text{cl}}^1(\Delta^k; \mathfrak{a}). \end{aligned} \quad (128)$$

Rationalizing Classifying Spaces. This *simplicial set of closed \mathfrak{a} -valued differential forms* is a famous construction in *rational homotopy theory* (RHT, for which cf. [FHT00; Hes07; FSS23, §5]):

For a simply connected $\mathcal{A} \in \text{Grpd}_\infty$, its \mathbb{R} -*rationalization* is $L^{\mathbb{R}}\mathcal{A} \in \text{Grpd}$ whose homotopy groups are the \mathbb{R} -rationalization of those of \mathcal{A} :

$$\forall_{n \geq 2} : \pi_n(L^{\mathbb{R}}\mathcal{A}) \simeq \pi_n(\mathcal{A}) \otimes_{\mathbb{Z}} \mathbb{R} \quad (129)$$

and which is equipped with a map

$$\mathcal{A} \xrightarrow{\eta_{\mathcal{A}}^{\mathbb{R}}} L^{\mathbb{R}}\mathcal{A} \quad (130)$$

that induces the canonical homomorphisms,

$$\pi_n(\mathcal{A}) \xrightarrow{(\eta_{\mathcal{A}}^{\mathbb{R}})_*} \pi_n(\mathcal{A}) \otimes_{\mathbb{Z}} \mathbb{R} \quad (131)$$

and is suitably universal with that property.

The *fundamental theorem of dg-algebraic RHT* says that for a simply connected \mathcal{A} whose rational cohomology/homotopy groups are finite-dimensional, there exists an essentially

unique minimal L_∞ -algebra

$$\begin{aligned} \mathfrak{L}\mathcal{A} &\in L_\infty\text{Alg}_{\mathbb{R}} \\ (\mathfrak{L}\mathcal{A})_\bullet &\simeq \pi_\bullet(\Omega\mathcal{A}) \otimes_{\mathbb{Z}} \mathbb{R} \end{aligned} \quad (132)$$

whose brackets are the \mathbb{R} -rationalized *higher Whitehead brackets* of \mathcal{A} (and whose CE-algebra is called *minimal Sullivan model* of \mathcal{A}) such that: ⁴

$$L^{\mathbb{R}}\mathcal{A} \simeq \text{Sing } \Omega_{\text{cl}}^1(*; \mathfrak{L}\mathcal{A}). \quad (133)$$

Regarding this under the tacit faithful embedding $\text{Dsc} : \text{Grpd}_\infty \hookrightarrow \text{SmthGrpd}_\infty$, it reads equivalently:

$$L^{\mathbb{R}}\mathcal{A} \simeq \int \Omega_{\text{cl}}^1(*; \mathfrak{L}\mathcal{A}). \quad (134)$$

In this form it immediately generalizes, by the smooth Oka principle (80) and using (126), to: ⁵

$$\{X \in \text{SmthMfd}\} \Rightarrow \text{Map}(X, L^{\mathbb{R}}\mathcal{A}) \simeq \int \Omega_{\text{cl}}^1(X; \mathfrak{L}\mathcal{A}). \quad (135)$$

Restriction to Solitonic Flux Densities. An important variant of (135) is Prop. 4.1 below. The *solitonic* on-shell flux densities are those on-shell flux densities (125) which on a Cauchy surface X are *compactly supported* or *vanishing at infinity* according to (91):

$$\Omega_{\text{cl},c}^1(X; \mathfrak{a}) := \varinjlim_{K \in \text{Cmp}(X)} \text{fib}\left(\Omega_{\text{cl}}^1(X; \mathfrak{a}) \xrightarrow{\iota_K^*} \Omega_{\text{cl}}^1(X - K^\circ; \mathfrak{a})\right). \quad (136)$$

We assume now that the Cauchy surface manifold X is non-pathologic at infinity. One way of saying this is that it is *finitary* in that it is the interior of a compact manifold with boundary.

In this case the compactly supported maps out of X with coefficients in a pointed space \mathcal{Y} ,

$$\text{Map}_c(X, \mathcal{Y}) := \varinjlim_{K \in \text{Cpt}(X)} \text{fib}\left(\text{Map}(X, \mathcal{Y}) \xrightarrow{\iota_K^*} \text{Map}(X - K^\circ, \mathcal{Y})\right), \quad (137)$$

are equivalently (cf. [AF25, p. 6]) the *pointed* maps out of the *one-point compactification* $X_{\cup\{\infty\}}$ (cf. [Jam84, §3; SS25d, §2.2]):

$$\left. \begin{array}{l} \text{finitary } X \in \text{SmthMfd}, \\ \mathcal{Y} \in \text{Grpd}_\infty^* \end{array} \right\} \Rightarrow \text{Map}_c(X, \mathcal{Y}) \xrightarrow{\sim} \text{Map}^*(X_{\cup\{\infty\}}, \mathcal{Y}). \quad (138)$$

Proposition 4.1. *The shape of the moduli of solitonic on-shell flux densities (136) on a finitary Cauchy surface X is the pointed mapping space: ⁶*

$$\text{Map}^*(X, L^{\mathbb{R}}\mathcal{A}) \simeq \int \Omega_{\text{cl},c}^1(X; \mathfrak{L}\mathcal{A}). \quad (139)$$

Proof. We have the following sequence of natural equivalences:

$$\int \Omega_{\text{cl},c}^1(X^d; \mathfrak{L}\mathcal{A})$$

⁴The equivalence (133) is traditionally discussed for polynomial differential forms on bounded simplices; but it holds just as well for the smooth differential forms on extended simplices used here, since these still satisfy (cf. [GM13, Cor. 9.9]) the relevant *extension lemma* (cf. [FSS23, Prop. 5.10; nLa26a]).

⁵Our one-line proof, via cohesion, of (135) reproduces the result of [Ber15, Thm. 1.4] (cf. also [Laz13, Thm. 8.1; BFM11, Thm. 3.1]) for the special case when the domain X is a smooth manifold and for rationalization over the real numbers.

⁶Our quick proof, via cohesion, of (139) reproduces essentially the result of [BFM11, Thm. 1.2] for the special case that the domain X is a smooth manifold and for rationalization over the real numbers.

$$\begin{aligned}
&\equiv \int \varinjlim_K \text{fib} \left(\Omega_{\text{cl}}^\bullet(X; \mathfrak{L}\mathcal{A}) \xrightarrow{\iota_K^*} \Omega_{\text{cl}}^\bullet(X - K^\circ; \mathfrak{L}\mathcal{A}) \right) && \text{by (136)} \\
&\simeq \varinjlim_K \int \text{fib} \left(\Omega_{\text{cl}}^\bullet(X; \mathfrak{L}\mathcal{A}) \xrightarrow{\iota_K^*} \Omega_{\text{cl}}^\bullet(X - K^\circ; \mathfrak{L}\mathcal{A}) \right) && \text{since } \int \text{ preserves colimits} \\
&\simeq \varinjlim_K \text{fib} \left(\int \Omega_{\text{cl}}^\bullet(X; \mathfrak{L}\mathcal{A}) \xrightarrow{\iota_K^*} \int \Omega_{\text{cl}}^\bullet(X - K^\circ; \mathfrak{L}\mathcal{A}) \right) && \text{using (79)} \\
&\simeq \varinjlim_K \text{fib} \left(\mathbf{Map}(X^d, L^{\mathbb{R}}\mathcal{A}) \xrightarrow{\iota_K^*} \mathbf{Map}(X^d - K^\circ, L^{\mathbb{R}}\mathcal{A}) \right) && \text{by (135)} \\
&\simeq \mathbf{Map}^*(X^d, L^{\mathbb{R}}\mathcal{A}) && \text{by (138).} \quad \square
\end{aligned}$$

Admissible Classifying Spaces. We hence ask for classifying spaces \mathcal{A} whose real Whitehead L_∞ -algebra $\mathfrak{L}\mathcal{A}$ (132) coincides with the characteristic L_∞ -algebra \mathfrak{a} (Def. 3.2) of a given Maxwell-type higher gauge theory. For example:

- (i) $\mathfrak{I}B^2\mathbb{Z}^2$, cf. (257), is the characteristic L_∞ -algebra of 4D vacuum electromagnetism (103a):

$$\text{CE}(\mathfrak{I}B^2\mathbb{Z}^2) \simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ g_2 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d g_2 = 0 \end{array} \right). \quad (140)$$

- (ii) $\mathfrak{I}B^{2k+1}\mathbb{Z}$, cf. (257), is the characteristic L_∞ -algebra of the chiral flux in $D = 4k + 2$ (103b):

$$\text{CE}(\mathfrak{I}B^{2k+1}\mathbb{Z}) \simeq \mathbb{R}_d [c_{2k+1}] / (d c_{2k+1} = 0). \quad (141)$$

- (iii) $\mathfrak{I}S^2$, cf. (269), is the characteristic L_∞ -algebra of 5D Maxwell-Chern-Simons theory (103c):

$$\text{CE}(\mathfrak{I}S^2) \simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ g_3 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d g_3 = \frac{1}{2} f_2^2 \end{array} \right). \quad (142)$$

- (iv) $\mathfrak{I}(\text{KU}_0 // \text{BU}(1))$, cf. (270), is almost the characteristic L_∞ -algebra of massive type IIA supergravity (103d), missing H_7 and its nonlinear Bianchi identity:

$$\text{CE}(\mathfrak{I}(\text{KU}_0 // \text{BU}(1))) \simeq \mathbb{R}_d \left[\begin{array}{c} f_{2\bullet} \\ h_3 \end{array} \right] / \left(\begin{array}{l} d f_{2\bullet} = h_3 f_{2\bullet-2} \\ d h_3 = 0 \end{array} \right). \quad (143)$$

- (v) $\mathfrak{I}\text{Cyc}(S^4)$, cf. (274), is the characteristic L_∞ -algebra of actual IIA supergravity (103e), including H_7 and its Bianchi identity:

$$\text{CE}(\mathfrak{I}\text{Cyc}(S^4)) \simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ f_4 \\ f_6 \\ h_3 \\ h_7 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d f_4 = h_3 f_2 \\ d f_6 = h_3 f_4 \\ d h_3 = 0 \\ d h_7 = \frac{1}{2} f_4 f_4 - f_2 f_6 \end{array} \right). \quad (144)$$

- (vi) $\mathfrak{I}S^4$, cf. (269), is the characteristic L_∞ -algebra of 11D supergravity (103f):

$$\text{CE}(\mathfrak{I}S^4) \simeq \mathbb{R}_d \left[\begin{array}{c} g_4 \\ g_7 \end{array} \right] / \left(\begin{array}{l} d g_4 = 0 \\ d g_7 = \frac{1}{2} g_4^2 \end{array} \right). \quad (145)$$

But beware that $\mathfrak{l}(-)$ forgets a lot of structure, namely all the “torsion information”. Accordingly, there is always an infinitude of inequivalent classifying spaces \mathcal{A} with the same rationalization $\mathfrak{L}\mathcal{A}$. For example, with any $K \in \text{Grp}(\text{FinSet})$ we have

$$\mathfrak{l}(\mathcal{A} \times BK) \simeq \mathfrak{L}\mathcal{A}. \quad (146)$$

Accordingly, when we come to *flux quantization* in § 4.1.2, the actual choice of classifying space \mathcal{A} , which is *admissible*, in that

$$\mathfrak{L}\mathcal{A} \simeq \mathfrak{a} \quad (147)$$

for the given characteristic Gauss law L_∞ -algebra \mathfrak{a} , is a substantial datum that needs to be provided/chosen in order to globally complete the higher gauge field theory.

4.1.2. Nonabelian Character

By combining the facts from § 4.1.1, we obtain the natural *differential character maps*

$$\left. \begin{array}{l} X \in \text{SmthMfd} \\ \mathcal{A} \in \text{Grpd}_{[2,\infty]}^{\text{ft}\mathbb{Q}} \end{array} \right\} \vdash \overbrace{\text{Map}(X, \mathcal{A}) \xrightarrow[\text{(130)}]{(\eta_{\mathcal{A}}^{\mathbb{R}})^*} \text{Map}(X, L^{\mathbb{R}}\mathcal{A}) \xrightarrow[\text{(135)}]{\sim} \int \Omega_{\text{cl}}^1(X; \mathfrak{L}\mathcal{A})}^{\text{ch}_X^{\mathfrak{A}}} \quad (148)$$

(where on the left we mean that \mathcal{A} is simply connected and of rational finite type).

These maps hence extract the underlying differential form (flux density!) shadow from maps to the classifying space (charges!).

To make this interpretation more tangible, consider:

- (i) The *nonabelian cohomology* with coefficients in the loop ∞ -group of \mathcal{A} is

$$H^1(X; \Omega\mathcal{A}) := \pi_0 \text{Map}(\int X, \mathcal{A}) \quad (149)$$

and, more generally, if \mathcal{A} is not necessarily connected:

$$H^0(X; \mathcal{A}) := \pi_0 \text{Map}(\int X, \mathcal{A}). \quad (150)$$

([Toë02, Def. 6.0.6; Lur14, Def. 6; FSS23, §2], more exposition in [SS25i, §1; SS26e, §4]).

This subsumes:

- (a) Ordinary nonabelian cohomology with coefficients in a Lie group G , taking $\mathcal{A} \equiv BG$,

$$H^1(X; G) \simeq \pi_0 \text{Map}(X, BG), \quad (151)$$

(cf. [FSS23, Ex. 2.2]).

- (b) Ordinary abelian cohomology with coefficients $A \in \text{AbGrp}$, taking $\mathcal{A} \equiv B^n A$:

$$H^n(X; A) \simeq H^0(X; B^n A) \quad (152)$$

(cf. [FSS23, Ex. 2.1]).

- (c) Complex topological K-theory, taking $\mathcal{A} \equiv BU(\infty) \times \mathbb{Z}$

$$K^0(X) \simeq H^0(X; BU(\infty) \times \mathbb{Z}), \quad (153)$$

(cf. [FSS23, Ex. 2.11]).

- (d) Any Whitehead-generalized abelian cohomology theory $E^n(-)$, taking $\mathcal{A} \equiv E_n$ to be a stage in the corresponding spectrum of classifying spaces:

$$E^n(X) \simeq H^0(X; E_n) \quad (154)$$

(cf. [Koc96, §3.4; AGP02, §12; FSS23, Ex. 2.10]).

(e) n -Cohomotopy, taking $\mathcal{A} \equiv S^n$:

$$\pi^n(X) := H^1(X; \Omega S^n). \quad (155)$$


(ii) The *nonabelian de Rham cohomology* [FSS23, Def. 6.3] with coefficients in $\mathfrak{a} \in L_\infty \text{Alg}_{\mathbb{R}}^{\text{ft}}$ is the set of *concordance classes* of closed forms:

$$H_{\text{dR}}^1(X; \mathfrak{a}) := \Omega_{\text{cl}}^1(X; \mathfrak{a}) / \text{concordance}. \quad (156)$$

This means that a pair of closed \mathfrak{a} -valued forms on X^d are in the same cohomology class iff they are the boundary values of a closed \mathfrak{a} -valued form on $[0, 1] \times X^d$.

By the discussion § 3.2, this means that the nonabelian de Rham class of on-shell flux densities is preserved by time evolution, hence reflects *conserved total flux* ([SS25d, §3.1] cf. Fig. 3):

Figure 3: Time evolution of on-shell flux is a *concordance* of Cauchy data. The concordance class in nonabelian de Rham cohomology (156) is the *conserved total flux*.



For $\mathfrak{a} \equiv \mathbb{I}B^n\mathbb{Z}$ this definition (156) reproduces [FSS23, Prop. 6.4] ordinary de Rham cohomology:

$$H_{\text{dR}}^1(X; \mathbb{I}B^n\mathbb{Z}) \simeq H_{\text{dR}}^n(X). \quad (157)$$

Generally, for $\mathfrak{a} \simeq \mathbb{I}\mathcal{A}$ (132), the following *nonabelian de Rham theorem* [FSS23, Thm. 6.5] naturally identifies nonabelian de Rham cohomology (156) with *nonabelian real cohomology*:

$$\begin{aligned} H_{\text{dR}}^1(X; \mathbb{I}\mathcal{A}) &\equiv \Omega_{\text{cl}}^1(X; \mathbb{I}\mathcal{A}) / \text{concordance} && \text{by (156)} \\ &\simeq \pi_0 \int \Omega_{\text{cl}}^1(X; \mathbb{I}\mathcal{A}) && \text{by (75)} \\ &\simeq \pi_0 \mathbf{Map}(X, L^{\mathbb{R}}\mathcal{A}) && \text{by (135)} \\ &\equiv H^1(X; \Omega L^{\mathbb{R}}\mathcal{A}) && \text{by (149)}. \end{aligned} \quad (158)$$

Therefore, on π_0 the differential character map (148) becomes a map from nonabelian integral to nonabelian real/de Rham cohomology, which as such is the plain *nonabelian character map* [FSS23, Def. IV.2]:

$$H^1(X; \Omega\mathcal{A}) \simeq \pi_0 \mathbf{Map}(X; \mathcal{A}) \xrightarrow{(\text{ch}_X^{\mathcal{A}})^*} \pi_0 \int \Omega_{\text{dR}}^1(X; \mathbb{I}\mathcal{A}) \simeq H_{\text{dR}}^1(X; \mathbb{I}\mathcal{A}). \quad (159)$$

For example:

- (i) $\text{ch}^{B^n\mathbb{Z}}$ is the classical *de Rham map*,
- (ii) ch^{KU_0} is the *Chern character* on K-theory,
- (iii) ch^{E_n} is the *Chern-Dold character* on the generalized abelian cohomology $E^n(-)$,
- (iv) ch^{S^4} is a nonabelian character map from 4-Cohomotopy (269) to classes of total C-field flux (145).

Hence, if we understand:

- (i) nonabelian cohomology as the home of *charge*,
- (ii) nonabelian de Rham cohomology as the home of *total flux*,

then the character map (159) is understood as mapping charges to the total flux which they source.

However, the charges in a cohomology theory are generally more constrained than total flux in real cohomology. For example, the image of the de Rham character

$$H^n(X; \mathbb{Z}) \xrightarrow{\text{ch}_X^{B^n\mathbb{Z}}} H_{\text{dR}}^1(X) \quad (160)$$

is a *lattice* inside a real vector space.

Therefore, asking that some total flux $[\vec{B}]$ *lifts* through the character map, hence that it is the *character image* of a charge χ ,

$$[\vec{B}] = \text{ch}_X^A(\chi), \quad (161)$$

is to ask that it is “quantized” (in the sense of *discretized*) in some form. This is the coarse form of *flux quantization*:

$$\begin{array}{ccc}
 & \xrightarrow{\text{charge } \chi} & H^1(X^d; \mathcal{A}) \\
 * & \xrightarrow{\vec{B} \text{ flux}} & \Omega_{\text{dR}}^1(X^d, \mathfrak{a}) \longrightarrow H_{\text{dR}}^1(X^d; \mathfrak{a}) \\
 & & \downarrow \text{ch}_X^A
 \end{array} \quad (162)$$

and the choice of

$$\mathcal{A} \in \text{Grpd}_\infty \text{ with } \mathfrak{L}\mathcal{A} \simeq \mathfrak{a} \quad (163)$$

is the corresponding *flux quantization law*.

However, the quantization of total flux $[\vec{B}]$ alone is not sufficient for defining a local higher gauge field: We need to *locally* impose structure on the actual flux \vec{B} which *implies* global charges quantizing the corresponding global flux as above.

This is a curious request: Locally the flux may take any value (as long as it satisfies the equations of motion), but globally it must “integrate” to certain prescribed discrete values. But this request has a natural solution in cohesive homotopy theory; this is what we turn to now in § 4.2.

4.2. Global Completion

We are now ready to put all the pieces together and construct the *phase space* stack of globally completed higher gauge fields for a given flux quantization law (§ 4.1).

4.2.1. The Phase Space

The Phase Space As Such. Namely, after a choice \mathcal{A} of flux quantization (163) for given Gauss laws \mathfrak{a} , we have now obtained a pair of coincident maps, one from the smooth moduli set of on-shell flux densities, the other from the moduli space of charges:

$$\begin{array}{ccc} \mathbf{Map}(X^d, \mathcal{A}) & & \\ \text{ch}^{\mathcal{A}} \downarrow (159) & & \\ \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \xrightarrow[\text{(76)}]{\eta^f} \int \Omega_{\text{cl}}^1(X^d; \mathcal{L}\mathcal{A}). & & \end{array} \quad (164)$$

This is the local refinement of the corresponding coincident maps in (162), in that it reproduces the latter on gauge equivalence classes (52):

$$\begin{array}{ccc} \mathbf{Map}(X^d, \mathcal{A}) & \xrightarrow{[-]_0} & H^1(X^d; \Omega\mathcal{A}) \\ \downarrow \text{ch}^{\mathcal{A}} & & \downarrow \text{ch}^{\mathcal{A}} \\ \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \xrightarrow{\eta^f} \int \Omega_{\text{cl}}^1(X^d; \mathcal{L}\mathcal{A}) & \longrightarrow & H_{\text{dR}}^1(X^d; \mathcal{L}\mathcal{A}). \end{array} \quad (165)$$

Therefore we may now ask for *local* flux quantization by asking for a lift as above in (162) but before passing to gauge equivalence classes, and hence up to a specified homotopy \hat{A} :

$$\begin{array}{ccc} & \xrightarrow{\text{charges}} & \mathbf{Map}(X^d, \mathcal{A}) \\ & \hat{A} \swarrow \text{potentials} & \downarrow \text{ch}^{\mathcal{A}} \\ * & \xrightarrow{\text{fluxes}} & \Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \xrightarrow{\eta^f} \int \Omega_{\text{cl}}^1(X^d; \mathcal{L}\mathcal{A}). \end{array} \quad (166)$$

That homotopy \hat{A} turns out to encode the *gauge potentials*. In this way, such a diagram is the proper enhancement of the flux densities to a globally defined higher gauge field configuration.

The entirety of choices of such data forms, by the universal property (59), the moduli stack, which is the homotopy fiber product (58) of our two maps:

$$\begin{array}{ccc} \text{Phs}(X^d; \mathcal{A}) & \longrightarrow & \mathbf{Map}(X^d, \mathcal{A}) \\ \downarrow \lrcorner & & \downarrow \text{ch}^{\mathcal{A}} \\ * & \xrightarrow{\hat{A}} & \Omega_{\text{cl}}^1(X^d; \mathcal{L}\mathcal{A}) \xrightarrow{\eta^f} \int \Omega_{\text{cl}}^1(X^d; \mathfrak{a}). \end{array} \quad (167)$$

This

$$\mathbf{Phs}(X^d; \mathcal{A}) \in \text{SmthGrpd}_{\infty} \quad (168)$$

is the globally completed *phase space* of the global completion of our Maxwell-type higher gauge theory by flux quantization in \mathcal{A} -cohomology, in that it is the gauged smooth set of (Cauchy data for) *on-shell* higher gauge fields, hence those solving their equations of motion (here: of Maxwell-type).

We notice that the above homotopy pullback is the image of a homotopy pullback under $\mathbf{Map}(X^d, -)$, by (126) and (127):

$$(167) \simeq \begin{array}{ccc} \mathbf{Map}(X^d, \mathcal{A}_{\text{diff}}) & \xrightarrow{\quad} & \mathbf{Map}(X^d, \mathcal{A}) \\ \downarrow & \lrcorner & \downarrow_{\mathbf{Map}(X^d, \text{ch}^d)} \\ \mathbf{Map}(X^d, \Omega_{\text{cl}}^1(*; \mathfrak{a})) & \xrightarrow{\mathbf{Map}(X^d, \eta^1)} & \mathbf{Map}(X^d, \int \Omega_{\text{cl}}^1(*; \mathfrak{L}\mathcal{A})) \end{array} \quad (169)$$

Here in the top left we used that $\mathbf{Map}(X^d, -)$ preserves homotopy limits, and identified the *differential moduli stack* of \mathcal{A} :

$$\mathcal{A}_{\text{diff}} := \mathcal{A} \times_{\int \Omega_{\text{cl}}^1(*; \mathfrak{L}\mathcal{A})} \Omega_{\text{cl}}^1(*; \mathfrak{L}\mathcal{A}) \in \text{SmthGrpd}_\infty. \quad (170)$$

This object neatly exhibits the combination of plain homotopy types (\mathcal{A} , on the left) with differential geometric spaces ($\Omega_{\text{cl}}^1(*; \mathfrak{a})$, on the right) that we achieve among smooth ∞ -groupoids (cf. Fig. 1).

The Gauge Potentials. Among the most profound aspects of this construction is that the homotopies in (166) are really locally equivalent to higher gauge potentials as known in the traditional literature. This is the case but requires non-trivial analysis:

Example 4.2.

- (i) **Differential ordinary cohomology.** For ordinary cohomology coefficients, $\mathcal{A} \equiv B^{n+1}\mathbb{Z}$, [FSS23, Prop. 9.5] shows that that the differential moduli stack (170) is that classifying ordinary differential $(n+1)$ -cohomology (Deligne cohomology, higher $U(1)$ -bundle $(n-1)$ -gerbes with connection, cf. [FSS13, §2.5; FSS15a, §3.1; FRS16, §2.8])

$$(B^{n+1}\mathbb{Z})_{\text{diff}} \simeq \mathbf{B}^n U(1)_{\text{conn}}. \quad (171)$$

This shows that the above construction reproduces the usual global completions of abelian higher gauge fields, such as the ordinary magnetic field in ordinary differential 2-cohomology (*Dirac charge quantization*, cf. [Alv85; Fre02, §2; LS22b]), the Kalb-Ramond B-field in ordinary differential 3-cohomology ([Gaw88; FW99]), and so on; see [SS25d, Ex. 3.10] for further pointers.

- (ii) **Differential K-theory.** For $\mathcal{A} \equiv KU_0$ (or $\mathcal{A} \equiv KU_0 // BU(1)$) flux-quantizing type IIA supergravity with the H_7 -Bianchi disregarded (143), [FSS23, Exs. 9.2, 11.2] shows that $\mathcal{A}_{\text{diff}}$ (170) is the moduli stack for (twisted) differential K-theory, a candidate model for the gauge potentials of the RR-field in 10D supergravity (cf. [Fre02; Sza13, §3; GS22]).
- (iii) **Differential abelian cohomology.** Generally, for $\mathcal{A} \equiv E_n$ a stage in a spectrum E_\bullet of spaces, representing a (Whitehead-generalized) abelian cohomology theory, E_{diff}^n is [FSS23, Ex. 9.1] the moduli stack for differential E -cohomology ([HS05, §4], cf. [Bun12]) in degree n .

But this construction generalizes now further to differential refinement of *nonabelian* cohomology (150), such as:

- (iv) **Differential 4-cohomotopy.** For $\mathcal{A} \equiv S^4$ flux-quantizing 11D SuGra (145), the differential cohomology theory classified by S_{diff}^4 (170) is *differential 4-Cohomotopy*

[FSS15c, §4; GS21, §3.1]. That this indeed induces the local gauge potentials (and their local gauge transformations) expected in 11D SuGra (cf. § 6.1) is shown in [GSS24a, Prop. 1.1; GSS24b, §4.1; Ban25].

The Shape of Phase Space. The shape of the phase space is the groupoid of *global symmetries* exhibited by *large gauge transformations*.

The properties (77) and (78) immediately imply the following crucial fact: *The shape of the phase space is the moduli space of the topological charges:*

$$\int \mathbf{Phs}(X^d; \mathcal{A}) \simeq \mathbf{Map}(X^d; \mathcal{A}). \quad (172)$$

Hence the gauge equivalence classes (52) of the shape (71) of the phase space (167) constitute the nonabelian cohomology (149) of the Cauchy surface with coefficients in the flux quantization law \mathcal{A} (163):

$$[\int \mathbf{Phs}(X^d; \mathcal{A})]_0 \simeq H^1(X^d; \mathcal{A}). \quad (173)$$

The Points of Phase Space. If we forget the smooth structure on phase space, by applying the flat modality $\flat(-)$ (81), and then pass to gauge equivalence classes (52), we get the discrete set of gauge equivalence classes of on-shell field configurations:

$$[\flat \mathbf{Phs}(X^d; \mathcal{A})]_0 \in \text{Set}. \quad (174)$$

This is equivalently the cohomology of the Cauchy surface with coefficients in the differential moduli stack $\mathcal{A}_{\text{diff}}$ (170)

$$\begin{aligned} [\flat \mathbf{Phs}(X^d; \mathcal{A})]_0 &\simeq [\flat \mathbf{Map}(X^d, \mathcal{A}_{\text{diff}})]_0 && \text{by (169)} \\ &\simeq [\text{Hom}(X^d, \mathcal{A}_{\text{diff}})]_0 \\ &=: H_{\text{diff}}^1(X^d; \Omega \mathcal{A}), \end{aligned} \quad (175)$$

which is the *differential nonabelian cohomology* [FSS23, Def. 9.3] of the Cauchy surface, with coefficients in \mathcal{A} .

For example:

- (i) For $\mathcal{A} \equiv B^2\mathbb{Z}^2$ flux-quantizing 4D vacuum Maxwell theory (140), the completed gauge fields have classes in (two copies of) *ordinary differential cohomology in degree 2*,

$$H_{\text{diff}}^1(X^d; \Omega B^2\mathbb{Z}) \simeq H_{\text{diff}}^2(X^d; \mathbb{Z}) \quad (176)$$

(cf. [FSS23, Prop. 9.5]).

This is equivalently the usual flux quantization that one finds in modern textbooks (going back to [Dir31; Sch66; Zwa68]), equivalently given by isomorphism classes of $U(1)$ -principal connections (cf. [Alv85; Bry93, §7.1; Fra11, §16.4e; Fre02, §2]).

The consideration of quantization not just of the magnetic but also of the electric flux, hence of differential coefficients in $B^2\mathbb{Z}^2$, is less widely appreciated but has been considered in [FMS07b; FMS07a; Bec+17, Rem. 2.3; LS22a; LS22b; SS24b, (21)].

Of course, one may also choose not to integrally quantize the electric fluxes at all, hence to take the flux quantization law to be $B^2\mathbb{Z} \times B^2\mathbb{Q}$. If one means to model real-world electromagnetism, then the “right” choice of flux quantization is a matter of comparison to experiment.

While integral magnetic flux quantization has been securely observed experimentally (in the form of flux quanta attached to *Abrikosov vortices* in type II superconductors, cf. [Tim23, §5.3]) the phenomenology of electric flux quantization may yet have to receive attention.

- (ii) For $\mathcal{A} \equiv B^3\mathbb{Z}$ flux-quantizing the NS field in 10D supergravity (143), the completed gauge fields have classes in *ordinary differential cohomology in degree 3* (again by [FSS23, Prop. 9.5]):

$$H_{\text{diff}}^1(X^d; \Omega B^3\mathbb{Z}) \simeq H_{\text{diff}}^3(X^d; \mathbb{Z}). \quad (177)$$

This is the flux quantization traditionally considered for this *NS B-field*, equivalently given by *Deligne cohomology in degree 3* and by *bundle gerbes with connection and curving* [Gaw88; FW99; GR02; CJM04; BFS08; Fer12; GS19].

- (iii) For $\mathcal{A} \equiv \text{KU}_0$ flux-quantizing massive 10D type IIA supergravity with trivial NS-field background (143), the completed gauge fields have classes [FSS23, Ex. 9.2] in *differential K-theory*:

$$H_{\text{diff}}^1(X^d; \Omega \text{KU}_0) \simeq K_{\text{diff}}^0(X^d) \quad (178)$$

(the twisting by the background NS-field is readily incorporated, too; we come to this in §§ 5.1 and 6.1.)

This is the RR-flux quantization traditionally considered in the string theory literature [Fre01, §3.7; Fre02; KV14; GS22], going back to suggestions in [GHM97; MM97; MW00], cf. [SS25d, §4.1].

- (iv) For $\mathcal{A} \equiv S^4$ flux-quantizing 11D supergravity (145), the completed gauge fields have classes in *differential 4-Cohomotopy* [FSS15c, §4; FSS23, Ex. 9.3; GS21, §3.1].

The Reduced Phase Space. The *reduced phase space* (cf. [HT92]) is the (concrete, § 2.1.2) quotient of the phase space by gauge symmetries (52):

$$\sharp_1[\mathbf{Phs}(X^d; \mathcal{A})]_0 \in \text{SmthSet}. \quad (179)$$

We now have the following fundamental result:

The points of the reduced phase space are the differential nonabelian cohomology classes of the Cauchy surface, with coefficients in the flux quantization law \mathcal{A} .

This holds by these natural isomorphisms:

$$\begin{aligned} \flat(\sharp_1[\mathbf{Phs}(X^d; \mathcal{A})]_0) &\simeq \flat[\mathbf{Phs}(X^d; \mathcal{A})]_0 && \text{by (38)} \\ &\simeq [\flat\mathbf{Phs}(X^d; \mathcal{A})]_0 && \text{by (82)} \\ &\simeq H_{\text{diff}}^1(X^d; \mathcal{A}) && \text{by (175)}. \end{aligned} \quad (180)$$

Restriction to Solitonic Fields. In variation of the construction (167), we may restrict to solitonic fields whose flux densities \vec{B} on the Cauchy surface X^d vanish at infinity (136):

$$\vec{B} \in \Omega_{\text{cl}, \mathbf{c}}^1(X^d; \mathfrak{a}) \hookrightarrow \Omega_{\text{cl}}^1(X^d; \mathfrak{a}). \quad (181)$$

By Prop. 4.1, the shape of these compactly supported flux moduli is the pointed mapping space out of the one-point compactification $X_{\{0, \infty\}}^d$ of phase space. Therefore, the variant of

the phase space (167) for solitonic fields is the following pullback:

$$\begin{array}{ccc} \mathbf{Phs}_c(X^d; \mathcal{A}) & \longrightarrow & \mathbf{Map}^*(X_{U\{\infty\}}^d, \mathcal{A}) \\ \downarrow & \lrcorner & \downarrow \mathbf{ch}^{\mathcal{A}} \\ \Omega_{\text{cl},c}^1(X^d; \mathfrak{a}) & \xrightarrow{\eta^f} & \int \Omega_{\text{cl},c}^1(X^d; \mathfrak{a}). \end{array} \quad (182)$$

The shape of this solitonic phase space is thus found, analogous to (172), to be the space of pointed maps into the classifying space:

$$\int \mathbf{Phs}_c(X^d; \mathcal{A}) \simeq \mathbf{Map}^*(X_{U\{\infty\}}^d, \mathcal{A}). \quad (183)$$

4.2.2. The Observables

Ordinary Observables. An ordinary *observable* is (cf. [SW76, §4; Kir90, §1.1]) a smooth map from phase space to the complex numbers, sending on-shell field configurations Φ to the given value $\mathcal{O}(\Phi)$ that is observed/measured:

$$\begin{array}{ccc} \mathbf{Phs}(X^d; \mathcal{A}) & \xrightarrow{\mathcal{O}} & \mathbb{C} \\ \Phi & \longmapsto & \mathcal{O}(\Phi). \end{array} \quad (184)$$

For definiteness: By

$$\mathbb{C} \in \text{Alg}(\text{SmthMfd}) \hookrightarrow \text{Alg}(\text{SmthGrpd}_\infty), \quad (185)$$

we mean the complex numbers with their canonical smooth structure as the Euclidean space \mathbb{R}^2 . Being a smooth manifold, as a gauged smooth set, the complex numbers are therefore both concrete (33) as well as 0-truncated (51):

$$\begin{array}{l} \mathbb{C} \simeq \sharp_1 \mathbb{C} \\ \mathbb{C} \simeq [\mathbb{C}]_0. \end{array} \quad (186)$$

But this immediately implies, by the universal properties (37) and (53), that ordinary observables (184) factor through the reduced phase space (179):

$$\mathbf{Phs}(X^d; \mathcal{A}) \longrightarrow \sharp_1[\mathbf{Phs}(X^d; \mathcal{A})]_0 \dashrightarrow \mathbb{C} \quad (187)$$

This is exactly the expected/desired physical property of ordinary observables: They are gauge invariant (and send smooth families of field configurations to smooth families of observed numbers).

Topological Observables. An observable (184) is *topological* (physics jargon for: *homotopical*) if it factors through the shape of the phase space.

$$\mathbf{Phs}(X^d; \mathcal{A}) \xrightarrow{\eta^f} \int \mathbf{Phs}(X^d; \mathcal{A}) \dashrightarrow \mathbb{C}. \quad (188)$$

Combined with the implied gauge invariance (187) and with the characterization (173) of the shape of phase space, this means that ordinary topological observables are functions of the nonabelian cohomology of the Cauchy surface with coefficients in $\Omega\mathcal{A}$, hence of the set

of topological charges:

$$\overbrace{\mathbf{Phs}(X^d; \mathcal{A}) \xrightarrow{\eta^f} \int \mathbf{Phs}(X^d; \mathcal{A}) \xrightarrow{\eta^{[-1]_0}} H^1(X^d; \Omega \mathcal{A}) \dashrightarrow \mathbb{C}}^{\mathcal{O}_{\text{top}}} \quad (189)$$

Moreover, *realistic* observables (those measurable in experiment) will have compact support on phase space, whence realistic topological observables are actually compactly supported functions on the connected components of the shape of phase space, as such identified with the 0-homology of phase space:

$$\begin{aligned} \text{TopObs}_0(X^d; \mathcal{A}) &\simeq H_0(\int \mathbf{PhsSp}(X^d; \mathcal{A}); \mathbb{C}) \\ &\simeq H_0(\text{Map}(X^d; \mathcal{A}); \mathbb{C}). \end{aligned} \quad (190)$$

Or rather, restricting to solitonic field configurations (183) whose flux densities vanish at infinity (136) along the Cauchy surface, the realistic topological observables (190) constitute homology of the pointed mapping space (183) out of the one-point compactification of the Cauchy surface:

$$\text{SolTopObs}_0(X^d; \mathcal{A}) = H_0(\text{Map}^*(X_{\cup\{\infty\}}^d, \mathcal{A}); \mathbb{C}). \quad (191)$$

Topological Light Cone Quantization. Consider then the case that spacetime is not just globally hyperbolic but of the form

$$X^{1,d} \simeq \mathbb{R}^{1,1} \times X^{d-1}, \quad (192)$$

whence the Cauchy surfaces are of the form

$$X^d \simeq \mathbb{R}^1 \times X^{d-1}. \quad (193)$$

On such spacetimes one may naturally consider *light-front evolution* ([Dir49, §5], cf. [WP94]).

Remarkably, the realistic solitonic topological observables (191) on such spacetimes (192) form the homology of a based loop space (cf. [SS25g, (18)])

$$\begin{aligned} \text{SolTopObs}_0(X^d; \mathcal{A}) &\simeq H_0(\text{Map}^*((\mathbb{R}^1 \times X^{d-1})_{\cup\{\infty\}}, \mathcal{A}); \mathbb{C}) \\ &\simeq H_0(\Omega \text{Map}^*(X_{\cup\{\infty\}}^{d-1}, \mathcal{A}); \mathbb{C}) \end{aligned} \quad (194)$$

and as such inherit [SS25g, (20)] a (generally) non-commutative (star-)algebra structure induced by loop concatenation (and reversal), the *Pontrjagin algebra* structure:

$$\text{SolTopObs}_0 \otimes_{\mathbb{C}} \text{SolTopObs}_0 \xrightarrow{\text{conc}^*} \text{SolTopObs}_0. \quad (195)$$

As such, SolTopObs_0 has the form of an algebra of quantum observables⁷, which one may understand as the result of *light-front quantization* [SS26c, §2.2] in the topological sector.

Concretely, with the restriction to homology in degree 0, this is the group algebra of the fundamental group of the pointed maps from $X_{\cup\{\infty\}}^{d-1}$:

$$\text{SolTopObs}_0(X^d; \mathcal{A}) \simeq \mathbb{C}[\pi_1 \text{Map}^*(X_{\cup\{\infty\}}^{d-1}, \mathcal{A})]. \quad (196)$$

⁷Beware that usual “canonical” deformation quantization of symplectic/Poisson structures cannot see non-trivial quantum algebra structure of topological observables, as these are locally constant functions on phase space and as such have vanishing Poisson brackets.

5. Enhancements

We briefly indicate some natural further enhancements of the above constructions.

For brevity we omit the important topic of higher gauge fields on *orbifolds*, flux-quantized in twisted *equivariant* nonabelian cohomology theory. For more on that see [SS25i; SS26e].

5.1. Sources and Branes

Maxwell Theory with Sources. The classical Maxwell equations of course generally have a source term J_3 , the electric current density 3-form, and are as such traditionally written as:

$$“d F_2 = 0, \quad d G_2 = J_3, \quad \star F_2 = G_2” \quad (197)$$

Here we have put this traditional expression in quotation marks because it glosses over the important subtlety that the topological class of J_3 is on a different footing compared to that of F_2 and G_2 :

- (i) The electric current density J_3 is meant to be integrable over the Cauchy surface, its integral being the net number of electrons.
- (ii) Its class is not in plain cohomology but subject to coboundaries that preserve this integral, and hence the net number of electrons.
- (iii) The Maxwell equation $d G_2 = J_3$ is meant to apply after forgetting these integrability conditions, so that it trivializes only the plain de Rham class of J_3 , without forcing the net number of electrons to vanish.

We may mathematically implement these desiderata by considering the *one-point compactification* $X_{\{ \infty \}}^d$ of the Cauchy surface, assuming for simplicity that this still has the structure of a smooth manifold (for instance $\mathbb{R}_{\{ \infty \}}^3 \simeq S^3$). Then we have a canonical smooth map

$$X^d \xrightarrow{\phi} X_{\{ \infty \}}^d \quad (198)$$

with $F_2, G_2 \in \Omega_{\text{dR}}^2(X^d)$ while $J_3 \in \Omega_{\text{dR}}^3(X_{\{ \infty \}}^d)$, and with the would-be Maxwell equations (197) really reading as follows:

$$\begin{aligned} d F_2 &= 0, \\ d G_2 &= \phi^* J_3, \quad \star F_2 = G_2. \\ d J_3 &= 0. \end{aligned} \quad (199)$$

Such a system of EoMs no longer has a characteristic L_∞ -algebra, but a *fibration* of such:

$$\begin{array}{ccc} \mathfrak{a} & \text{CE}(\mathfrak{a}) := \mathbb{R}_d \begin{bmatrix} f_2 \\ g_2 \\ j_3 \end{bmatrix} / \begin{pmatrix} d f_2 = 0 \\ d g_2 = j_3 \\ d j_3 = 0 \end{pmatrix} & \\ \downarrow \iota_\phi & \uparrow \iota_\phi^* & \begin{array}{c} \uparrow j_3 \\ \uparrow \\ \downarrow j_3 \end{array} \\ \mathfrak{b} & \text{CE}(\mathfrak{b}) := \mathbb{R}_d [j_3] / (d j_3 = 0), & \end{array} \quad (200)$$

in that germs of solutions of Maxwell's equations with sources (199) are equivalent to commuting squares of this form:

$$\begin{array}{ccc} X^d & \dashrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{a}) \\ \downarrow \phi & & \downarrow \phi^* \\ X_{\cup\{\infty\}}^d & \dashrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{b}). \end{array} \quad (201)$$

Background Fluxes. Generally, given

(i) an embedding of smooth manifolds

$$\Sigma^p \xrightarrow{\phi} X^d, \quad (202)$$

(ii) a fibration of finite-type L_∞ -algebras

$$\mathfrak{a} \xrightarrow{\iota_\phi} \mathfrak{b}, \quad (203)$$

then a dashed map completing a commuting diagram of the form

$$\begin{array}{ccc} \Sigma^p & \dashrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{a}) \\ \downarrow \phi & & \downarrow (\iota_\phi)^* \\ X^d & \longrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{b}) \end{array} \quad (204)$$

encodes on-shell flux densities on Σ^p whose Bianchi identities may be *twisted* by *background fluxes* on X^d .

Examples:

(i) *Massive RR-Fluxes Twisted by NS-Fluxes:*

$$\begin{array}{ccc} X^9 & \dashrightarrow^{F_{2\bullet}} & \Omega_{\text{cl}}^1(*; \mathfrak{l}(\text{KU}_0 // \text{BU}(1))) & d F_{2\bullet} = F_{2\bullet-2} \wedge \phi^* H_3 \\ \downarrow & & \downarrow & \\ X_{\text{cpt}}^9 & \xrightarrow{H_3} & \Omega_{\text{cl}}^1(*; \mathfrak{l}B^2\text{U}(1)) & d H_3 = 0. \end{array} \quad (205)$$

(ii) *Self-Dual Flux on M5 Twisted By C-Field Flux:*

$$\begin{array}{ccc} \Sigma^5 & \xrightarrow{H_3} & \Omega_{\text{cl}}^1(*; \mathfrak{l}_{S^4} S^7) & d H_3 = \phi^* G_4 \\ \downarrow \phi & & \downarrow (\iota_{\mathfrak{C}})^* & \\ X^{10} & \xrightarrow{G_4, G_7} & \Omega_{\text{cl}}^1(\mathfrak{l}S^4) & d G_4 = 0 \\ & & & d G_7 = \frac{1}{2} G_4 \wedge G_4. \end{array} \quad (206)$$

Twisted Relative Gauge Fields. Our construction above of globally completed higher gauge fields carries over straightforwardly to this case of relative Bianchi identities twisted by background fluxes, simply by passing to the *arrow category* of SmthGrpd_∞ :

The choice of classifying space for (now: twisted relative) flux quantization is enhanced to a choice of fibration

$$\mathcal{A} \xrightarrow{\wp} \mathcal{B}, \quad (207)$$

whose *relative Whitehead* L_∞ -algebra

$$\mathfrak{l}_{\mathcal{B}} \mathcal{A} \xrightarrow{\iota_\wp} \mathfrak{l} \mathcal{B} \quad (208)$$

characterizes the twisted Bianchi identities as above (204).

Now given such a pair of maps ϕ, φ , consider their *twisted relative mapping space*

$$\mathbf{Map}(\phi, \varphi) := \mathbf{Map}(\Sigma; \mathcal{A}) \times_{\mathbf{Map}(\Sigma, \mathcal{B})} \mathbf{Map}(X; \mathcal{B}) = \left\{ \begin{array}{ccc} \Sigma & \dashrightarrow & \mathcal{A} \\ \downarrow \phi & & \downarrow \varphi \\ X & \dashrightarrow & \mathcal{B} \end{array} \right\}, \quad (209)$$

and similarly the *twisted relative flux moduli*

$$\Omega_{\text{cl}}^1(\phi; \mathfrak{l}\varphi) := \Omega_{\text{cl}}^1(\Sigma; \mathfrak{l}_{\mathcal{B}}\mathcal{A}) \times_{\Omega_{\text{cl}}^1(\Sigma; \mathfrak{l}\mathcal{B})} \Omega_{\text{cl}}^1(X; \mathfrak{l}\mathcal{B}) = \left\{ \begin{array}{ccc} \Sigma & \dashrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{l}_{\mathcal{B}}\mathcal{A}) \\ \downarrow \phi & & \downarrow (\mathfrak{l}\varphi)_* \\ X & \longrightarrow & \Omega_{\text{cl}}^1(*; \mathfrak{l}\mathcal{B}) \end{array} \right\}. \quad (210)$$

Applying rationalization componentwise, we obtain the twisted relative version of the differential character map and from this the twisted relative version of the phase space, as the homotopy pullback:

$$\begin{array}{ccc} \mathbf{Phs}(\phi; \varphi) & \longrightarrow & \mathbf{Map}(\phi; \varphi) \\ \downarrow & \lrcorner & \downarrow \text{ch}^\varphi \\ \Omega_{\text{cl}}^1(\phi; \mathfrak{l}\varphi) & \xrightarrow{\eta^f} & \int \Omega_{\text{cl}}^1(\phi; \mathfrak{l}\varphi) \end{array} \quad (211)$$

(We are using here that shape preserves the fiber product (210), by (79), since $\mathfrak{l}\varphi$ is a fibration.)

For more details on such twisted (differential nonabelian) cohomology see [FSS23, §3 & §V]. For early discussion of applications to string/M-theory see [Sat11].

5.2. CS/BF-Like Fields

Above we discussed equations of motion of higher Maxwell-type. Here we generalize to the situation where some of the flux densities $F^{(i)}$ in addition satisfy the simple but consequential equation of motion of the form:

$$F^{(i)} = 0.$$

Such equations of motion famously hold in abelian Chern-Simons theories of p -form fluxes in $D = 2p - 1$, and for higher BF-theories.

It is a truism that such EoM imply a *topological field theory*, with the entire field content consisting of the gauge potentials and their topological charges. But we are now positioned to accurately say what this means globally, particularly when there is coupling to further non-topological fields.

The relevant generalization of the above discussion is straightforward: In § 4.2.1 we bluntly took the map from the moduli of on-shell flux densities to the rationalized classifying space to be the canonical one, the shape unit

$$\Omega_{\text{cl}}^1(X^d; \mathfrak{a}) \xrightarrow{\eta^f} \int \Omega_{\text{cl}}^1(X^d; \mathfrak{a}). \quad (212)$$

But in order to construct a phase space along the lines of (166), all it takes is any map of this form.

In particular, we may consider that the characteristic L_∞ -algebra of the Maxwell-type

higher gauge fields is not actually isomorphic to $\mathfrak{L}\mathcal{A}$, but just equipped with a morphism

$$\mathfrak{a} \xrightarrow{\prime} \mathfrak{L}\mathcal{A}. \quad (213)$$

Forming the corresponding homotopy pullback immediately entails that generators in $\mathfrak{L}\mathcal{A}$ that are not in the image of (213) correspond to gauge fields whose gauge potentials do appear in the phase space, but whose flux densities vanish.

In this case the conclusion (172) may no longer hold, and the 0-truncated phase space may coincide with plain \mathcal{A} -cohomology only up to rational corrections.

A physical application of this situation is discussed in [SS26a].

6. Examples and Applications

6.1. 11D/10D Supergravity

M-Flux Quantization in 4-Cohomotopy. We have seen that the characteristic L_∞ -algebra of 11D supergravity is $\mathfrak{a} \equiv \mathfrak{L}S^4$. This means that one admissible flux quantization law in 11D (in fact the minimal one, in numbers of cells) is [Sat18, §2.5; GSS24a] the actual 4-sphere

$$\mathcal{A} \equiv \int S^4, \quad (214)$$

which is the classifying space for *4-Cohomotopy* cohomology theory (cf. [Hu59, §VII; FSS23, Ex. 2.7]):

$$\pi^4(X) \simeq H^1(X; \Omega S^4). \quad (215)$$

This may naturally be twisted by $\mathrm{Sp}(2)$ -tangential structure, in which guise it implies [FSS20] several subtle topological conditions expected in M-theory (cf. [Wit97b; Wit97a]), such as:

- [FSS20, Prop. 3.13]: the $\frac{1}{4}p_1$ -shifted half-integral flux quantization of G_4 ,
- [FSS21a; SS21]: the cancellation of M5-brane anomalies.

Double Dimensional Reduction. The operation of *double dimensional reduction* along principal circle fiber bundles is implemented on flux quantized higher gauge fields by *cyclification* of their classifying spaces ([BSS19, §2.2; SS24a, §2.2]):

$$\mathrm{Cyc}(\mathcal{A}) := \mathbf{Map}(S^1, \mathcal{A}) // S^1 \quad (216)$$

in that for S^1 -principal bundles

$$\begin{array}{ccc} X^{10} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ X^9 & \xrightarrow{c_1} & BS^1 \end{array} \quad (217)$$

there is a natural equivalence of classifying maps

$$\left\{ X^{10} \dashrightarrow \mathcal{A} \right\} \simeq \left\{ \begin{array}{ccc} X^9 & \dashrightarrow & \mathrm{Cyc}(\mathcal{A}) \\ & \searrow^{c_1} & \swarrow \\ & BS^1 & \end{array} \right\} \quad (218)$$

with compatible rationalization. In particular, the characteristic L_∞ -algebra (Def. 3.2) of actual type IIA supergravity (103e) is [FSS17, §3; SV23, Ex. 2.5; GSS25, Ex. 3.10] the

Whitehead bracket L_∞ -algebra of the cyclic loop space of the 4-sphere:

$$\mathfrak{a} \equiv \mathbb{I}\text{Cyc}(S^4), \quad \text{CE}(\mathbb{I}\text{Cyc}(S^4)) \simeq \mathbb{R}_d \left[\begin{array}{c} f_2 \\ f_4 \\ f_6 \\ h_3 \\ h_7 \end{array} \right] / \left(\begin{array}{l} d f_2 = 0 \\ d f_4 = h_3 \wedge f_2 \\ d f_6 = h_3 \wedge f_4 \\ d h_3 = 0 \\ d h_7 = \frac{1}{2} f_4 f_4 - f_2 f_6 \end{array} \right). \quad (219)$$

IIA Flux Quantization in Twisted Unstable K-Theory. Another admissible flux quantization law in 11D turns out to be $\mathcal{A} \equiv \widehat{BSU(2)}$, the homotopy fiber of the square of the second Chern class c_2 on $BSU(2)$:

$$\begin{array}{ccc} \widehat{BSU(2)} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ BSU(2) & \xrightarrow{(c_2)^2} & B^8\mathbb{Z}. \end{array} \quad (220)$$

The canonical comparison map induces isomorphisms on $\pi_{\leq 7}$:

$$S^4 \xrightarrow{\sim_{\leq 7}} \widehat{BSU(2)}. \quad (221)$$

Under double dimensional reduction, this entails a flux quantization of type IIA supergravity in $\text{Cyc}(\widehat{BSU(2)})$, which receives a canonical map

$$\widehat{BU(2)} \longrightarrow \text{Cyc}(\widehat{BSU(2)}), \quad (222)$$

exhibiting it as a twisted deformation of “unstable K-theory” [BSS26a].

This yields a consistent global completion of actual (instead of massive) 10D supergravity of type IIA which retains an (unstable, deformed) semblance of the famously conjectured flux quantization in twisted K-theory, compatible with oxidation to globally completed 11D supergravity.

Superspace formulation. The development of the higher gauge theory in § 4 capitalized on the fact (Thm. 3.3) that (germs of) solutions to higher Maxwell-type equations of motion are equivalent to closed L_∞ -valued forms when restricted to a Cauchy surface (solutions to the electromagnetic Gauss law constraints). Here it is the implicit time-evolution (107) away from the Cauchy surface which absorbs the Hodge duality constraint from the higher Maxwell-type equations and leaves the purely cohomological L_∞ -closure condition that allows for cohomological flux quantization of the theory (§ 4.1.2).

Remarkably, for some supergravity theories there is an alternative *covariant* form — meaning: not requiring existence or choice of Cauchy surfaces — of this “cohomologization by metric decoupling”.

This requires formulating the theory on super-spacetime and constraining the fermionic form of the super-flux densities (cf. [GSS24a, §2] for relevant background on supergeometry):

Specifically, consider a super-spacetime $X^{(1,10|\mathbf{32})}$ with super-torsion free super-coframe

$$\left(\begin{array}{l} (E^a \in \Omega^{1,\text{evn}}(X^{1,10|\mathbf{32}}))_{a=0}^{10} \\ (\Psi^\alpha \in \Omega^{1,\text{odd}}(X^{1,10|\mathbf{32}}))_{\alpha=1}^{32} \end{array} \right) \quad (223)$$

and promote the ordinary duality-symmetric flux density forms G_4 and G_7 (103f) to differential superforms G_4^s and G_7^s with these specific components:

$$\begin{aligned} G_4^s &:= \frac{1}{4!}(G_4)_{a_1 \dots a_4} E^{a_1} \dots E^{a_4} + \frac{1}{2}(\overline{\Psi} \Gamma_{a_1 a_2} \Psi) E^{a_1} E^{a_2} \\ G_7^s &:= \frac{1}{7!}(G_7)_{a_1 \dots a_7} E^{a_1} \dots E^{a_7} + \frac{1}{5!}(\overline{\Psi} \Gamma_{a_1 \dots a_5} \Psi) E^{a_1} \dots E^{a_5}. \end{aligned} \quad (224)$$

Then we have the following remarkable result, following [CF80; BH80; CL94, §6; CDF91, §III.8.5]:

Proposition 6.1 ([GSS24a, Thm. 3.1]). *Fields E (graviton), Ψ (gravitino) and G_4, G_7 (duality-symmetric C-field) satisfy the equations of motion of 11D Supergravity iff the corresponding super-flux densities (224) are $\mathfrak{L}S^4$ -closed, hence iff*

$$(G_4^s, G_7^s) \in \Omega_{\text{cl}}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4). \quad (225)$$

In particular the Hodge-duality relation (103f) is implied [CDF91, p. 878; GSS24a, Lem. 3.3(iii)]:

$$(G_4^s, G_7^s) \in \Omega_{\text{cl}}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4) \quad \Rightarrow \quad G_7 = \star G_4. \quad (226)$$

This entails that 11D SuGra admits *covariant* flux quantization (not requiring existence or choice of Cauchy surfaces) along the same lines as § 4 but replacing the Cauchy surface X^{10} by super-spacetime $X^{1,10|\mathbf{32}}$ and using only the sub-sheaf of $\mathfrak{L}S^4$ -closed super-forms whose components are of the form (224):

$$\begin{aligned} \Omega_{\text{cl},s}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4) &:= \{(G_4^s, G_7^s) \in \Omega_{\text{cl}}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4) \mid \text{of the form (224)}\}, \\ \Omega_{\text{cl},s}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4) &:= \Omega_{\text{cl}}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4) \times_{\sharp_1 \Omega_{\text{cl}}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4)} C^\infty(-, \Omega_{\text{cl},s}^1(X^{1,10|\mathbf{32}}; \mathfrak{L}S^4)). \end{aligned} \quad (227)$$

Beware that this smooth moduli set of constrained super-flux densities depends on the choice of super-coframe (223) which encodes the super-gravitational background field. But in this form it is straightforward to generalize to the smooth moduli set of compatible coframe fields and flux fields. This leads to flux quantization of 11D SuGra beyond just its gauge sector.

Such super-space flux quantization is not restricted to 11D supergravity:

- (i) After promoting the M5-brane embedding (more on which in § 6.2) to a $1/2$ -BPS *super-embedding*

$$\Sigma^{1,5|2\cdot 8} \xrightarrow{\phi_s} X^{1,10|\mathbf{32}} \quad (228)$$

of a super-worldvolume with coframe field $((e^a)_{a=0}^5, (\psi^\alpha)_{\alpha=1}^{16})$, the notorious non-linear self-duality (cf. [FSS21b, Rem. 3.19]) of the 3-form flux H_3 on the 5-brane is *implied* ([FSS21b, Prop. 3.18(iii), Rem. 3.20] following [HS97; HSW97]) by imposing on its super-flux

$$H_3^s := (H_3)_{a_1 a_2 a_3} e^{a_1 a_2 a_3} \quad (229)$$

the $\mathfrak{L}_{S^4} S^7$ -valued Bianchi identity

$$dH_3^s = \phi_s^* G_4^s, \quad (230)$$

hence by the condition that

$$((G_4^s, G_7^s), H_3^s) \in \Omega_{\text{cl}}^1(\phi_s; \mathfrak{h}_{\mathbb{C}}) \quad (231)$$

constitute twisted relative closed differential forms (210) with coefficients in the quaternionic Hopf fibration $S^7 \xrightarrow{h_{\mathbb{H}}} S^4$ (cf. [FSS20, §3.7; FSS21a]).

- (ii) After dimensional reduction on superspace, an analogous statement [GS26] characterizes the actual equations of motion (103e) of 10D IIA supergravity as equivalent to the constraint that corresponding super-flux densities constitute a closed $\mathbf{ICyc}(S^4)$ -valued (219) differential form on super-spacetime $X^{1,9|\mathbf{16}\oplus\overline{\mathbf{16}}}$:

$$(F_s^2, F_4^s, F_7^s, H_3^s, H_7^s) \in \Omega_{\text{cl}}^1(X^{1,9|\mathbf{16}\oplus\overline{\mathbf{16}}}; \mathbf{ICyc}(S^4)). \quad (232)$$

In this manner, the phenomenon will propagate to at least all those supergravity theories that arise as reductions from 11D, rendering them all amenable to global completion by covariant electromagnetic flux quantization.

6.2. M5-Brane Probes

We have seen in (206) that the characteristic L_∞ -fibration for the (non-linearly) self-dual flux density on M5-probes of the 11D bulk is the relative real Whitehead L_∞ -algebra of the quaternionic Hopf fibration. This means that the initial flux quantization of such M5-probes is classified by the actual quaternionic Hopf fibration.

Putting the M5 thus flux-quantized on an A-type orbifold turns out [SS25b; BSS26b] to yield the geometric engineering of topological order which reproduces in fine detail properties of anyons in fractional quantum Hall (FQH) systems (cf. [SS26b; SS25e; SS26a], exposition in [SS25a; SS25c]). Briefly:

- (i) For an M5 worldvolume embedding of the form (cf. [CLS26, p. 5])

$$\begin{aligned} \Sigma^{1,5} &\equiv \mathbb{R}^{1,1} \times \Sigma^2 \times \overset{G}{\mathbb{C}} \\ \downarrow \phi & \\ X^{1,10} &\equiv \mathbb{R}^{1,1} \times \mathbb{R}^5 \times \overset{G}{\mathbb{C}} \times \overset{\overline{G}}{\mathbb{C}} \end{aligned} \quad (233)$$

(where $G \equiv \mathbb{Z}/n \subset \text{SU}(2)$, $n \geq 2$, and \overline{G} its opposite).

- (ii) For flux quantization by the *twistor fibration* with its canonical G -action on one of the two \mathbb{C}^2 -factors:

$$\begin{array}{ccc} \overset{G}{\mathbb{C}P^3} & \xrightarrow{t_{\mathbb{H}}} & \overset{G}{S^4} \\ \parallel & & \parallel \\ (\mathbb{C}^2 \times \overset{G}{\mathbb{C}^2 - \{0\}}) / \mathbb{C}^\times & \longrightarrow & (\mathbb{C}^2 \times \overset{G}{\mathbb{C}^2 - \{0\}}) / \mathbb{H}^\times. \end{array} \quad (234)$$

For trivial background C-field charge, the resulting classifying space on the orbi-singularity $(\Sigma^{1,5})^G \simeq \mathbb{R}^{1,1} \times \Sigma^2$ turns out to be the 2-sphere $S^2 \simeq (\mathbb{C}P^3)^G$, so that the solitonic charges in this situation are equivalently classified by pointed maps

$$\mathbb{R}_{\cup\{\infty\}}^1 \wedge \Sigma_{\cup\{\infty\}}^2 \dashrightarrow S^2. \quad (235)$$

Therefore, the light-cone quantum algebra of the realistic solitonic topological observables (194) and (196) on such M5 brane probes is the group algebra

$$\text{SolTopObs}_0(\mathbb{R}^1 \times \Sigma_{\cup\{\infty\}}^2; S^2) \simeq \mathbb{C}[\pi_1 \text{Map}^*(\Sigma_{\cup\{\infty\}}^2, S^2)]. \quad (236)$$

of the fundamental group of the pointed mapping space from the given surface, with the point at infinity adjoined, to the 2-sphere.

This turns out to reproduce in fine detail [SS26f; SS25e; SS25f] the quantum observables of abelian Chern-Simons theory, hence of the topological quantum order of fractional quantum Hall (FQH) systems on Σ^2 (cf. string theoretic conjectures along these lines in [CGK20] and see review in [SS25a]).

Notably:

- (i) For $\Sigma^2 \equiv \mathbb{R}^2$ the plane, one finds ([SS26b; SS25h]) that:

$$\text{SolTopObs}_0(\mathbb{R}^1 \times \mathbb{R}^2; S^2) \simeq \left\{ \text{renormalized Wilson loop observables} \right\}. \quad (237)$$

is exactly the algebra of observables whose pure states detect the *braiding phases* $\zeta = e^{\pi i n / K}$ of anyon worldlines (framed links) with net crossing number (writhe) $n \in \mathbb{Z}$ in the FQH state with filling fraction $\nu = 1/K$.

- (ii) For $\Sigma^2 \equiv T^2$ the torus, one finds ([KSS26, Thm. 3.5] following [Kal25; LT80; Han74]) that

$$\text{SolTopObs}_0(\mathbb{R}^1 \times T^2; S^2) \simeq \mathbb{C}[\text{integer Heisenberg group at level } 2] \quad (238)$$

is the group algebra of the *integer Heisenberg group* at level=2. This is exactly the algebra of quantum observables for FQH anyons on the torus (cf. [Ton16, (5.28)]).

We intentionally do not go into further detail here, not to overburden these lecture notes. Extensive discussion may be found in the references cited above.

A. Basic Background

For reference in the main text, here we briefly recall some relevant basic notions from higher homotopy and higher algebra.

A.1. Higher Lie Algebras

We briefly recall the dual incarnation of finite-type L_∞ -algebras in the guise of their Chevalley-Eilenberg dgc-algebras (cf. [LS93; SSS09, §6.1; FSS23, §4]). This dual formulation is much more concise and transparent than the “direct” one. In particular, it makes immediate the generalization to super- L_∞ -algebras [FSS15b; FSS19, §3; GSS25, §2.1], which in this dual guise are (mis-)named “FDAs” in the supergravity literature (cf. [CDF91; CD25]).

Our ground field is the real numbers, \mathbb{R} . By *graded* we mean \mathbb{Z} -graded (though all constructions considered here happen to be just \mathbb{N} -graded). We say *gc-algebras* for graded-commutative algebras and *dgc-algebras* for differential graded-commutative algebras with differential of degree +1.

The base case of ordinary Lie algebras. For $(\mathfrak{g}, [-, -])$ a finite-dimensional Lie algebra, recall the construction of its *Chevalley-Eilenberg algebra* $\text{CE}(\mathfrak{g})$:

Write $\mathfrak{g}^\vee := \mathfrak{g}^*$ for the linear dual of the underlying vector space and

$$\wedge^\bullet \mathfrak{g}^\vee := \text{Sym}(\mathfrak{g}^\vee[1]) \quad (239)$$

for its *Grassmann algebra*, the free graded-commutative algebra for which $\wedge^n \mathfrak{g}^\vee$ is the subspace of degree n .

A graded derivation d on $\wedge^\bullet \mathfrak{g}^\vee$ is thus determined by its restriction $\wedge^1 \mathfrak{g}^\vee \rightarrow \wedge^\bullet \mathfrak{g}^\vee$. Taking this restriction to be the linear dual of the Lie bracket, $d|_{\wedge^1 \mathfrak{g}^\vee} := [-, -]^*$ hence yields a graded derivation of degree 1. One checks that its nilpotency is equivalent to the Jacobi identity of the bracket:

$$d^2 = 0 \quad \Leftrightarrow \quad \text{Jacobi identity on } [-, -]. \quad (240)$$

The resulting dgc-algebra is the *Chevalley-Eilenberg algebra* of the Lie algebra:

$$\text{CE}(\mathfrak{g}, [-, -]) := (\wedge^\bullet \mathfrak{g}^\vee, d := [-, -]^*). \quad (241)$$

The key observation now is that this construction is a *fully faithful* functor on the category $\text{LieAlg}_{\mathbb{R}}^{\text{fd}}$ of finite-dimensional Lie algebras:

$$\text{LieAlg}_{\mathbb{R}}^{\text{fd}} \xrightarrow{\text{CE}} \text{dgcAlg}_{\mathbb{R}}^{\text{op}}. \quad (242)$$

This means that we may work (op-)equivalently with the dgc-algebras $\text{CE}(\mathfrak{g})$. But for these there is an evident generalization, simply by allowing the vector space \mathfrak{g} to be \mathbb{N} -graded:

The case of L_∞ -algebras. For \mathfrak{g} an \mathbb{N} -graded vector space of finite type (meaning that each degree \mathfrak{g}_n is a finite-dimensional vector space), write \mathfrak{g}^\vee for its degreewise dual vector space, hence with

$$(\mathfrak{g}^\vee)_n := (\mathfrak{g}_n)^* \quad (243)$$

and write

$$\wedge^\bullet \mathfrak{g}^\vee := \text{Sym}(\mathfrak{g}^\vee[1]) \quad (244)$$

for the graded Grassmann gc-algebra, which in low degrees is

$$(\wedge^\bullet \mathfrak{g}^\vee)_1 = \wedge^1 \mathfrak{g}_0^\vee, \quad (\wedge^\bullet \mathfrak{g}^\vee)_2 = \wedge^2 \mathfrak{g}_0^\vee \oplus \wedge^1 \mathfrak{g}_1^\vee, \quad \text{etc.} \quad (245)$$

Then the category of (connective) finite-type L_∞ -algebras is the full subcategory of $\text{dgcAlg}_{\mathbb{R}}^{\text{op}}$ on those whose underlying gc-algebra is of the form (244):

$$L_\infty \text{Alg}_{\mathbb{R}}^{\text{ft}} \xrightarrow{\text{CE}} \text{dgcAlg}_{\mathbb{R}}^{\text{op}}. \quad (246)$$

Unwinding this: Since a differential d on (244) is again determined by its restriction to $\wedge^1 \mathfrak{g}^\vee$, it is given by a sequence of linear duals of n -ary graded-skew symmetric linear maps for $n \in \mathbb{N}$:

$$d|_{\wedge^1 \mathfrak{g}^\vee} = [-]^* + [-, -]^* + [-, -, -]^* + \cdots : \wedge^1 \mathfrak{g}^\vee \rightarrow \wedge^1 \mathfrak{g}^\vee \oplus \wedge^2 \mathfrak{g}^\vee \oplus \wedge^3 \mathfrak{g}^\vee \oplus \cdots, \quad (247)$$

and the nilpotency condition $d^2 = 0$ is hence equivalently a (somewhat complicated looking) generalization of the Jacobi identity on this system of brackets. Such a system of brackets on graded vector spaces \mathfrak{g} that satisfies these conditions is exactly what makes $(\mathfrak{g}, ([-], [-, -], [-, -, -], \cdots))$ an L_∞ -algebra.

Closed L_∞ -Algebra Valued Forms. Consider $X \in \text{SmthMfd}$. Given again an ordinary Lie algebra \mathfrak{g} of finite dimension, it is immediate that the \mathfrak{g} -valued differential 1-forms on X are equivalently gc-algebra homomorphisms out of the dual Grassmann algebra (239), like this:

$$\Omega_{\text{dR}}^1(X; \mathfrak{g}) \simeq \text{Hom}(\wedge^\bullet \mathfrak{g}^\vee, \Omega_{\text{dR}}^\bullet(X)). \quad (248)$$

Moreover, a moment of reflection shows that the *flat* (here we will say: *closed*) \mathfrak{g} -valued forms among these correspond exactly to those gc-homomorphisms which are in fact dgc-homomorphisms out of the Chevalley-Eilenberg algebra (241):

$$\begin{aligned} \Omega_{\text{cl}}^1(X; \mathfrak{g}) &:= \{A \in \Omega_{\text{dR}}^1(X; \mathfrak{g}) \mid dA + \frac{1}{2}[A \wedge A] = 0\} \\ &\simeq \text{Hom}(\text{CE}(\mathfrak{g}), \Omega_{\text{dR}}^\bullet(X)). \end{aligned} \quad (249)$$

Taking this characterization as the definition, it immediately generalizes to the case where \mathfrak{g} is a finite-type L_∞ -algebra. We say that dgc-homomorphisms out of its Chevalley-Eilenberg algebra (246) into the de Rham dgc-algebra are the closed (flat) \mathfrak{g} -valued differential forms:

$$\Omega_{\text{dR}}^1(X; \mathfrak{g}) := \text{Hom}(\text{CE}(\mathfrak{g}), \Omega_{\text{dR}}^\bullet(X)). \quad (250)$$

A.2. Classifying Spaces

We briefly recall the notion of *classifying space*, beginning with its origin in ordinary (non-)abelian cohomology, and then passing to classifying spaces for generalized abelian and nonabelian cohomology.

Looping and Delooping. For $\mathcal{X}, \mathcal{A} \in \text{Grpd}_\infty$ equipped with base points $\infty \in \mathcal{A}$ and $0 \in \mathcal{A}$, their *pointed mapping space* is the homotopy fiber (58) of the point evaluation map on the plain mapping space:

$$\begin{array}{ccc} \mathbf{Map}^*(\mathcal{X}, \mathcal{A}) & \longrightarrow & \mathbf{Map}(\mathcal{X}, \mathcal{A}) \\ \downarrow & \lrcorner & \downarrow \text{ev}_\infty \\ \{0\} & \longrightarrow & \mathcal{A}. \end{array} \quad (251)$$

In particular, the *based loop space* of $(\mathcal{A}, 0)$ is

$$\Omega\mathcal{A} \simeq \mathbf{Map}^*(S^1, \mathcal{A}). \quad (252)$$

Under concatenation and reversal of loops, the based loop space is an ∞ -group (an A_∞ -monoid whose connected components form a group), and on pointed connected ∞ -groupoids this construction is actually an equivalence:

$$\text{Grpd}_{[1, \infty]}^* \begin{array}{c} \xrightarrow{\Omega} \\ \xrightarrow[\sim]{} \\ \xleftarrow{B} \end{array} \text{Grp}(\text{Grpd}_\infty). \quad (253)$$

Here the inverse equivalence $B(-)$ is hence also called the *delooping* operation.

The analogous equivalence holds in every ∞ -topos ([Lur09, Lem. 7.2.2.1; Lur17, §5.2.6] cf. [NSS15, Thm. 2.19]), hence in particular among smooth ∞ -groupoids (§ 2.3.1), where we denote the delooping in boldface:

$$\text{SmthGrpd}_{[1, \infty]}^* \begin{array}{c} \xrightarrow{\Omega} \\ \xrightarrow[\sim]{} \\ \xleftarrow{\mathbf{B}} \end{array} \text{Grp}(\text{SmthGrpd}_\infty). \quad (254)$$

The shape operation (71) preserves ∞ -group structure and deloopings, due to (70) and (78), cf. [SS26e, Prop. 9.1.4].

Traditional Classifying Spaces. For G a Hausdorff, Lie or Fréchet-Lie group, the traditional *classifying space* for G -principal bundles (cf. [RS17, Thm. 3.5.1]) is, as a homotopy

type ([SS26d, Thm. 5.2.13])

$$BG \equiv \int \mathbf{B}G \simeq B \int G, \quad (255)$$

in that for $X \in \text{SmthMfd}$ it classifies isomorphism classes of G -principal bundles and hence classifies ordinary nonabelian cohomology:

$$H^1(X; G) := \text{GPncplBndl}_X / \text{iso} \simeq \pi_0 \mathbf{Map}(X, BG). \quad (256)$$

If here $G \equiv A \in \text{AbGrp}$ is an abelian group, then BA inherits itself ∞ -group structure (2-group structure) and so on recursively, whence for $n \in \mathbb{N}$ we have the *Eilenberg-MacLane space* (cf. [AGP02, §6])

$$B^n A := BB^{n-1} A, \quad (257)$$

traditionally denoted $K(A, n)$, which classifies ordinary abelian cohomology (cf. [AGP02, §7.1; FSS23, Ex. 2.1]):

$$H^n(X; A) \simeq \pi_0 \mathbf{Map}(X, B^n A). \quad (258)$$

Spectra of Classifying Spaces. By construction, the sequence of EM-spaces $(B^n A)_{n \in \mathbb{N}}$ (257) forms a *spectrum of spaces*

$$E_\bullet := \left((E_n \in \text{Grpd}_\infty^*)_{n \in \mathbb{N}}, \text{ with } (E_n \xrightarrow{\sim} \Omega E_{n+1})_{n \in \mathbb{N}} \right). \quad (259)$$

The *Brown representability theorem* says (cf. [Koc96, §3.4; AGP02, Thm. 12.2.18]) that such spectra are exactly the classifying spaces of Whitehead-generalized abelian cohomology theories $E^\bullet(-)$ (cf. [FSS23, Ex. 2.10]):

$$E^n(X) \simeq \pi_0 \mathbf{Map}(X, E_n). \quad (260)$$

A key example beyond ordinary cohomology, the classifying space for *complex topological K-theory* $K^0(-)$ is

$$\text{KU}_0 \simeq \text{BU}(\infty) \times \mathbb{Z} := \varinjlim_n \text{BU}(n) \times \mathbb{Z}. \quad (261)$$

Remarkably, while each finite stage $\text{BU}(n)$ fails to have further delooping, the colimiting space does deloop, by the homotopy-theoretic version of *Bott periodicity* (cf. [AGP02, §B.2; Hus+08, §15]):

$$\text{BU}(\infty) \times \mathbb{Z} \xrightarrow{\sim} \text{U}(\infty) \xrightarrow{\sim} \text{BU}(\infty) \times \mathbb{Z} \xrightarrow{\sim} \dots \quad (262)$$

Hence

$$\text{KU}_\bullet := \begin{cases} \text{BU}(\infty) \times \mathbb{Z} & \text{if } \bullet \text{ is even} \\ \text{U}(\infty) & \text{if } \bullet \text{ is odd} \end{cases} \quad (263)$$

makes a spectrum of spaces, which exhibits K-theory as a Whitehead-generalized abelian cohomology theory.

Unstable Classifying Spaces. In this vein, one may understand the finite/unstable stages $\text{BU}(n)$ as classifying forms of *unstable K-theory*.

In fact, since every connected space \mathcal{A} is the classifying space of its loop ∞ -group (253)

$$\mathcal{A} \simeq B(\Omega \mathcal{A}), \quad (264)$$

every connected space is the classifying space of the evident generalization of ordinary non-abelian cohomology (256):

$$H^1(X; \Omega \mathcal{A}) := \pi_0 \mathbf{Map}(X, \mathcal{A}). \quad (265)$$

Moreover, here the connectedness assumption only serves to give the cohomology in degree 1. More generally we may say that *nonabelian cohomology* is what is classified by *any* (pointed) $\mathcal{A} \in \text{Grpd}_\infty$:

$$H^0(X; \mathcal{A}) := \pi_0 \mathbf{Map}(X, \mathcal{A}) \quad (266)$$

([Toë02, Def. 6.0.6; Lur14, Def. 6; FSS23, §2], more exposition in [SS25i, §1; SS26e, §4]).

In this pattern, yet more generally we have nonabelian cohomology in every negative degree $n \in \mathbb{N}$

$$H^{-n}(X; \mathcal{A}) := \pi_0 \mathbf{Map}(X, \Omega^n \mathcal{A}), \quad (267)$$

while the existence of nonabelian cohomology in positive degrees $n \in \mathbb{N}$ is conditional on higher deloopings

$$H^n(\Omega \mathcal{A}) := \pi_0 \mathbf{Map}(X; B^n \Omega \mathcal{A}) \quad (268)$$

which need not exist — and in examples rarely exists for finite n without already existing for all n and thus reducing to the stable case (259).

The most basic example of an unstable classifying space is the n -sphere for $n \geq 2$. The corresponding nonabelian cohomology theory is *n-Cohomotopy*:

$$\pi^n(X) := \pi_0 \mathbf{Map}(X, \mathbb{S}^n). \quad (269)$$

Fibrations of Classifying Spaces. Unstable classifying spaces appear already when *twisting* abelian cohomology theories.

For instance the K-theory classifying space KU_0 (261) carries an ∞ -action by $\text{BU}(1)$, whose homotopy quotient is a KU_0 -fibration over $B^2\text{U}(1)$:

$$\begin{array}{ccc} \text{KU}_0 & \longrightarrow & \text{KU}_0 // \text{BU}(1) \\ \downarrow & \lrcorner & \downarrow p \\ * & \longrightarrow & B^2\text{U}(1). \end{array} \quad (270)$$

Hence, while the homotopy quotient is not itself a stage in a spectrum, it is so *fiberwise* over $B^2\text{U}(1)$. Accordingly, the *fiberwise* maps to this quotient, over a fixed *twisting* map $\tau : X \rightarrow B^2\text{U}(1)$, classify τ -*twisted K-theory* ([AS04], cf. [FSS23, Ex. 3.4; SS26e, Def. 5.1.22; SS26d, Ex. 6.2.5]):

$$K^\tau(X) := \pi_0 \mathbf{Map}\left((X, \tau), (\text{KU}_0 // \text{BU}(1), p)\right)_{B^2\text{U}(1)} = \pi_0 \left\{ \begin{array}{ccc} & \text{KU}_0 // \text{BU}(1) & \\ & \nearrow & \downarrow p \\ X & \xrightarrow{\tau} & B^2\text{U}(1) \end{array} \right\}. \quad (271)$$

Cyclic Loop Spaces. A particularly important operation for obtaining new unstable classifying spaces from given spaces \mathcal{A} is *cyclification*.

Given $\mathcal{G} \in \text{Grp}(\text{Grpd}_\infty)$, there is an induced \mathcal{G} -action on $\mathbf{Map}(\mathcal{G}, \mathcal{A})$. The homotopy quotient by that action turns out to be the classifying space for \mathcal{A} -cohomology on \mathcal{G} -principal fibrations

$$\begin{array}{ccc} \mathcal{P} & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ X & \xrightarrow{c} & B\mathcal{G} \end{array} \quad (272)$$

in that we have a natural equivalence ([BSS19, §2.2; SS24a, §2.2])

$$\left\{ \mathcal{P} \dashrightarrow \mathcal{A} \right\} \begin{array}{c} \xrightarrow{\text{reduction}} \\ \xleftarrow{\sim} \\ \xleftarrow{\text{oxidation}} \end{array} \left\{ X \begin{array}{c} \dashrightarrow \mathbf{Map}(\mathcal{G}, \mathcal{A}) // \mathcal{G} \\ \xrightarrow{c} B\mathcal{G} \end{array} \right\}. \quad (273)$$

In particular, when $\mathcal{G} \equiv \int S^1$ is the circle, then \mathcal{P} is (the homotopy type of) a circle-principal bundle and this homotopy quotient is the *cyclic loop space*

$$\mathrm{Cyc}(\mathcal{A}) := \mathbf{Map}(S^1, \mathcal{A}) // \int S^1. \quad (274)$$

But if we think of \mathcal{A} here as classifying charges of a field theory on \mathcal{P} , then this equivalence (272) says that $\mathrm{Cyc}(\mathcal{A})$ classifies the same information, but now transmuted to a larger set of charges on the base space X . This is the global topological guise of what in physics is known as *double dimensional reduction* of fluxes (cf. [FSS18, §3; GSS25, §2.2]).

References

- [AF25] D. Ayala and J. Francis. *A parametrized Pontryagin–Thom theorem*. 2025. DOI: [10.48550/arXiv.2512.10274](#). arXiv: [2512.10274](#) [math.AT] (cit. on p. 25).
- [AGP02] M. Aguilar, S. Gitler, and C. Prieto. *Algebraic Topology from a Homotopical Viewpoint*. Universitext. Springer, 2002. ISBN: 9780387224893. DOI: [10.1007/b97586](#) (cit. on pp. 27, 46).
- [Alf25] L. Alfonsi. “Higher Geometry in Physics”. In: *Encyclopedia of Mathematical Physics*. Ed. by R. Szabo and M. Bojowald. 2nd ed. Academic Press, 2025, pp. 39–61. ISBN: 978-0-323-95706-9. DOI: [10.1016/b978-0-323-95703-8.00209-3](#). arXiv: [2312.07308](#) [hep-th] (cit. on pp. 3–5, 23).
- [Alv85] O. Alvarez. “Topological Quantization and Cohomology”. In: *Commun. Math. Phys.* 100.2 (June 1985), pp. 279–309. DOI: [10.1007/bf01212452](#) (cit. on pp. 3, 31, 32).
- [AS04] M. Atiyah and G. Segal. “Twisted K-theory”. In: *Ukrainian Mathematical Bulletin* 1.3 (2004), pp. 291–330. eprint: [math/0407054](#) (math.KT). URL: <http://iamm.su/en/journals/j879/?VID=10> (cit. on p. 47).
- [Awo10] S. Awodey. *Category Theory*. 2nd. Vol. 52. Oxford Logic Guides. Oxford: Oxford University Press, 2010. ISBN: 978-0-19-923718-0. DOI: [10.1093/acprof:oso/9780198568612.001.0001](#) (cit. on p. 5).
- [Bag19] M. Baggioli. *Applied Holography – A Practical Mini-Course*. SpringerBriefs in Physics. Springer, 2019. DOI: [10.1007/978-3-030-35184-7](#) (cit. on p. 5).
- [Ban25] P. Banerjee. “Gauge potentials on the M5 brane in twisted equivariant cohomotopy”. In: *Journal of High Energy Physics* 2025.12 (2025), p. 162. DOI: [10.1007/JHEP12\(2025\)162](#). arXiv: [2507.07049](#) [hep-th] (cit. on p. 32).
- [BBP24] D. Berwick-Evans, P. Boavida de Brito, and D. Pavlov. “Classifying spaces of ∞ -sheaves”. In: *Algebraic & Geometric Topology* 24.9 (2024), pp. 4891–4937. DOI: [10.2140/agt.2024.24.4891](#). arXiv: [1912.10544](#) [math.AT] (cit. on pp. 16, 17).
- [Bec+17] C. Becker et al. “Abelian Duality on Globally Hyperbolic Spacetimes”. In: *Communications in Mathematical Physics* 349.1 (May 2017), pp. 361–392. ISSN: 1432-0916. DOI: [10.1007/s00220-016-2669-9](#). arXiv: [1511.00316](#) [hep-th] (cit. on p. 32).
- [Ber15] A. Berglund. “Rational homotopy theory of mapping spaces via Lie theory for L_∞ algebras”. In: *Homology, Homotopy and Applications* 17.2 (2015). DOI: [10.4310/HHA.2015.v17.n2.a16](#). arXiv: [1110.6145](#) [math.AT] (cit. on p. 25).
- [BFM11] U. Buijs, Y. Félix, and A. Murillo. “ L_∞ -models of based mapping spaces”. In: *J. Math. Soc. Japan* 63.2 (2011), pp. 503–524. DOI: [10.2969/jmsj/06320503](#). arXiv: [1209.4756](#) [math.AT] (cit. on p. 25).

- [BFS08] L. Bonora, F. Ferrari Ruffino, and R. Savelli. “Classifying A-field and B-field configurations in the presence of D-branes”. In: *JHEP* 12 (2008), p. 078. DOI: [10.1088/1126-6708/2008/12/078](https://doi.org/10.1088/1126-6708/2008/12/078). arXiv: [0810.4291](https://arxiv.org/abs/0810.4291) [hep-th] (cit. on p. 33).
- [BH11] J. C. Baez and A. E. Hoffnung. “Convenient Categories of Smooth Spaces”. In: *Transactions of the American Mathematical Society* 363.11 (2011), pp. 5789–5825. DOI: [10.1090/S0002-9947-2011-05107-X](https://doi.org/10.1090/S0002-9947-2011-05107-X). arXiv: [0807.1704](https://arxiv.org/abs/0807.1704) [math.DG] (cit. on p. 8).
- [BH25] H. Bursztyn and M. del Hoyo. “Lie Groupoids”. In: *Encyclopedia of Mathematical Physics*. 2nd. Elsevier, 2025, p. 469. DOI: [10.1016/B978-0-323-95703-8.00024-0](https://doi.org/10.1016/B978-0-323-95703-8.00024-0). arXiv: [2309.14105](https://arxiv.org/abs/2309.14105) [math.DG] (cit. on p. 4).
- [BH80] L. Brink and P. Howe. “Eleven-Dimensional Supergravity on the Mass-Shell in Superspace”. In: *Phys. Lett. B* 91 (1980), pp. 384–386. DOI: [10.1016/0370-2693\(80\)91002-3](https://doi.org/10.1016/0370-2693(80)91002-3) (cit. on p. 41).
- [Bor+25] L. Borsten et al. “Higher Gauge Theory”. In: *Encyclopedia of Mathematical Physics*. 2nd. Vol. 4. 2025, pp. 159–185. DOI: [10.1016/B978-0-323-95703-8.00217-2](https://doi.org/10.1016/B978-0-323-95703-8.00217-2). arXiv: [2401.05275](https://arxiv.org/abs/2401.05275) [hep-th] (cit. on p. 5, 23).
- [Bor94] F. Borceux. *Handbook of Categorical Algebra: Volume 1, Basic Category Theory*. Vol. 50. Encyclopedia of Mathematics and its Applications. Cambridge: Cambridge University Press, 1994. DOI: [10.1017/CB09780511525858](https://doi.org/10.1017/CB09780511525858) (cit. on p. 10).
- [Bro73] K. S. Brown. “Abstract homotopy theory and generalized sheaf cohomology”. In: *Transactions of the American Mathematical Society* 186 (1973), pp. 419–458. DOI: [10.2307/1996573](https://doi.org/10.2307/1996573) (cit. on p. 5).
- [Bry+91] R. L. Bryant et al. *Exterior Differential Systems*. Vol. 18. Mathematical Sciences Research Institute Publications. New York, NY: Springer-Verlag, 1991. ISBN: 978-1-4613-9716-8. DOI: [10.1007/978-1-4613-9714-4](https://doi.org/10.1007/978-1-4613-9714-4) (cit. on p. 21).
- [Bry93] J.-L. Brylinski. *Loop Spaces, Characteristic Classes and Geometric Quantization*. Vol. 107. Progress in Mathematics. Boston, MA: Birkhäuser, 1993. ISBN: 978-0-8176-3644-9. DOI: [10.1007/978-0-8176-4731-5](https://doi.org/10.1007/978-0-8176-4731-5) (cit. on p. 3, 32).
- [BS07] J. C. Baez and U. Schreiber. “Higher Gauge Theory”. In: *Categories in Algebra, Geometry and Mathematical Physics*. Vol. 431. Contemporary Mathematics. American Mathematical Society, 2007, pp. 7–30. DOI: [10.1090/conm/431/08270](https://doi.org/10.1090/conm/431/08270). arXiv: [math/0511710](https://arxiv.org/abs/math/0511710) [math.DG] (cit. on p. 5, 23).
- [BSS18] M. Benini, A. Schenkel, and U. Schreiber. “The Stack of Yang–Mills Fields on Lorentzian Manifolds”. In: *Communications in Mathematical Physics* 359.2 (Mar. 2018), pp. 765–820. DOI: [10.1007/s00220-018-3120-1](https://doi.org/10.1007/s00220-018-3120-1). arXiv: [1704.01378](https://arxiv.org/abs/1704.01378) [math-ph] (cit. on p. 3).
- [BSS19] V. Braunack-Mayer, H. Sati, and U. Schreiber. “Gauge enhancement of Super M-Branes via rational parameterized stable homotopy theory”. In: *Communications in Mathematical Physics* 371 (2019), pp. 197–265. DOI: [10.1007/s00220-019-03441-4](https://doi.org/10.1007/s00220-019-03441-4). arXiv: [1806.01115](https://arxiv.org/abs/1806.01115) [hep-th] (cit. on p. 39, 48).
- [BSS26a] P. Banerjee, H. Sati, and U. Schreiber. *Flux Quantization of Type IIA in Unstable K-Theory*. to appear next week. 2026. URL: <https://ncatlab.org/schreiber/show/Flux+Quantization+of+Type+IIA> (cit. on p. 40).
- [BSS26b] P. Banerjee, H. Sati, and U. Schreiber. “Flux Quantization on M-Strings”. In: *Journal of Physics A: Mathematical and Theoretical* (June 2026). arXiv: [2603.14440](https://arxiv.org/abs/2603.14440) [hep-th] (cit. on p. 42).
- [Bun+26] S. Bunk et al. *Symmetries and Higher-Form Connections in Derived Differential Geometry*. 2026. DOI: [10.48550/arXiv.2602.03441](https://doi.org/10.48550/arXiv.2602.03441). arXiv: [2602.03441](https://arxiv.org/abs/2602.03441) [math.DG] (cit. on p. 5, 23).
- [Bun12] U. Bunke. “Differential cohomology”. In: (2012). Course notes. arXiv: [1208.3961](https://arxiv.org/abs/1208.3961) [math.AT] (cit. on p. 31).
- [CD25] L. Castellani and R. D’Auria. “The L_∞ -structure of Free Differential Algebras and Supergravity”. In: *International Journal of Modern Physics A* (2025). DOI: [10.1142/S0217751X25501787](https://doi.org/10.1142/S0217751X25501787). arXiv: [2507.20344](https://arxiv.org/abs/2507.20344) [hep-th] (cit. on p. 43).

- [CDF91] L. Castellani, R. D’Auria, and P. Fré. *Supergravity and Superstrings – A Geometric Perspective*. World Scientific, 1991. DOI: [10.1142/0224](https://doi.org/10.1142/0224) (cit. on pp. 41, 43).
- [CF80] E. Cremmer and S. Ferrara. “Formulation of Eleven-Dimensional Supergravity in Super-space”. In: *Phys. Lett. B* 91 (1980), pp. 61–66. DOI: [10.1016/0370-2693\(80\)90662-0](https://doi.org/10.1016/0370-2693(80)90662-0) (cit. on p. 41).
- [CGK20] G. Y. Cho, D. Gang, and H.-C. Kim. “M-theoretic Genesis of Topological Phases”. In: *Journal of High Energy Physics* 2020.11 (2020), p. 115. DOI: [10.1007/JHEP11\(2020\)115](https://doi.org/10.1007/JHEP11(2020)115). arXiv: [2007.01532](https://arxiv.org/abs/2007.01532) [[hep-th](#)] (cit. on p. 43).
- [CGX11] A. S. Cattaneo, A. Giaquinto, and P. Xu, eds. *Higher Structures in Geometry and Physics: In Honor of Murray Gerstenhaber and Jim Stasheff*. Vol. 287. Progress in Mathematics. Boston, MA: Birkhäuser, 2011. DOI: [10.1007/978-0-8176-4735-3](https://doi.org/10.1007/978-0-8176-4735-3) (cit. on p. 4).
- [CJM04] A. Carey, S. Johnson, and M. Murray. “Holonomy on D-Branes”. In: *Journal of Geometry and Physics* 52.2 (2004), pp. 186–216. DOI: [10.1016/j.geomphys.2004.02.008](https://doi.org/10.1016/j.geomphys.2004.02.008). arXiv: [hep-th/0204199](https://arxiv.org/abs/hep-th/0204199) (cit. on p. 33).
- [CL94] A. Candiello and K. Lechner. “Duality in supergravity theories”. In: *Nuclear Physics B* 412 (1994), pp. 479–501. DOI: [10.1016/0550-3213\(94\)90389-1](https://doi.org/10.1016/0550-3213(94)90389-1). arXiv: [hep-th/9309143](https://arxiv.org/abs/hep-th/9309143) [[hep-th](#)] (cit. on p. 41).
- [CLS26] C. Couzens, A. Lüscher, and J. Sparks. “Localizing punctures in M-theory”. In: *J. High Energy Phys.* 2026.2 (2026), p. 180. DOI: [10.1007/JHEP02\(2026\)180](https://doi.org/10.1007/JHEP02(2026)180). arXiv: [2511.03397](https://arxiv.org/abs/2511.03397) [[hep-th](#)] (cit. on p. 42).
- [DGP13] A. Donos, J. P. Gauntlett, and C. Pantelidou. “Semi-local quantum criticality in string/M-theory”. In: *J. High Energy Phys.* 2013.3 (2013), p. 103. DOI: [10.1007/JHEP03\(2013\)103](https://doi.org/10.1007/JHEP03(2013)103). arXiv: [1212.1462](https://arxiv.org/abs/1212.1462) [[hep-th](#)] (cit. on p. 5).
- [Dir31] P. A. M. Dirac. “Quantised Singularities in the Electromagnetic Field”. In: *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character* 133.821 (Sept. 1931), pp. 60–72. DOI: [10.1098/rspa.1931.0130](https://doi.org/10.1098/rspa.1931.0130) (cit. on pp. 3, 24, 32).
- [Dir49] P. A. M. Dirac. “Forms of Relativistic Dynamics”. In: *Reviews of Modern Physics* 21.3 (July 1949), pp. 392–399. DOI: [10.1103/RevModPhys.21.392](https://doi.org/10.1103/RevModPhys.21.392). URL: <https://doi.org/10.1103/RevModPhys.21.392> (cit. on p. 35).
- [Don+13] A. Donos et al. “Competing orders in M-theory: superfluids, stripes and metamagnetism”. In: *J. High Energy Phys.* 2013.3 (2013), p. 108. DOI: [10.1007/JHEP03\(2013\)108](https://doi.org/10.1007/JHEP03(2013)108). arXiv: [1212.0871](https://arxiv.org/abs/1212.0871) [[hep-th](#)] (cit. on p. 5).
- [Duf99] M. J. Duff. *The World in Eleven Dimensions: Supergravity, Supermembranes and M-Theory*. IOP Publishing, 1999. ISBN: 978-0-750-30672-0. DOI: [10.1201/9781482268737](https://doi.org/10.1201/9781482268737) (cit. on pp. 3, 5).
- [Fer12] F. Ferrari Ruffino. “Classifying A-field and B-field configurations in the presence of D-branes - Part II: Stacks of D-branes”. In: *Nucl. Phys. B* 858 (2012), pp. 377–404. DOI: [10.1016/j.nuclphysb.2012.01.013](https://doi.org/10.1016/j.nuclphysb.2012.01.013). arXiv: [1104.2798](https://arxiv.org/abs/1104.2798) [[hep-th](#)] (cit. on p. 33).
- [FHT00] Y. Félix, S. Halperin, and J.-C. Thomas. *Rational Homotopy Theory*. Vol. 205. Graduate Texts in Mathematics. Springer, 2000. DOI: [10.1007/978-1-4613-0105-9](https://doi.org/10.1007/978-1-4613-0105-9) (cit. on p. 24).
- [FMS07a] D. S. Freed, G. W. Moore, and G. Segal. “Heisenberg Groups and Noncommutative Fluxes”. In: *Annals Phys.* 322 (2007), pp. 236–285. DOI: [10.1016/j.aop.2006.07.014](https://doi.org/10.1016/j.aop.2006.07.014). arXiv: [hep-th/0605200](https://arxiv.org/abs/hep-th/0605200) (cit. on p. 32).
- [FMS07b] D. S. Freed, G. W. Moore, and G. Segal. “The Uncertainty of Fluxes”. In: *Communications in Mathematical Physics* 271.1 (Jan. 2007), pp. 247–274. ISSN: 1432-0916. DOI: [10.1007/s00220-006-0181-3](https://doi.org/10.1007/s00220-006-0181-3). arXiv: [hep-th/0605198](https://arxiv.org/abs/hep-th/0605198) [[hep-th](#)] (cit. on p. 32).
- [Fra11] T. Frankel. *The Geometry of Physics: An Introduction*. Cambridge University Press, 2011. ISBN: 9781139061377. DOI: [10.1017/CB09781139061377](https://doi.org/10.1017/CB09781139061377) (cit. on pp. 3, 32).
- [Fre01] D. S. Freed. “K-Theory in Quantum Field Theory”. In: *Current Developments in Mathematics, 2001*. Vol. 2001. Current Developments in Mathematics 1. International Press

- of Boston, 2001, pp. 41–87. DOI: [10.4310/cdm.2001.v2001.n1.a2](https://doi.org/10.4310/cdm.2001.v2001.n1.a2). arXiv: [math-ph/0206031](https://arxiv.org/abs/math-ph/0206031) (cit. on p. 33).
- [Fre02] D. S. Freed. “Dirac Charge Quantization and Generalized Differential Cohomology”. In: *Surveys in Differential Geometry* 7.1 (2002), pp. 129–194. DOI: [10.4310/SDG.2002.v7.n1.a6](https://doi.org/10.4310/SDG.2002.v7.n1.a6). arXiv: [hep-th/0011220](https://arxiv.org/abs/hep-th/0011220) [[hep-th](#)] (cit. on pp. 31–33).
- [Fri12] G. Friedman. “An elementary illustrated introduction to simplicial sets”. In: *Rocky Mountain Journal of Mathematics* 42.2 (2012), pp. 353–423. DOI: [10.1216/RMJ-2012-42-2-353](https://doi.org/10.1216/RMJ-2012-42-2-353). arXiv: [0809.4221](https://arxiv.org/abs/0809.4221) [[math.AT](#)] (cit. on p. 12).
- [FRS16] D. Fiorenza, C. Rogers, and U. Schreiber. “Higher $U(1)$ -gerbe connections in geometric prequantization”. In: *Reviews in Mathematical Physics* 28.06 (2016), p. 1650012. DOI: [10.1142/S0129055X16500124](https://doi.org/10.1142/S0129055X16500124). arXiv: [1304.0236](https://arxiv.org/abs/1304.0236) [[math.DG](#)] (cit. on p. 31).
- [FSS12] D. Fiorenza, U. Schreiber, and J. Stasheff. “Čech cocycles for differential characteristic classes: an ∞ -Lie theoretic construction”. In: *Advances in Theoretical and Mathematical Physics* 16.1 (2012), pp. 149–250. DOI: [10.4310/atmp.2012.v16.n1.a5](https://doi.org/10.4310/atmp.2012.v16.n1.a5). arXiv: [1011.4735](https://arxiv.org/abs/1011.4735) [[math.AT](#)] (cit. on pp. 5, 23).
- [FSS13] D. Fiorenza, H. Sati, and U. Schreiber. “Extended higher cup-product Chern-Simons theories”. In: *Journal of Geometry and Physics* 74 (2013), pp. 130–163. DOI: [10.1016/j.geomphys.2013.07.011](https://doi.org/10.1016/j.geomphys.2013.07.011). arXiv: [1207.5449](https://arxiv.org/abs/1207.5449) [[hep-th](#)] (cit. on p. 31).
- [FSS15a] D. Fiorenza, H. Sati, and U. Schreiber. “A Higher Stacky Perspective on Chern–Simons Theory”. In: *Mathematical Aspects of Quantum Field Theories*. Ed. by D. Calaque and T. Strobl. Springer, 2015, pp. 153–211. ISBN: 9783319099491. DOI: [10.1007/978-3-319-09949-1_6](https://doi.org/10.1007/978-3-319-09949-1_6). arXiv: [1301.2580](https://arxiv.org/abs/1301.2580) [[hep-th](#)] (cit. on pp. 4, 31).
- [FSS15b] D. Fiorenza, H. Sati, and U. Schreiber. “Super Lie n -algebra extensions, higher WZW models, and super p -branes with tensor multiplet fields”. In: *International Journal of Geometric Methods in Modern Physics* 12.02 (2015), p. 1550018. DOI: [10.1142/S021987815500188](https://doi.org/10.1142/S021987815500188). arXiv: [1308.5264](https://arxiv.org/abs/1308.5264) [[hep-th](#)] (cit. on p. 43).
- [FSS15c] D. Fiorenza, H. Sati, and U. Schreiber. “The WZW term of the M5-brane and differential cohomotopy”. In: *J. Math. Phys.* 56.10 (2015), p. 102301. DOI: [10.1063/1.4932618](https://doi.org/10.1063/1.4932618). arXiv: [1506.07557](https://arxiv.org/abs/1506.07557) [[math-ph](#)] (cit. on pp. 32, 33).
- [FSS17] D. Fiorenza, H. Sati, and U. Schreiber. “Rational sphere valued supercocycles in M-theory and type IIA string theory”. In: *Journal of Geometry and Physics* 114 (Apr. 2017), pp. 91–108. DOI: [10.1016/j.geomphys.2016.11.024](https://doi.org/10.1016/j.geomphys.2016.11.024). arXiv: [1606.03206](https://arxiv.org/abs/1606.03206) [[hep-th](#)] (cit. on p. 39).
- [FSS18] D. Fiorenza, H. Sati, and U. Schreiber. “T-Duality from super Lie n -algebra cocycles for super p -branes”. In: *Adv. Theor. Math. Phys.* 22.5 (2018), pp. 1209–1270. DOI: [10.4310/ATMP.2018.v22.n5.a3](https://doi.org/10.4310/ATMP.2018.v22.n5.a3). arXiv: [1611.06536](https://arxiv.org/abs/1611.06536) [[hep-th](#)] (cit. on p. 48).
- [FSS19] D. Fiorenza, H. Sati, and U. Schreiber. “The Rational Higher Structure of M-theory”. In: *Fortschritte der Physik* 67.8–9 (2019), p. 1910017. DOI: [10.1002/prop.201910017](https://doi.org/10.1002/prop.201910017). arXiv: [1903.02834](https://arxiv.org/abs/1903.02834) [[hep-th](#)] (cit. on pp. 4, 5, 43).
- [FSS20] D. Fiorenza, H. Sati, and U. Schreiber. “Twisted Cohomotopy Implies M-Theory Anomaly Cancellation on 8-Manifolds”. In: *Commun. Math. Phys.* 377.3 (Apr. 2020), pp. 1961–2025. DOI: [10.1007/s00220-020-03707-2](https://doi.org/10.1007/s00220-020-03707-2). arXiv: [1904.10207](https://arxiv.org/abs/1904.10207) [[hep-th](#)] (cit. on pp. 3, 39, 41).
- [FSS21a] D. Fiorenza, H. Sati, and U. Schreiber. “Twisted Cohomotopy Implies Level Quantization of the Full 6d Wess-Zumino Term of the M5-Brane”. In: *Commun. Math. Phys.* 384.1 (Apr. 2021), pp. 403–432. DOI: [10.1007/s00220-021-03951-0](https://doi.org/10.1007/s00220-021-03951-0). arXiv: [1906.07417](https://arxiv.org/abs/1906.07417) [[hep-th](#)] (cit. on pp. 3, 39, 41).
- [FSS21b] D. Fiorenza, H. Sati, and U. Schreiber. “Twisted Cohomotopy implies Twisted String Structure on M5-branes”. In: *Journal of Mathematical Physics* 62.4 (2021), p. 042301. DOI: [10.1063/5.0037786](https://doi.org/10.1063/5.0037786). arXiv: [2002.11093](https://arxiv.org/abs/2002.11093) [[hep-th](#)] (cit. on p. 41).
- [FSS23] D. Fiorenza, H. Sati, and U. Schreiber. *The Character Map in Non-abelian Cohomology: Twisted, Differential, and Generalized*. <https://ncatlab.org/schreiber/show>

- w/[The+Character+Map+in+Non-Abelian+Cohomology](#). World Scientific, 2023. ISBN: 9789811276705. DOI: [10.1142/13422](#) (cit. on pp. 4, 14, 23–25, 27, 28, 31–33, 38, 39, 43, 46, 47).
- [FW99] D. S. Freed and E. Witten. “Anomalies in String Theory with D-Branes”. In: *Asian Journal of Mathematics* 3.4 (Dec. 1999), pp. 819–852. DOI: [10.4310/AJM.1999.v3.n4.a6](#). arXiv: [hep-th/9907189](#) [[hep-th](#)] (cit. on pp. 31, 33).
- [Gaw88] K. Gawedzki. “Topological Actions in Two-Dimensional Quantum Field Theories”. In: *Nonperturbative Quantum Field Theory*. Ed. by G. ’t Hooft et al. Vol. 185. NATO ASI Series. New York, NY: Springer, 1988, pp. 101–141. DOI: [10.1007/978-1-4613-0729-7_5](#) (cit. on pp. 31, 33).
- [GHM97] M. B. Green, J. A. Harvey, and G. W. Moore. “I-brane inflow and anomalous couplings on D-branes”. In: *Class. Quant. Grav.* 14 (1997), pp. 47–52. DOI: [10.1088/0264-9381/14/1/008](#). arXiv: [hep-th/9605033](#) [[hep-th](#)] (cit. on p. 33).
- [GJ09] P. Goerss and J. F. Jardine. *Simplicial homotopy theory*. Modern Birkhäuser Classics. Reprint of the 1999 edition (Progress in Mathematics, Vol. 174). Birkhäuser, 2009. DOI: [10.1007/978-3-0346-0189-4](#) (cit. on p. 17).
- [GM13] P. Griffiths and J. Morgan. *Rational Homotopy Theory and Differential Forms*. Vol. 16. Progress in Mathematics. Boston, MA: Birkhäuser, 2013. ISBN: 978-1-4614-8467-7. DOI: [10.1007/978-1-4614-8468-4](#) (cit. on p. 25).
- [GMS09] G. Giachetta, L. Mangiarotti, and G. Sardanashvily. *Advanced Classical Field Theory*. Singapore: World Scientific, 2009. ISBN: 978-981-283-914-5. DOI: [10.1142/7189](#) (cit. on p. 3).
- [GPR10] S. S. Gubser, S. S. Pufu, and F. D. Rocha. “Quantum critical superconductors in string theory and M-theory”. In: *Phys. Lett. B* 683 (2010), pp. 201–204. DOI: [10.1016/j.physletb.2009.12.017](#) (cit. on p. 5).
- [GR02] K. Gawedzki and N. Reis. “WZW Branes and Gerbes”. In: *Rev. Math. Phys.* 14.12 (2002), pp. 1281–1334. DOI: [10.1142/S0129055X02001557](#). arXiv: [hep-th/0205233](#) (cit. on p. 33).
- [GS19] D. Grady and H. Sati. “Higher-twisted periodic smooth Deligne cohomology”. In: *Homology, Homotopy and Applications* 21.1 (2019), pp. 129–159. ISSN: 1532-0081. DOI: [10.4310/hha.2019.v21.n1.a7](#). URL: [http://dx.doi.org/10.4310/HHA.2019.v21.n1.a7](#) (cit. on p. 33).
- [GS21] D. Grady and H. Sati. “Differential cohomotopy versus differential cohomology for M-theory and differential lifts of Postnikov towers”. In: *Journal of Geometry and Physics* 165 (2021), p. 104203. DOI: [10.1016/j.geomphys.2021.104203](#). arXiv: [2001.07640](#) [[hep-th](#)] (cit. on pp. 32, 33).
- [GS22] D. Grady and H. Sati. “Ramond–Ramond fields and twisted differential K-theory”. In: *Advances in Theoretical and Mathematical Physics* 26.5 (2022), pp. 1097–1155. ISSN: 1095-0753. DOI: [10.4310/atmp.2022.v26.n5.a2](#). URL: [http://dx.doi.org/10.4310/ATMP.2022.v26.n5.a2](#) (cit. on pp. 31, 33).
- [GS25a] G. Giotopoulos and H. Sati. “Field Theory via Higher Geometry I: Smooth Sets of Fields”. In: *Journal of Geometry and Physics* 213 (July 2025), p. 105462. DOI: [10.1016/j.geomphys.2025.105462](#). arXiv: [2312.16301](#) [[math-ph](#)] (cit. on pp. 4, 7).
- [GS25b] G. Giotopoulos and H. Sati. “Field Theory via Higher Geometry II: Thickened Smooth Sets as Synthetic Foundations”. In: (2025). arXiv:2512.22816 [[math-ph](#)]. arXiv: [2512.22816](#) [[math-ph](#)] (cit. on pp. 4, 7).
- [GS26] G. Giotopoulos and H. Sati. “Flux Quantization on 10D IIA Superspace Via Cyclification from 11D”. in preparation. 2026 (cit. on p. 42).
- [GSS24a] G. Giotopoulos, H. Sati, and U. Schreiber. “Flux quantization on 11D superspace”. In: *Journal of High Energy Physics* 2024.7 (July 2024), p. 82. DOI: [10.1007/JHEP07\(2024\)082](#). arXiv: [2403.16456](#) [[hep-th](#)] (cit. on pp. 3, 32, 39–41).

- [GSS24b] G. Giotopoulos, H. Sati, and U. Schreiber. “Flux Quantization on M5-Branes”. In: *Journal of High Energy Physics* 2024.10 (Oct. 2024), p. 140. DOI: [10.1007/JHEP10\(2024\)140](https://doi.org/10.1007/JHEP10(2024)140). arXiv: [2406.11304](https://arxiv.org/abs/2406.11304) [hep-th] (cit. on p. 32).
- [GSS25] G. Giotopoulos, H. Sati, and U. Schreiber. “Super-Lie $_{\infty}$ T-Duality and M-Theory”. In: *Rev. Math. Phys.* (2025). DOI: [10.1142/S0129055X25500187](https://doi.org/10.1142/S0129055X25500187). arXiv: [2411.10260](https://arxiv.org/abs/2411.10260) [hep-th] (cit. on pp. 39, 43, 48).
- [GST25] G. Gagliardo, C. Saemann, and R. Tellez-Dominguez. *Principal 3-Bundles with Adjusted Connections*. 2025. arXiv: [2505.13368](https://arxiv.org/abs/2505.13368) [math-ph]. URL: <https://arxiv.org/abs/2505.13368> (cit. on p. 3).
- [GSW09] J. P. Gauntlett, J. Sonner, and T. Wiseman. “Holographic superconductivity in M-Theory”. In: *Phys. Rev. Lett.* 103 (2009), p. 151601. DOI: [10.1103/PhysRevLett.103.151601](https://doi.org/10.1103/PhysRevLett.103.151601) (cit. on p. 5).
- [GSW10] J. Gauntlett, J. Sonner, and T. Wiseman. “Quantum Criticality and Holographic Superconductors in M-theory”. In: *J. High Energ. Phys.* 2010.2 (2010), p. 060. DOI: [10.1007/JHEP02\(2010\)060](https://doi.org/10.1007/JHEP02(2010)060). arXiv: [0912.0512](https://arxiv.org/abs/0912.0512) [hep-th] (cit. on p. 5).
- [Han74] V. L. Hansen. “On the space of maps of a closed surface into the 2-sphere”. In: *Mathematica Scandinavica* 35.2 (1974), pp. 149–158. URL: <https://www.jstor.org/stable/24490694> (cit. on p. 43).
- [Her+07] C. P. Herzog et al. “Quantum critical transport, duality, and M-theory”. In: *Phys. Rev. D* 75 (2007), p. 085020. DOI: [10.1103/PhysRevD.75.085020](https://doi.org/10.1103/PhysRevD.75.085020). arXiv: [hep-th/0701036](https://arxiv.org/abs/hep-th/0701036) [hep-th] (cit. on p. 5).
- [Hes07] K. Hess. “Rational homotopy theory: a brief introduction”. In: *Interactions between Homotopy Theory and Algebra*. Ed. by L. L. Avramov et al. Vol. 436. Contemporary Mathematics. Providence, RI: American Mathematical Society, 2007, pp. 175–202. DOI: [10.1090/conm/436/08409](https://doi.org/10.1090/conm/436/08409). arXiv: [math/0604626](https://arxiv.org/abs/math/0604626) [math.AT] (cit. on p. 24).
- [HS05] M. J. Hopkins and I. M. Singer. “Quadratic Functions in Geometry, Topology, and M-Theory”. In: *Journal of Differential Geometry* 70.3 (July 2005), pp. 329–452. DOI: [10.4310/jdg/1143642908](https://doi.org/10.4310/jdg/1143642908). arXiv: [math/0211216](https://arxiv.org/abs/math/0211216) [math.AT] (cit. on pp. 3, 31).
- [HS97] P. S. Howe and E. Sezgin. “ $D = 11, p = 5$ ”. In: *Phys. Lett. B* 394 (1997), pp. 62–66. DOI: [10.1016/S0370-2693\(96\)01672-3](https://doi.org/10.1016/S0370-2693(96)01672-3). arXiv: [hep-th/9611008](https://arxiv.org/abs/hep-th/9611008) [hep-th] (cit. on p. 41).
- [HSW97] P. S. Howe, E. Sezgin, and P. C. West. “The six-dimensional self-dual tensor”. In: *Physics Letters B* 400.3–4 (1997), pp. 255–259. DOI: [10.1016/S0370-2693\(97\)00365-1](https://doi.org/10.1016/S0370-2693(97)00365-1). arXiv: [hep-th/9702111](https://arxiv.org/abs/hep-th/9702111) [hep-th] (cit. on p. 41).
- [HT92] M. Henneaux and C. Teitelboim. *Quantization of Gauge Systems*. Princeton, NJ: Princeton University Press, 1992. ISBN: 978-0691037691. DOI: [10.2307/j.ctv10crg0r](https://doi.org/10.2307/j.ctv10crg0r) (cit. on pp. 3, 33).
- [Hu59] S.-T. Hu. *Homotopy Theory*. Vol. 8. Pure and Applied Mathematics. New York and London: Academic Press, 1959. URL: <https://webhomes.maths.ed.ac.uk/~v1ranick/papers/hu2.pdf> (cit. on p. 39).
- [Hus+08] D. Husemoller et al. *Basic Bundle Theory and K-Cohomology Invariants*. Vol. 726. Lecture Notes in Physics. Springer, 2008. DOI: [10.1007/978-3-540-74956-1](https://doi.org/10.1007/978-3-540-74956-1) (cit. on p. 46).
- [Igl13] P. Iglesias-Zemmour. *Diffeology*. Vol. 185. Mathematical Surveys and Monographs. Providence, RI: American Mathematical Society (AMS), 2013. ISBN: 978-0-8218-9131-5. URL: <https://bookstore.ams.org/surv-185> (cit. on pp. 8, 10).
- [IM25] A. Ibort and A. Mas. “Smooth sets of fields: A pedagogical introduction”. In: *Geometric Mechanics* 2.3 (Aug. 2025), pp. 251–274. DOI: [10.1142/s2972458925400052](https://doi.org/10.1142/s2972458925400052). arXiv: [2510.20422](https://arxiv.org/abs/2510.20422) [math-ph] (cit. on pp. 4, 7).
- [Jam84] I. M. James. *General Topology and Homotopy Theory*. Springer, 1984. DOI: [10.1007/978-1-4613-8283-6](https://doi.org/10.1007/978-1-4613-8283-6) (cit. on p. 25).

- [Jar87] J. F. Jardine. “Simplicial presheaves”. In: *Journal of Pure and Applied Algebra* 47.1 (1987), pp. 35–87. ISSN: 0022-4049. DOI: [10.1016/0022-4049\(87\)90100-9](https://doi.org/10.1016/0022-4049(87)90100-9) (cit. on p. 5).
- [Joy08] A. Joyal. *The theory of quasi-categories and its applications*. Quaderns 45. Lectures at Advanced Course on Simplicial Methods in Higher Categories, Vol. II. Barcelona: Centre de Recerca Matemàtica, Feb. 2008. URL: <http://mat.uab.cat/~kock/crm/hocat/advanced-course/Quadern45-2.pdf> (cit. on p. 13).
- [Jur+19] B. Jurčo et al. “Higher Structures in M-Theory”. In: *Fortschritte der Physik* 67.8–9 (2019), p. 1910001. DOI: [10.1002/prop.201910001](https://doi.org/10.1002/prop.201910001). arXiv: [1903.02807 \[hep-th\]](https://arxiv.org/abs/1903.02807) (cit. on pp. 4, 5).
- [Kal25] S. Kallel. “Configuration Spaces of Points: A User’s Guide”. In: *Encyclopedia of Mathematical Physics*. Ed. by R. Szabo and M. Bojowald. 2nd ed. Academic Press, 2025, pp. 98–135. ISBN: 9780323957069. DOI: [10.1016/b978-0-323-95703-8.00211-1](https://doi.org/10.1016/b978-0-323-95703-8.00211-1). arXiv: [2407.11092 \[math-ph\]](https://arxiv.org/abs/2407.11092) (cit. on p. 43).
- [Kir90] A. Kirillov. “Geometric Quantization”. In: *Dynamical Systems IV*. Vol. 4. Encyclopedia of Mathematical Sciences. Springer, 1990, pp. 137–172. DOI: [10.1007/978-3-662-06793-2_2](https://doi.org/10.1007/978-3-662-06793-2_2) (cit. on p. 34).
- [Koc96] S. O. Kochman. *Bordism, Stable Homotopy and Adams Spectral Sequences*. Vol. 7. Fields Institute Monographs. Providence, RI: American Mathematical Society, 1996. ISBN: 978-1-4704-3134-1. DOI: [10.1090/fim/007](https://doi.org/10.1090/fim/007). URL: <https://bookstore.ams.org/fim-7> (cit. on pp. 27, 46).
- [KS26] I. Khavkine and U. Schreiber. “Synthetic Geometry of Differential Equations: Jets and Comonad Structure”. In: *Journal of Geometry and Physics* (2026). to appear. arXiv: [1701.06238 \[math.DG\]](https://arxiv.org/abs/1701.06238) (cit. on p. 4).
- [KSS26] S. Kallel, H. Sati, and U. Schreiber. *Higher-Dimensional Anyons via Higher Cohomotopy*. Jan. 2026. arXiv: [2601.03150 \[cond-mat.str-el\]](https://arxiv.org/abs/2601.03150) (cit. on p. 43).
- [KV14] A. Kahle and A. Valentino. “T-Duality and Differential K-Theory”. In: *Communications in Contemporary Mathematics* 16.02 (2014), p. 1350014. DOI: [10.1142/S0219199713500144](https://doi.org/10.1142/S0219199713500144). arXiv: [0912.2516 \[math.KT\]](https://arxiv.org/abs/0912.2516) (cit. on p. 33).
- [Law07] F. W. Lawvere. “Axiomatic cohesion”. In: *Theory and Applications of Categories* 19.3 (2007), pp. 41–49. URL: <http://www.tac.mta.ca/tac/volumes/19/3/19-03abs.html> (cit. on pp. 8, 11).
- [Laz13] A. Lazarev. “Maurer–Cartan moduli and models for function spaces”. In: *Advances in Mathematics* 235 (2013), pp. 296–320. DOI: [10.1016/j.aim.2012.11.009](https://doi.org/10.1016/j.aim.2012.11.009). arXiv: [1109.3715 \[math.AT\]](https://arxiv.org/abs/1109.3715) (cit. on p. 25).
- [Lee12] J. M. Lee. *Introduction to Smooth Manifolds*. Second. Vol. 218. Graduate Texts in Mathematics. New York: Springer, 2012. ISBN: 9781441999825. DOI: [10.1007/978-1-4419-9982-5](https://doi.org/10.1007/978-1-4419-9982-5) (cit. on p. 6).
- [LS22a] C. Lazaroiu and C. S. Shahbazi. “The duality covariant geometry and DSZ quantization of abelian gauge theory”. In: *Advances in Theoretical and Mathematical Physics* 26.7 (2022), pp. 2213–2312. ISSN: 1095-0753. DOI: [10.4310/atmp.2022.v26.n7.a5](https://doi.org/10.4310/atmp.2022.v26.n7.a5). arXiv: [2101.07236 \[math.DG\]](https://arxiv.org/abs/2101.07236) (cit. on pp. 3, 32).
- [LS22b] C. Lazaroiu and C. S. Shahbazi. “The geometry and DSZ quantization of four-dimensional supergravity”. In: *Letters in Mathematical Physics* 113.1 (Dec. 2022), p. 9. ISSN: 1573-0530. DOI: [10.1007/s11005-022-01626-y](https://doi.org/10.1007/s11005-022-01626-y). arXiv: [2101.07778 \[math.DG\]](https://arxiv.org/abs/2101.07778) (cit. on pp. 3, 31, 32).
- [LS93] T. Lada and J. Stasheff. “Introduction to sh Lie algebras for physicists”. In: *Int. J. Theor. Phys.* 32 (1993), pp. 1087–1103. DOI: [10.1007/BF00671791](https://doi.org/10.1007/BF00671791). arXiv: [hep-th/9209099 \[hep-th\]](https://arxiv.org/abs/hep-th/9209099) (cit. on p. 43).
- [LT80] L. L. Larmore and E. Thomas. “On the fundamental group of a space of sections”. In: *Mathematica Scandinavica* 47 (1980), pp. 232–246. URL: <http://www.jstor.org/stable/24491393> (cit. on p. 43).

- [Lur09] J. Lurie. *Higher Topos Theory*. Vol. 170. Annals of Mathematics Studies. Princeton University Press, 2009. ISBN: 978-0691140490. URL: <https://press.princeton.edu/titles/8957.html> (cit. on pp. 5, 17, 45).
- [Lur14] J. Lurie. “Nonabelian Poincaré Duality”. In: (2014). URL: <http://www.math.harvard.edu/~lurie/282ynotes/LectureVIII-Poincare.pdf> (cit. on pp. 27, 47).
- [Lur17] J. Lurie. “Higher Algebra”. Sept. 2017. URL: <https://www.math.ias.edu/~lurie/papers/HA.pdf> (cit. on p. 45).
- [Max65] J. C. Maxwell. “A Dynamical Theory of the Electromagnetic Field”. In: *Philosophical Transactions of the Royal Society of London* 155 (1865), pp. 459–512. DOI: 10.1098/rs.tl.1865.0008. URL: <https://www.jstor.org/stable/108892> (cit. on p. 23).
- [McK19] B. McKay. “Introduction to exterior differential systems”. In: *Banach Center Publications* 117 (2019), pp. 45–55. DOI: 10.4064/bc117-2. arXiv: 1706.09697 [math.DG] (cit. on p. 21).
- [MM03] I. Moerdijk and J. Mrčun. *Introduction to Foliations and Lie Groupoids*. Cambridge: Cambridge University Press, 2003. ISBN: 9780511615450. DOI: 10.1017/cbo9780511615450 (cit. on p. 4).
- [MM97] R. Minasian and G. Moore. “K-theory and Ramond-Ramond charge”. In: *Journal of High Energy Physics* 1997.11 (1997), p. 002. DOI: 10.1088/1126-6708/1997/11/002. arXiv: hep-th/9710230 [hep-th] (cit. on p. 33).
- [MW00] G. Moore and E. Witten. “Self-Duality, Ramond-Ramond Fields, and K-Theory”. In: *JHEP* 05 (2000), p. 032. DOI: 10.1088/1126-6708/2000/05/032. arXiv: hep-th/9912279 [hep-th] (cit. on p. 33).
- [nLa26a] nLab. “extension lemma for differential forms”. In: (2026). URL: <https://ncatlab.org/nlab/show/extension+lemma+for+differential+forms> (cit. on p. 25).
- [nLa26b] nLab. “Supergravity”. In: (2026). URL: <https://ncatlab.org/nlab/show/supergravity> (cit. on p. 3).
- [NSS15] T. Nikolaus, U. Schreiber, and D. Stevenson. “Principal ∞ -bundles: general theory”. In: *Journal of Homotopy and Related Structures* 10.4 (June 2015), pp. 749–801. DOI: 10.1007/s40062-014-0083-6. arXiv: 1207.0248 [math.AT] (cit. on p. 45).
- [Ric20] B. Richter. *From Categories to Homotopy Theory*. Vol. 188. Cambridge Studies in Advanced Mathematics. Cambridge: Cambridge University Press, 2020. DOI: 10.1017/9781108855891 (cit. on p. 5).
- [RS17] G. Rudolph and M. Schmidt. *Differential Geometry and Mathematical Physics: Part II. Fibre Bundles, Topology and Gauge Fields*. Theoretical and Mathematical Physics. Dordrecht: Springer Netherlands, 2017. ISBN: 9789402409598. DOI: 10.1007/978-94-024-0959-8 (cit. on p. 45).
- [RSW26] D. Rist, C. Saemann, and M. Wolf. “Explicit Non-Abelian Gerbes with Connections”. In: *Journal of Physics A: Mathematical and Theoretical* 59.3 (2026), p. 035201. DOI: 10.1088/1751-8121/ae2e60. arXiv: 2203.00092 [math.AT] (cit. on pp. 5, 23).
- [Sat10] H. Sati. “Geometric and Topological Structures Related to M-Branes”. In: *Superstrings, Geometry, Topology, and C*-Algebras*. Vol. 81. Proceedings of Symposia in Pure Mathematics. Providence, RI: American Mathematical Society, 2010, pp. 181–236. ISBN: 978-0-8218-4887-6. DOI: 10.1090/pspum/081/2681765. arXiv: 1001.5020 [math.DG] (cit. on pp. 3, 5).
- [Sat11] H. Sati. “Geometric and topological structures related to M-branes II: Twisted String and String^c structures”. In: *Journal of the Australian Mathematical Society* 90 (2011), pp. 93–108. DOI: 10.1017/S1446788711001261. arXiv: 1007.5419 [hep-th] (cit. on p. 38).
- [Sat18] H. Sati. “Framed M-branes, corners, and topological invariants”. In: *Journal of Mathematical Physics* 59.6 (2018), p. 062304. DOI: 10.1063/1.5007185. arXiv: 1310.1060 [hep-th] (cit. on p. 39).

- [Sch05] U. Schreiber. “From Loop Space Mechanics to Nonabelian Strings”. PhD thesis. Universität Duisburg-Essen, 2005. arXiv: [hep-th/0509163](https://arxiv.org/abs/hep-th/0509163) [[hep-th](#)] (cit. on pp. 5, 23).
- [Sch13] U. Schreiber. “Differential cohomology in a cohesive ∞ -topos”. In: (2013). v2 (2017): <https://ncatlab.org/schreiber/files/dcct170811.pdf>. arXiv: [1310.7930](https://arxiv.org/abs/1310.7930) [[math-ph](#)] (cit. on pp. 4, 5, 8, 11, 15, 17, 18).
- [Sch14] U. Schreiber. *Higher field bundles for Gauge fields*. Talk given at *Operator and geometric analysis on quantum theory*, Levico Terme (Trento), Italy. September 15–19, 2014. Sept. 2014. URL: <https://ncatlab.org/schreiber/show/Higher+Field+Bundles> (cit. on p. 3).
- [Sch16] U. Schreiber. *Higher Structures in Mathematics and Physics*. Talk at Oberwolfach Workshop 1651a: Higher Structures in Geometry and Physics. Mathematisches Forschungsinstitut Oberwolfach, Dec. 2016 (cit. on p. 4).
- [Sch18] U. Schreiber. “Categories and Toposes – Differential Cohesive Higher Toposes”. In: (June 2018). URL: <https://ncatlab.org/schreiber/show/Categories+and+Toposes> (cit. on p. 5).
- [Sch25] U. Schreiber. “Higher Topos Theory in Physics”. In: *Encyclopedia of Mathematical Physics*. Ed. by R. Szabo and M. Bojowald. 2nd ed. Academic Press, 2025, pp. 62–76. ISBN: 9780323957069. DOI: [10.1016/B978-0-323-95703-8.00210-X](https://doi.org/10.1016/B978-0-323-95703-8.00210-X). arXiv: [2311.11026](https://arxiv.org/abs/2311.11026) [[math-ph](#)] (cit. on p. 4).
- [Sch66] J. Schwinger. “Magnetic Charge and Quantum Field Theory”. In: *Phys. Rev.* 144 (1966), pp. 1087–1093. DOI: [10.1103/PhysRev.144.1087](https://doi.org/10.1103/PhysRev.144.1087) (cit. on p. 32).
- [SS21] H. Sati and U. Schreiber. “Twisted cohomotopy implies M5-brane anomaly cancellation”. In: *Letters in Mathematical Physics* 111.5 (Sept. 2021). DOI: [10.1007/s11005-021-01452-8](https://doi.org/10.1007/s11005-021-01452-8). arXiv: [2002.07737](https://arxiv.org/abs/2002.07737) [[hep-th](#)] (cit. on p. 39).
- [SS24a] H. Sati and U. Schreiber. “Cyclification of Orbifolds”. In: *Communications in Mathematical Physics* 405 (2024), p. 67. DOI: [10.1007/s00220-023-04929-w](https://doi.org/10.1007/s00220-023-04929-w). arXiv: [2212.13836](https://arxiv.org/abs/2212.13836) [[math.AT](#)] (cit. on pp. 39, 48).
- [SS24b] H. Sati and U. Schreiber. “Flux Quantization on Phase Space”. In: *Annales Henri Poincaré* 26 (May 2024), pp. 895–919. DOI: [10.1007/s00023-024-01438-x](https://doi.org/10.1007/s00023-024-01438-x). arXiv: [2312.12517](https://arxiv.org/abs/2312.12517) [[hep-th](#)] (cit. on pp. 3, 19, 21, 23, 32).
- [SS25a] H. Sati and U. Schreiber. “Non-Lagrangian Construction of Anyons via Flux Quantization in Cohomotopy”. In: *Journal of Physics: Conference Series* 3152.1 (July 2025). The XXIX International Conference on Integrable Systems and Quantum Symmetries, p. 012024. DOI: [10.1088/1742-6596/3152/1/012024](https://doi.org/10.1088/1742-6596/3152/1/012024). arXiv: [2509.02577](https://arxiv.org/abs/2509.02577) [[math-ph](#)] (cit. on pp. 42, 43).
- [SS25b] H. Sati and U. Schreiber. “Anyons on M5-probes of Seifert 3-orbifolds via Flux Quantization”. In: *Letters in Mathematical Physics* 115.36 (Mar. 2025). DOI: [10.1007/s11005-025-01918-z](https://doi.org/10.1007/s11005-025-01918-z). arXiv: [2411.16852](https://arxiv.org/abs/2411.16852) [[hep-th](#)] (cit. on p. 42).
- [SS25c] H. Sati and U. Schreiber. “Engineering of Anyons on M5-Probes via Flux Quantization”. In: *SciPost Physics Lecture Notes* 107 (2025). DOI: [10.21468/SciPostPhysLectNotes.107](https://doi.org/10.21468/SciPostPhysLectNotes.107). arXiv: [2501.17927](https://arxiv.org/abs/2501.17927) [[hep-th](#)] (cit. on p. 42).
- [SS25d] H. Sati and U. Schreiber. “Flux Quantization”. In: *Encyclopedia of Mathematical Physics*. Ed. by R. Szabo and M. Bojowald. 2nd ed. Vol. 4. Academic Press, 2025, pp. 281–324. ISBN: 9780323957069. DOI: [10.1016/b978-0-323-95703-8.00078-1](https://doi.org/10.1016/b978-0-323-95703-8.00078-1). arXiv: [2402.18473](https://arxiv.org/abs/2402.18473) [[hep-th](#)] (cit. on pp. 3, 25, 28, 31, 33).
- [SS25e] H. Sati and U. Schreiber. “Fractional Quantum Hall Anyons via the Algebraic Topology of Exotic Flux Quanta”. In: (2025). arXiv: [2505.22144](https://arxiv.org/abs/2505.22144) [[cond-mat.mes-hall](#)] (cit. on pp. 42, 43).
- [SS25f] H. Sati and U. Schreiber. “Fragile Topological Phases and Topological Order of 2D Crystalline Chern Insulators”. In: (Dec. 2025). arXiv: [2512.24709](https://arxiv.org/abs/2512.24709) [[cond-mat.str-el](#)] (cit. on p. 43).

- [SS25g] H. Sati and U. Schreiber. “Quantum Observables of Quantized Fluxes”. In: *Annales Henri Poincaré* 26 (2025), pp. 4241–4269. DOI: [10.1007/s00023-024-01517-z](https://doi.org/10.1007/s00023-024-01517-z). arXiv: [2306.01214](https://arxiv.org/abs/2306.01214) [hep-th] (cit. on p. 35).
- [SS25h] H. Sati and U. Schreiber. “Renormalization of Chern-Simons Wilson Loops via Flux Quantization in Cohomotopy”. In: *Reviews in Mathematical Physics* (2025). DOI: <https://doi.org/10.1142/S0129055X25500382>. arXiv: [2509.25336](https://arxiv.org/abs/2509.25336) [hep-th] (cit. on p. 43).
- [SS25i] H. Sati and U. Schreiber. “The Character Map in Twisted Equivariant Nonabelian Cohomology”. In: *Beijing Journal of Pure and Applied Mathematics* 2.2 (2025), pp. 515–617. DOI: [10.4310/bpam.250908174706](https://doi.org/10.4310/bpam.250908174706). arXiv: [2011.06533](https://arxiv.org/abs/2011.06533) [hep-th] (cit. on pp. 27, 36, 47).
- [SS26a] H. Sati and U. Schreiber. *Bulk-Edge Correspondence via Higher Gauge Theory*. 2026. arXiv: [2605.10232](https://arxiv.org/abs/2605.10232) [hep-th] (cit. on pp. 39, 42).
- [SS26b] H. Sati and U. Schreiber. “Cohomotopy, Framed Links, and Abelian Anyons”. In: *Proceedings of the Focus Program on Algebraic Topology in Memory of Fred Cohen*. Fields Institute Communications. In press. Springer, 2026. arXiv: [2408.11896](https://arxiv.org/abs/2408.11896) [hep-th] (cit. on pp. 42, 43).
- [SS26c] H. Sati and U. Schreiber. “Complete Topological Quantization of Higher Gauge Fields”. In: *SciPost Physics Lecture Notes* 127 (2026). DOI: [10.21468/SciPostPhysLectNotes.127](https://doi.org/10.21468/SciPostPhysLectNotes.127). arXiv: [2512.12431](https://arxiv.org/abs/2512.12431) [hep-th] (cit. on p. 35).
- [SS26d] H. Sati and U. Schreiber. *Equivariant Principal ∞ -Bundles*. Cambridge University Press, 2026. ISBN: 9781009698559. arXiv: [2112.13654](https://arxiv.org/abs/2112.13654) [math.AT]. URL: <https://ncatlab.org/schreiber/show/EPBund> (cit. on pp. 4, 5, 16, 17, 46, 47).
- [SS26e] H. Sati and U. Schreiber. *Geometric Orbifold Cohomology*. CRC Press, 2026. ISBN: 9781041147510. URL: <https://ncatlab.org/schreiber/show/OrbCoh> (cit. on pp. 4, 5, 27, 36, 45, 47).
- [SS26f] H. Sati and U. Schreiber. “Identifying Anyonic Topological Order in Fractional Quantum Anomalous Hall Systems”. In: *Applied Physics Letters* 128 (2026). Special issue: Quantum Geometry in Condensed Matter: Fundamentals and Applications, p. 023101. DOI: [10.1063/5.0305441](https://doi.org/10.1063/5.0305441). arXiv: [2507.00138](https://arxiv.org/abs/2507.00138) [cond-mat.str-el] (cit. on p. 43).
- [SSS09] H. Sati, U. Schreiber, and J. Stasheff. “ L_∞ -Algebra Connections and Applications to String- and Chern-Simons n -Transport”. In: *Quantum Field Theory*. Ed. by G. Battle et al. Vol. 272. Progress in Mathematics. Birkhäuser Basel, 2009, pp. 303–424. ISBN: 978-3-7643-8736-5. DOI: [10.1007/978-3-7643-8736-5_17](https://doi.org/10.1007/978-3-7643-8736-5_17). arXiv: [0801.3480](https://arxiv.org/abs/0801.3480) [math.DG] (cit. on p. 43).
- [SSS12] H. Sati, U. Schreiber, and J. Stasheff. “Twisted Differential String and Fivebrane Structures”. In: *Commun. Math. Phys.* 315.1 (2012), pp. 169–213. DOI: [10.1007/s00220-012-1510-3](https://doi.org/10.1007/s00220-012-1510-3). arXiv: [0910.4001](https://arxiv.org/abs/0910.4001) [math.AT] (cit. on pp. 5, 23).
- [Ste67] N. E. Steenrod. “A convenient category of topological spaces”. In: *Michigan Mathematical Journal* 14.2 (1967), pp. 133–152. DOI: [10.1307/mmj/1028999711](https://doi.org/10.1307/mmj/1028999711) (cit. on p. 8).
- [SV23] H. Sati and A. A. Voronov. “Mysterious Triality and Rational Homotopy Theory”. In: *Communications in Mathematical Physics* 400.3 (June 2023), pp. 1915–1960. DOI: [10.1007/s00220-023-04643-7](https://doi.org/10.1007/s00220-023-04643-7). arXiv: [2111.14810](https://arxiv.org/abs/2111.14810) [hep-th] (cit. on p. 39).
- [SW13] U. Schreiber and K. Waldorf. “Connections on non-abelian gerbes and their holonomy”. In: *Theory and Applications of Categories* 28.17 (2013), pp. 476–540. arXiv: [0808.1923](https://arxiv.org/abs/0808.1923) [math.DG]. URL: <http://www.tac.mta.ca/tac/volumes/28/17/28-17abs.html> (cit. on pp. 5, 23).
- [SW76] D. J. Simms and N. M. J. Woodhouse. “Observables”. In: *Lectures on Geometric Quantization*. Vol. 53. Lecture Notes in Physics. Springer, 1976. Chap. 4. ISBN: 978-3-540-07860-9. DOI: [10.1007/3-540-07860-6_4](https://doi.org/10.1007/3-540-07860-6_4) (cit. on p. 34).
- [Sza13] R. J. Szabo. “Quantization of Higher Abelian Gauge Theory in Generalized Differential Cohomology”. In: *Proceedings of 7th International Conference on Mathematical Methods*

- in Physics — PoS(ICMP 2012)* (2013), p. 009. DOI: [10.22323/1.175.0009](https://doi.org/10.22323/1.175.0009) (cit. on p. 31).
- [Tim23] C. Timm. *Theory of Superconductivity*. Lecture notes, TU Dresden. 2023. URL: <https://ncatlab.org/nlab/files/Timm-Superconductivity.pdf> (cit. on p. 33).
- [Toë02] B. Toën. “Stacks and Non-abelian cohomology”. In: (2002). URL: <https://perso.math.univ-toulouse.fr/btoen/files/2015/02/msri2002.pdf> (cit. on pp. 27, 47).
- [Ton16] D. Tong. *Lectures on the Quantum Hall Effect*. 2016. URL: <http://www.damtp.cam.ac.uk/user/tong/qhe.html> (cit. on p. 43).
- [Tu11] L. W. Tu. *An Introduction to Manifolds*. 2nd. Universitext. New York, NY: Springer, 2011. ISBN: 978-1-4419-7399-3. DOI: [10.1007/978-1-4419-7400-6](https://doi.org/10.1007/978-1-4419-7400-6) (cit. on p. 6).
- [TV05] B. Toën and G. Vezzosi. “Homotopical Algebraic Geometry I: Topos Theory”. In: *Advances in Mathematics* 193.2 (2005), pp. 257–372. DOI: [10.1016/j.aim.2004.05.004](https://doi.org/10.1016/j.aim.2004.05.004). arXiv: [math/0207028](https://arxiv.org/abs/math/0207028) [math.AG] (cit. on p. 5).
- [Wal26] K. Waldorf. “Adjusted connections on non-abelian bundle gerbes”. In: (2026). arXiv: [2604.24513](https://arxiv.org/abs/2604.24513) [math.DG] (cit. on pp. 5, 23).
- [Wit97a] E. Witten. “Five-Brane Effective Action in M-Theory”. In: *Journal of Geometry and Physics* 22.2 (May 1997), pp. 103–133. DOI: [10.1016/S0393-0440\(97\)80160-X](https://doi.org/10.1016/S0393-0440(97)80160-X). arXiv: [hep-th/9610234](https://arxiv.org/abs/hep-th/9610234) [hep-th] (cit. on pp. 3, 39).
- [Wit97b] E. Witten. “On Flux Quantization in M-Theory and the Effective Action”. In: *Journal of Geometry and Physics* 22.1 (Apr. 1997), pp. 1–13. DOI: [10.1016/S0393-0440\(96\)00042-3](https://doi.org/10.1016/S0393-0440(96)00042-3). arXiv: [hep-th/9609122](https://arxiv.org/abs/hep-th/9609122) [hep-th] (cit. on pp. 3, 39).
- [WP94] K. K. Wan and J. J. Powis. “Quantum mechanics in Dirac’s front form”. In: *International Journal of Theoretical Physics* 33 (1994), pp. 553–574. DOI: [10.1007/BF00670516](https://doi.org/10.1007/BF00670516) (cit. on p. 35).
- [Zwa68] D. Zwanziger. “Quantum Field Theory of Particles with Both Electric and Magnetic Charges”. In: *Phys. Rev.* 176 (1968), pp. 1489–1495. DOI: [10.1103/PhysRev.176.1489](https://doi.org/10.1103/PhysRev.176.1489) (cit. on p. 32).