

Cyclification of Orbifolds

Hisham Sati¹, Urs Schreiber¹

October 3, 2023

Abstract

Inertia orbifolds homotopy-quotiented by rotation of *geometric* loops play a fundamental role not only in ordinary cyclic cohomology, but more recently in constructions of equivariant Tate-elliptic cohomology and generally of transchromatic characters on generalized cohomology theories. Nevertheless, existing discussion of such *cyclified stacks* has been relying on ad-hoc component presentations with intransparent and unverified stacky homotopy type.

Following our previous formulation of transgression of cohomological charges (“double dimensional reduction”), we explain how cyclification of ∞ -stacks is a fundamental and elementary base-change construction over moduli stacks in cohesive higher topos theory (cohesive homotopy type theory). We prove that Ganter/Huan’s extended inertia groupoid used to define equivariant quasi-elliptic cohomology is indeed a model for this intrinsically defined cyclification of orbifolds, and we show that cyclification implements transgression in group cohomology in general, and hence in particular the transgression of degree-4 twists of equivariant Tate-elliptic cohomology to degree-3 twists of orbifold K-theory on the cyclified orbifold.

As an application, we show that the universal shifted integral 4-class of equivariant 4-Cohomotopy theory on ADE-orbifolds induces the Platonic 4-twist of ADE-equivariant Tate-elliptic cohomology; and we close by explaining how this should relate to elliptic M5-brane genera, under our previously formulated *Hypothesis H*.

Contents

1	Introduction and Overview	2
2	Cyclic inertia orbifolds	10
3	Transgression as cyclification	18
4	Integral 4-classes of equivariant 4-Cohomotopy	21
A	Appendix: Technical background	24

¹Mathematics, Division of Science; and
Center for Quantum and Topological Systems (CQTS),
NYUAD Research Institute,
New York University Abu Dhabi, UAE.

Acknowledgements. We thank Zhen Huan for useful discussion and two anonymous referees for careful reading of our manuscript. The authors acknowledge support by *Tamkeen* under the *NYU Abu Dhabi Research Institute grant* CG008.

1 Introduction and Overview

Classical cyclic loop spaces. Topological spaces of free loops (e.g. [CO15]) in a given topological space \mathcal{X} , but homotopy-quotiented by the rigid rotation action of the topological circle group S^1 on itself,

$$\text{Cyc}(\mathcal{X}) := \underbrace{\text{Map}(S^1, \mathcal{X})}_{\text{free loop space}} // S^1 \quad \begin{array}{l} \text{homotopy quotient} \\ \text{by loop rotations} \end{array}$$

cyclic loop space circle topological space

have a long tradition in the study of the elliptic cohomology (see eg. [Re15]) of \mathcal{X} , at least in the extreme but still surprisingly rich limit of restricting to those loops which are effectively constant, and traditionally perceived through their S^1 -equivariant K-cohomology ([KM04, §5], review in [Do19, §6.2]).

In the generality where $\mathcal{X} = X // G$ is itself a homotopy quotient space, this goes back to [Wit88], where a topological point-set model for $\text{Map}(S^1, X // G)$ is called a “twisted loop space” of X . There this is thought of as a model for the configuration space of closed strings propagating on the orbifold $X // G$ (whose non-trivial orbifold transition functions are traditionally called their “twisted sectors” [DHVW85, p. 3], whence Witten’s terminology, see also e.g. [St15]). Accordingly, some authors call $\text{Cyc}(\mathcal{X})$ the *string space of \mathcal{X}* [Cha05, §4.8.1][BO05, p. 1].

However, this terminology is neither widely adopted nor quite appropriate, since not all free loop spaces arise as configuration spaces of strings – not even in string theory; and even when they do one is going to be interested in their twisted cohomology in addition to, and in a sense different from, the string’s “twisted sectors”. But *Jones’ theorem* ([Jo87, Thm. A], review in [Lo92, Cor. 7.3.14][Lo15, §3,4]) shows that the ordinary cohomology of $\text{Cyc}(X)$ for a simply-connected topological space X is its *cyclic cohomology* — which is meaningful and standard mathematical terminology. Therefore we call $\text{Cyc}(\mathcal{X})$ the *cyclification* of \mathcal{X} (following [FSS16TDI, §3][BSS18, §2.2]), also in line with the modern terminology of *cyclotomic spectra* [BM15][NS18], obtained from approximating the free loop S^1 -space $\text{Map}(S^1_{\text{coh}}, X)$ by its suspension spectrum.

Cyclic inertia orbifolds? In any case, when $G \curvearrowright X$ carries the geometric structure of a G -manifold (as it certainly does already in the motivating examples from string theory), one should expect a more fine-grained incarnation of $\text{Cyc}(X // G)$, lifting it from plain homotopy theory to the geometric homotopy theory (homotopy topos theory [Lu09][Re10]) of *orbifolds* regarded (cf. [Le10][SS20Orb]) as *stacks* (useful background references for our purposes are [Ho08][Ja15]), specifically topological stacks or differentiable stacks (cf., e.g., [Ca11]). Concretely, one should expect to make sense of the cyclification of $X // G$ regarded as a smooth orbifold, so that the orbifold K-theory of the *cyclic loop stack* $\text{Cyc}(X // G)$ would reflect the properly G -equivariant elliptic cohomology of X .

Finally, one should expect that such a cyclified orbifold may canonically be restricted to its “essentially constant loops”, which in themselves ought to constitute the familiar “inertia stack” (recalled in §2.1) $\Lambda(X // G) = \text{Map}(\mathbf{B}\mathbb{Z}, X // G)$ of the orbifold. In conclusion then, one should expect that the “essentially constant”-cyclification of an orbifold should be a homotopy quotient by the *topological* (“geometric”, “stacky”, “cohesive”) circle group S^1_{coh} (Ntn. A.48) of the inertia orbifold embedded inside the cyclified orbifold:


$$\underbrace{\text{Map}(\mathbf{B}\mathbb{Z}, X // G)}_{\text{inertia orbifold}} // S^1_{\text{coh}} \quad \xrightarrow{\substack{\text{include as essentially} \\ \text{constant loops}}} \quad \underbrace{\text{Map}(S^1_{\text{coh}}, X // G)}_{\text{smooth loop stack}} // S^1_{\text{coh}} \quad =: \text{Cyc}(X // G). \quad (1)$$

stacky S^1_{coh} -quotient stacky S^1_{coh} -quotient

For a long time it had been unclear how to bear out these expectations, nor had they been approached with tools from geometric homotopy theory. After crucial proposals in [Ga07, Def. 2.3 & 3.1][Ga13, Def. 2.6], and following advice by C. Rezk, a candidate component model for the expected stack was finally given in [Hua18a, Def. 2.5, 2.9] – see (27) below – and justified by demonstrating that it does support a satisfactory notion of “quasi-elliptic” cohomology (further discussed in [Hua18b][HS20][HY22]).

What has been left open is a proof that the component presentation (27) from [Hua18a] does present the abstract stacky construction (1), hence does satisfy expected abstract properties. This is what we prove here – in Thm. 2.3.

Embedding into cohesive homotopy theory. We approach this issue by embedding the situation into the “extremely convenient” (in the technical sense going back to [St67]) higher topos of smooth ∞ -groupoids (∞ -stacks over a site of smooth manifolds), laid out¹ in [SS20Orb][SS21EPB] (a reference list for our notations is given below in Ntn. A.48).



As indicated by this homotopy-commutative diagram of ∞ -categories (reproduced from [SS21EPB], where the relevant proofs are given in §3.3.1), each stage in this sequence of ever more “convenient” categories of spaces comes with its own notion of underlying homotopy types. All these notions are compatible and culminate in the operation of sending a smooth ∞ -groupoid \mathcal{X} to its *pure shape* $\int \mathcal{X}$ (cf. [SS21EPB, fn 1]), which generalizes ([BBP19][SS21EPB, pp. 144]) the traditional construction of *singular simplicial complexes* (thought of as higher path ∞ -groupoids) from topological spaces to smooth higher stacks:

$$\begin{array}{c}
 \text{shape unit} \\
 \text{transformation} \\
 \eta_{\mathcal{X}}^{\int}
 \end{array}
 : \mathcal{X} \xrightarrow{\text{includes constant paths into}} \int \mathcal{X} \xrightarrow{\text{pure shape of smooth type}} \lim_{n:\mathbb{N}} \underbrace{\text{Map}(\Delta_{\text{smth}}^n, \mathcal{X})}_{n\text{-dim. paths in } \mathcal{X}}
 \quad (3)$$

\simeq is equivalently amalgamation of smooth images of smooth simplices in the smooth type

One of the simplest non-trivial examples of the shape operation is already the most important one for the purpose of cyclification: Namely, the shape of the smooth circle is equivalent to the groupoid with a single object $*$ (witnessing that the circle is connected) which has \mathbb{Z} -worth of automorphisms (witnessing the winding number of paths starting and ending at this point, see Lem. 2.8):

$$\begin{array}{ccc}
 \mathbb{R}_{\text{smth}}^1 & \longrightarrow & * \\
 \downarrow & \text{(pb)} & \downarrow \\
 \text{the geometric circle } S_{\text{coh}}^1 & \xrightarrow{\eta_{S_{\text{coh}}^1}^{\int}} & \int S_{\text{coh}}^1 \simeq * // \mathbb{Z} \simeq \mathbf{B}\mathbb{Z} \quad \text{and its pure shape}
 \end{array}
 \quad (4)$$

On general abstract grounds (33), this immediately induces the desired structure (1), by the following homotopy pullback construction (which we explain below in §2.2):

$$\begin{array}{ccccc}
 \text{Map}(\mathbf{B}\mathbb{Z}, \mathcal{X}) // \mathbf{B}\mathbb{Z} & \longleftarrow & \text{Map}(\mathbf{B}\mathbb{Z}, \mathcal{X}) // S_{\text{coh}}^1 & \xrightarrow{\text{include as essentially constant loops}} & \text{Map}(S_{\text{coh}}^1, \mathcal{X}) // S_{\text{coh}}^1 \\
 \downarrow & & \downarrow & \text{Map}(\eta_{S_{\text{coh}}^1}^{\int}, \mathcal{X}) // S_{\text{coh}}^1 & \downarrow \\
 \mathbf{B}^2\mathbb{Z} & \longleftarrow & \mathbf{B}S_{\text{coh}}^1 & \xrightarrow{\text{include as essentially constant loops}} & \mathbf{B}S_{\text{coh}}^1 \\
 & & \text{B}\eta_{S_{\text{coh}}^1}^{\int} & &
 \end{array}
 \quad (5)$$

¹The articles [SS20Orb][SS21EPB] go further to the “singular-cohesive” homotopy theory of equivariant smooth stacks, ultimately needed for their properly equivariant cohomology theory. For brevity, here we do not dwell on this further step, but our results immediately make cyclified orbifolds available in this context of proper equivariant homotopy theory.

As the terms colored in purple in (5) are meant to highlight, the problem (1) of forming S_{coh}^1 -cyclic inertia orbifolds is abstractly solved by the shape unit (4) provided by cohesive homotopy theory: this is what knows about “restriction to essentially constant loops” in a way compatible with their S_{coh}^1 -action.

With this good abstract conception of cyclification of orbifolds in hand, and having recovered from it the existing component constructions via Thm. 2.3, we discover a wealth of interesting induced phenomena:

Transgression as cyclification of cocycles. One should expect that any kind of cohomology of orbifolds $X // G$ “transgresses to” (i.e.: functorially induces subject to degree shifts) cohomology of the cyclification – generalizing the familiar notion of transgression in group cohomology, which is the special case where $X = *$. In particular, for the purpose of twisted elliptic cohomology one imagines transgressing integral 4-cocycles on an orbifold to integral 3-cocycles on its cyclification, which there serve as twists for topological K-theory. Again, some component formulas have been considered ([ARZ07, Def. 4.1][Wi08, §1.3.3], recalled as Def. 3.1 below), but a general abstract formulation seems to have been missing.

However, we may observe that any decent generalized cohomology theory on orbifolds will be represented by some (equivariant) moduli stack \mathcal{A} (cf. [FSS20Cha, §II][SS20Orb, p. 6 & §5][SS21EPB, §4.3]), in that its cocycles are maps of (equivariant) smooth ∞ -stacks from the orbifold to this moduli ∞ -stack:

$$\begin{array}{ccc}
 & \text{cocycle in} & \\
 & \text{\textit{G-equivariant } \mathcal{A}\text{-cohomology}} & \\
 & F & \\
 \text{differentiable stack} & \curvearrowright & \text{differential} \\
 \text{/ orbifold} & & \text{moduli} \\
 X // G & & \infty\text{-stack} \\
 & & \mathcal{A} \\
 & \text{homotopy/} & \\
 & \text{coboundary} & \\
 & \Downarrow & \\
 & F' & \\
 & \text{cohomologous cocycle} &
 \end{array} \tag{6}$$

For example [FSS20Cha, Ex. 2.2], in the simple case of ordinary cohomology in some degree $n \in \mathbb{N}$ with coefficients in an abelian group A , the moduli stack is the n -fold delooping stack $\mathbf{B}^n A$ (we assume A to be discrete just for the purpose of exposition):

$$\begin{array}{c}
 \text{ordinary} \\
 \text{equivariant cohomology} \\
 H^n(X // G; A) \simeq \pi_0 \left\{ X // G \xrightarrow{F} \mathbf{B}^n A \right\}.
 \end{array} \tag{7}$$

But with cyclification of orbifolds identified as the abstract stacky construction (1), it immediately extends to an ∞ -functor on all ∞ -stacks. This allows to readily define/construct the cyclic transgression of any generalized orbifold cocycles (6) simply as the image formed under this ∞ -functor:

$$\begin{array}{ccc}
 & \text{cyclified cocycle} & \\
 & \text{in } \text{Cyc}(\mathcal{A})\text{-cohomology} & \\
 & \text{Cyc}(F) & \\
 \text{cyclified} & \curvearrowright & \text{cyclified} \\
 \text{orbifold} & & \text{moduli } \infty\text{-stack} \\
 \text{Cyc}(X // G) & & \text{Cyc}(\mathcal{A}) \\
 & \Downarrow \wr & \\
 & \text{Cyc}(F') &
 \end{array} \tag{8}$$

Moreover, in the example (7) of ordinary cohomology with coefficients in a torsion-free abelian group A (such as the integers \mathbb{Z}), one finds a retraction of the cyclification of the classifying stack in degree $n + 1$ onto that in degree n , which has the interpretation of “integrating cocycles along loops”:

$$\begin{array}{c}
 A : \text{AbGrp}, \\
 n : \mathbb{N}_{\geq 2}
 \end{array}
 \vdash
 \mathbf{B}^n A \xrightarrow{(\text{id}, 0)} \mathbf{B}^n A \times \mathbf{B}^{n+1} A \twoheadrightarrow \overbrace{(\mathbf{B}^n A \times \mathbf{B}^{n+1} A) // S_{\text{coh}}^1}^{\text{Cyc}(\mathbf{B}^{n+1} A)} \xrightarrow{\int_{S_{\text{coh}}^1}} \mathbf{B}^n A. \tag{9}$$

This may nicely be seen, due to the assumption that A is torsion-free, with tools from rational homotopy theory (comprehensive review and further pointers are given in [FSS20Cha, §3.2]): Using the general formula for Sullivan models (see particularly [Me15]) of cyclic loop spaces from [VPB85, Thm. A] (described in our context in [FSS17Sph, Prop. 3.2][FSS16TDI, Rem. 3.1], and iterated and extended to the nilpotent case in [SV21]), we have the following evident corresponding retraction of rational Sullivan dg-algebras (we are showing the polynomial generators, with degrees in subscript, and their differential relations):

$$\begin{array}{l}
\text{Retraction of cyclification of} \\
\text{torsion-free classifying space} \\
\text{onto its based loop space} \\
\\
\text{witnessed by the respective} \\
\text{Sullivan model dgc-algebras}
\end{array}
\quad
\begin{array}{c}
\mathbf{B}^n \mathbb{Z} \longrightarrow \mathbf{B}^n \mathbb{Z} \times \mathbf{B}^{n+1} \mathbb{Z} \longrightarrow \text{Cyc}(\mathbf{B}^{n+1} \mathbb{Z}) \xrightarrow{\int_{S^1_{\text{coh}}}} \mathbf{B}^n \mathbb{Z} . \\
\\
(d c_n = 0) \longleftarrow \left(\begin{array}{l} d c_{n+1} = 0 \\ d c_n = 0 \end{array} \right) \longleftarrow \left(\begin{array}{l} d c_{n+1} = c_n \wedge \omega_2 \\ d c_n = 0 \\ d \omega_2 = 0 \end{array} \right) \longleftarrow (d c_n = 0)
\end{array}
\quad (10)$$

Now under this retraction, the cyclification operation (8) reproduces the traditional transgression formula in discrete group cohomology (recalled as Def. 3.1 below) and shows that it descends from the free loop stack to the cyclification:

$$\begin{array}{c}
\begin{array}{ccc}
G : \text{Grp}(\text{Set}), & * // G & \xrightarrow[\text{cocycle in group cohomology}]{F} & \mathbf{B}^{n+1} A & \vdash \\
\\
\text{inertia orbifold} \\
\Lambda B G = \text{Map}(S^1_{\text{coh}}, * // G) & \xrightarrow[\text{looped cocycle}]{\text{Map}(S^1_{\text{coh}}, F)} & \text{Map}(S^1_{\text{coh}}, \mathbf{B}^n A) & \simeq & \mathbf{B}^n A \times \mathbf{B}^{n+1} A & \xrightarrow{\text{pr}_1} & \mathbf{B}^n A & (11) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & \nearrow \\
\text{cyclified orbifold} \\
\text{Cyc}(* // G) & \xrightarrow[\text{cyclified cocycle}]{\text{Cyc}(c)} & \text{Cyc}(\mathbf{B}^{n+1} A) & \simeq & (\mathbf{B}^n A \times \mathbf{B}^{n+1} A) // S^1_{\text{coh}} & \xrightarrow[\text{loop integration}]{\int_{S^1_{\text{coh}}}} & \mathbf{B}^n A & (9) \\
& & \text{cyclified coefficient } \infty\text{-stack} & & & & &
\end{array}
\end{array}$$

This is Thm. 3.4 below, proving a suggestion in [Wi08, §1.3.3]. But notice that the construction (11) works for $* // G = \mathbf{B}G$ replaced by any ∞ -stack \mathcal{X} and hence defines transgression in this generality. In the special case when \mathcal{X} is at most an orbifold, transgression is considered in the literature as pullback in cohomology along the evaluation map $S^1_{\text{coh}} \times \text{Map}(S^1_{\text{coh}}, \mathcal{X}) \xrightarrow{\text{ev}} \mathcal{X}$ (61) followed by suitable fiber integration over the S^1_{coh} -factor (e.g. [LU06, p. 2], following analogous discussion for manifolds [Br93, §3.5]). The proof of Thm. 3.4 brings out that this is what (11) reduces to in these cases; see around (48) below.

Therefore, cyclification subsumes transgression in broad generality, but it retains more information. We next see that this extra information is such as to recover the original cocycle:

Fiber integration via cyclification. In giving higher topos-theoretic meaning to the cyclification-construction of orbifolds, a web of further structure surrounding the construction becomes manifest, related to the topic of higher transformation groups and higher principal bundles (cf. [SS20Orb, §2.2][SS21EPB, §3.2.3]):

First (this is the content of §2 below), cyclification immediately generalizes from the circle group S^1_{coh} to any group ∞ -stack $\mathcal{T} \in \text{Grp}(\text{SmthGrpd}_{\infty})$, since the mapping stacks of the form $\text{Map}(\mathcal{T}, \mathcal{X})$ carry a canonical \mathcal{T} -action by precomposition with the multiplication action of \mathcal{T} on itself. Moreover, the general theory of principal ∞ -bundles shows that the resulting homotopy quotient projection is a \mathcal{T} -principal bundle which is classified by the homotopy quotient of the terminal map $\text{Map}(\mathcal{T}, \mathcal{X})$, in that we have homotopy pullback squares of this form:

$$\begin{array}{ccccc}
\mathcal{T} & \longrightarrow & \text{Map}(\mathcal{T}, \mathcal{X}) & \longrightarrow & * \\
\downarrow & \text{(pb)} & \downarrow & \text{(pb)} & \downarrow \\
* & \longrightarrow & \underbrace{\text{Map}(\mathcal{T}, \mathcal{X}) // \mathcal{T}}_{\text{Cyc}_{\mathcal{T}}} & \xrightarrow{c_1} & * // \mathcal{T} \\
& & & & \underbrace{\hspace{2cm}}_{\mathbf{B}\mathcal{T}}
\end{array}
\quad (12)$$

In fact, these \mathcal{T} -principal bundles over \mathcal{T} -cyclic stacks have a fundamental universal property (31): their construction is *right adjoint* (in the ∞ -category theoretic sense of, e.g., [RV20, Def. 1.1.2]) to sending a \mathcal{T} -principal bundle to its total space (i.e.: total ∞ -stack):

$$\begin{array}{ccc}
 \text{SmthGrpd}_\infty & \xleftarrow[\perp]{\text{total space}} & \mathcal{T}\text{PrncplBundl}(\text{SmthGrpd}_\infty) & \xrightarrow{\text{base space}} & \text{SmthGrpd}_\infty & (13) \\
 & & \mathcal{X} \mapsto (\text{Map}(\mathcal{T}, \mathcal{X}) \rightarrow \text{Map}(\mathcal{T}, \mathcal{X}) // \mathcal{T}) & & & \\
 & & & & \mathcal{X} \mapsto \text{Cyc}_{\mathcal{T}}(\mathcal{X}) & \\
 & & & & \parallel & \\
 & & & & \mathcal{T}\text{-cyclification} &
 \end{array}$$

This adjunction means that if a domain \mathcal{X} is a \mathcal{T} -principal bundle over a base $\mathcal{Y} \simeq \mathcal{X} // \mathcal{T}$ (as such classified by a cocycle $\mathcal{Y} \rightarrow \mathbf{B}\mathcal{T}$), then its \mathcal{A} -cohomology (6) is equivalently the cohomology of \mathcal{Y} with coefficients in $\text{Cyc}(\mathcal{A})$ subject to identification of the underlying \mathcal{T} -bundles on both sides:

$$\begin{array}{ccccc}
 \mathcal{T} & \xrightarrow{\mathcal{T}\text{-principal bundle}} & \mathcal{X} & \xrightarrow[\text{cocycle on total space}]{F} & \mathcal{A} & \xleftarrow{\text{oxidation}} & \text{generalized cohomology coefficients} \\
 & & \downarrow & & \updownarrow & & \\
 & & \mathcal{Y} & \xrightarrow[\text{Cyc-adjoint cocycle on base}]{\tilde{F}} & \text{Cyc}_{\mathcal{T}}(\mathcal{A}) & \xleftarrow{\text{reduction}} & \\
 & & \downarrow \wr & & \downarrow c_1 & & \\
 & & \mathbf{B}\mathcal{T} & & & &
 \end{array}$$

In the case (9) of ordinary cohomology of S^1_{coh} -bundles, this yields the fiber integration of cocycles:

$$\begin{array}{ccccc}
 S^1_{\text{coh}} & \xrightarrow[S^1_{\text{coh}}\text{-principal bundle}]{\mathcal{X}} & \xrightarrow[\text{(n+1)-cocycle}]{F} & \mathbf{B}^{n+1}A & \xrightarrow{\text{ordinary cohomology coefficients}} & \\
 & & \downarrow & & \updownarrow & \\
 & & \mathcal{Y} & \xrightarrow[\text{n-cocycle}]{\tilde{F}} & \text{Cyc}(\mathbf{B}^{n+1}A) & \xrightarrow{\int_{S^1_{\text{coh}}}} & \mathbf{B}^n A & \xleftarrow{\text{fiber integration / double dim-reduction}} & \\
 & & & & & & & &
 \end{array}$$

Double dimensional reduction. If in (15) we think of

- \mathcal{X} as a spacetime-orbifold (e.g. as in [DHVW85][Ac99][SS20Tad]),
- $\mathbf{B}^{n+1}A$ as the coefficients of charges of some solitonic physical objects (“branes”) in higher generalization (cf. [FSS20Cha, (2)]) of Dirac’s classical magnetic charge quantization (reviewed e.g. in [Al85, §2]),
- $\mathcal{T} = S^1_{\text{coh}}$ as the “Kaluza-Klein compactification”-space (see [Du94] for traditional pointers and [Al20] for discussion in our higher differential geometric context)

then (14) captures the “double dimensional reduction” [DHIS87] of these brane charges ([FSS16TDI, §][Sc16, §4] [BSS18]), namely their descent to charges of lower-dimensional branes in the lower-dimensional base spacetime:

$$\begin{array}{ccc}
 S^1_{\text{coh}} & \xrightarrow[\text{1+(d+1)-dimensional } S^1\text{-fibred spacetime orbifold}]{\mathcal{X}} & \xrightarrow[\text{magnetic } p\text{-brane charge coupling to electric } d-p-2\text{-branes}]{F} & \mathbf{B}^{d-p}\mathbb{Z} & & \text{e.g.} & \begin{array}{l} 1+(d+1) = 11 \text{ M-theory spacetime} \\ p = 5 \text{ M5-brane} \\ d-p-2 = 2 \text{ M2-brane} \end{array} \\
 & & \downarrow & & \updownarrow & & \\
 \text{“KK-compactified” } & & \mathcal{Y} & \xrightarrow[\text{magnetic } p\text{-brane charge coupling to electric } d-p-3\text{-branes}]{\text{Cyc}(F)} & \text{Cyc}(\mathbf{B}^{d-p}\mathbb{Z}) & \xrightarrow{\int_{S^1_{\text{coh}}}} & \mathbf{B}^{d-p-1}\mathbb{Z} & \text{e.g.} & \begin{array}{l} 1+d = 10 \text{ type IIA spacetime} \\ p = 5 \text{ NS5-brane} \\ d-p-3 = 1 \text{ NS1-branes (string)} \end{array} \\
 \text{1+d-dimensional spacetime orbifold} & & & & & & & &
 \end{array}$$


Here on the right, we are indicating the example of the double dimensional reduction of M2-brane charge to NS1-brane/string charge (see [MaSa04, §4]) for which we consider a more refined model further below, around (19).

(In a different but related context, a dimensional reduction procedure for supersymmetric Euclidean field theories over an orbifold is proposed in [Sto19].)

It is expected that if such a dimensional reduction is carried out with due care, then it does not lose information and may be reversed (“oxidized”, e.g. [LPSS95]). This is exactly what we find here, formalized by the cyclification adjunction with its hom-isomorphism (14). Notice that this means (i.e.: proves) that brane charges on spacetimes which are iterated principal fiber bundles may be reduced all the way down, possibly all the way to the point, by iterated cyclification, without losing information; see [SV21].

Generalized T-duality. Notice the subtle distinction between the bare cyclification $\text{Cyc}_{\mathcal{T}}(\mathcal{A}) \in \text{SmthGrpd}_{\infty}$ and its incarnation as the base of an \mathcal{T} -principal bundle (50), $\text{Cyc}(\mathcal{A}) \in (\text{SmthGrpd}_{\infty})_{/\mathbf{B}\mathcal{T}}$. The reduction/oxidation-equivalence (14) says that with this \mathcal{T} -bundle data retained, maps from a base space \mathcal{Y} into the cyclification of \mathcal{A} fully recover the \mathcal{A} -valued cocycles on \mathcal{X} . But it may happen that two in-equivalent moduli stacks $\mathcal{A} \neq \mathcal{B}$ have equivalent bare \mathcal{T} -cyclifications, $\text{Cyc}_{\mathcal{T}}(\mathcal{A}) \simeq \text{Cyc}_{\mathcal{T}}(\mathcal{B}) \in \text{SmthGrpd}_{\infty}$.

In this case the two maps c_1^A, c_1^B to $\mathbf{B}\mathcal{T}$ are necessarily in-equivalent, so that $\text{Cyc}(\mathcal{A})$ -valued cocycles on \mathcal{Y} have two different interpretations as cocycles F^A, F^B on different \mathcal{T} -bundles $\mathcal{X}_A, \mathcal{X}_B$. In such a situation, while pairs of cocycles F^A, F^B are in general different (not even of the same type), they are essentially equivalent after reduction along the \mathcal{T} -fibers. As such it makes sense to call them \mathcal{T} -dual to each other:



In the approximation of (super-)rational homotopy theory², this notion of \mathcal{T} -duality has been shown ([FSS16TDI], reviewed in [FSS19Rat, §9]) to reproduce the expected notion of (topological) T-duality from string theory: Here $\mathcal{A} = (\text{KU} // \mathbf{B}^2\mathbb{Z})^{\mathbb{Q}}$ and $\mathcal{B} = (\Sigma\text{KU} // \mathbf{B}^2\mathbb{Z})^{\mathbb{Q}}$ are the rationalizations of the twisted complex K-theory spectra in degree 0 and in degree 1, respectively. The full lift of this situation beyond the rational approximation remains to be discussed elsewhere, but we may readily spell out the comparatively simple but crucial sector of just the *twisting*: by passing along the projection

$$\begin{aligned} \text{KU}_0 // \mathbf{B}^2\mathbb{Z} &\longrightarrow * // \mathbf{B}^2\mathbb{Z} \simeq \mathbf{B}^3\mathbb{Z} \\ \text{Cyc}(\text{KU}_0 // \mathbf{B}^2\mathbb{Z}) &\longrightarrow \text{Cyc}(\mathbf{B}^3\mathbb{Z}) \end{aligned}$$

(i.e., focusing on the the “B-field” while ignoring the “RR-field” for the moment): From (10) and standard facts about homotopy (co-)fibers of maps of Sullivan models (e.g. [FSS16WZW, Prop. 3.5]) one finds that the cyclification of $\mathbf{B}^3\mathbb{Z}$ is the delooping of the shape of the *T-duality 2-group* [FSS12CS, §3.2.1] [FSS16TDI, Def. 7.1],

²Here *super homotopy theory* refers to the ∞ -topos not just over the site of smooth manifolds, as considered in (2), but further embedded into that over super-manifolds [Sc18][Gi23], where (super-)rational homotopy theory is modeled by super-dgc-algebras [HSS18, §3.2] known in the supergravity literature as “FDA”s (see [FSS16WZW]). All our discussion here immediately passes to that context, but for brevity we shall not further dwell on this point here.

defined to be the homotopy fiber of the cup product on degree-2 cohomology:

$$\begin{array}{l}
\text{cyclification of deg=3 cohomology is} \\
\text{classifying space for T-duality pairs} \\
\\
\text{witnessed by homotopy (co)fiber} \\
\text{of cup product on deg=2}
\end{array}
\quad
\begin{array}{c}
\text{Cyc}(\mathbf{B}^3\mathbb{Z}) \xrightarrow{\text{fib}(\cup)} \mathbf{B}^2\mathbb{Z} \times \mathbf{B}^2\mathbb{Z} \xrightarrow{\cup} \mathbf{B}^4\mathbb{Z} \\
\\
\left(\begin{array}{l} d c_3 = c_2 \wedge \omega_2 \\ d c_2 = 0 \\ d \omega_2 = 0 \end{array} \right) \xleftarrow{\text{hocofib}} \left(\begin{array}{l} d c_2 = 0 \\ d \omega_2 = 0 \end{array} \right) \xleftarrow{c_2 \wedge \omega_2 \mapsto c_4} \left(d c_4 = 0 \right) .
\end{array}
\quad (18)$$

This homotopy fiber is known to be the classifying space for topological T-duality pairs ([BuSc05, Thm. 2.17] [Ro09, §6.2][FSS16TDI, Rem. 7.2]); and from (17) we transparently see how this comes about: Given a $\text{Cyc}(\mathbf{B}^3\mathbb{Z})$ -valued cocycle ϕ on a base orbifold \mathcal{Y} , then by iterative application of the pasting law (Fact A.46), we may extract the following pasting diagram of homotopy cartesian squares which makes appear, on the left of the diagram, two total spaces $\mathcal{X}_{A/B}$ carrying “gerbes” $\mathcal{G}_{A/B}$ (here incarnated as $\mathbf{B}^2\mathbb{Z}$ -principal ∞ -bundles) subject to the constraint that their pullback to the correspondence orbifold $\mathcal{X}_A \times_{\mathcal{Y}} \mathcal{X}_B$ are both equivalent to the “Poincaré gerbe” \mathcal{G} :

This recovers the basic axiomatics [BuSc05] of “topological T-duality” (going back to [BEM04], good review is in [Wa22, §1]) in the familiar situation and generalizes it to orbifolds and higher stacks.

By way of further outlook, we close this overview by indicating the following concrete application, which deserves to be discussed in more detail elsewhere:

Application to brane physics. While much of the interest in elliptic cohomology goes back to the suggestion [Wit88] that the partition function of the heterotic string is an elliptic genus of the string’s target spacetime \mathcal{X} , realized via S^1 -equivariant K-theory of its free loop space $\text{Map}(S^1_{\text{coh}}, \mathcal{X})$, only a couple of authors ([KS04][KS05]) have investigated the idea that elliptic cohomology might relate to the widely expected hypothesis that the topological K-theory of \mathcal{X} itself measures the brane charges in (not heterotic but) type I/II string theories (a comprehensive list of references for this classical hypothesis may be found in [BSS18, §1]).

One circumstantial hint for what may really be going on comes the fact that the elliptic genus of the heterotic string is expected (at the physics level of rigor) to really just be a special case of the general notion of elliptic genera of toroidally compactified super-branes, notably of the M5-brane [GSY07][GY07][AHHKRW15] [GPPV21]. This leads us to ask whether elliptic cohomology measures aspects of M-brane charges?

One coherent answer to what may be going on is suggested by *Hypothesis H* [FSS19HypH][SS20Tad][SS21MF]: This postulates that the charges of branes in M-theory are measured in tangentially $\text{Sp}(2) \simeq \text{Spin}(5)$ -twisted equivariant unstable 4-Cohomotopy (see [FSS20Cha] for details), i.e. in the non-abelian cohomology theory whose plain moduli stack is the 4-sphere homotopy quotiented by its canonical $\text{Spin}(5) \rightarrow \text{SO}(5)$ -action, with the ordinary cohomology charge shown in (16) being only a subtle integral characteristic class $\tilde{\Gamma}_4$ ([FSS19HypH, Lem. 3.12][FSS22GS, (7)][FSS21Str, (4)]) of this twisted non-abelian cohomology theory:

(19)

If \mathcal{X} is an S^1_{coh} -principal bundle over a 10d spacetime orbifold \mathcal{Y} , as usually assumed (starting with [DHIS87] [Wi95, p. 10], see [MaSa04, p. 2]), then the Cyc-adjunction (13) says that such charges on \mathcal{X} are equivalently twisted equivariant $\text{Cyc}(S^4)$ -valued charges on \mathcal{Y} . Curiously, these cyclic 4-sphere coefficients are close to the twisted K-theory expected for D-brane charge ([FSS17Sph, Ex. 3.3][BSS18, Ex. 2.47]):

$$\begin{array}{ccc} \text{rational 6-truncation of} & & \text{rational 6-truncation of} \\ \text{cyclified 4-sphere} & & \text{twisted K-theory spectrum} \\ \text{[Cyc}(S^4)\text{]}_6^{\mathbb{Q}} & \simeq & \text{[(Fred}_{\mathbb{C}}^0\text{//PU}(\mathcal{H}))\text{]}_6^{\mathbb{Q}}. \end{array}$$

This suggests that the pullback of the **twisted ADE-equivariant quasi-elliptic cohomology of the 4-sphere** along twisted Cohomotopy cocycles produces good observables on M-brane charge in a Tate-elliptic enhancement of D-brane charge in twisted equivariant K-theory (for which our notation follows [SS21EPB, Ex. 4.5.4]):

(20)

We compute the relevant twisting 4-class below in §4, cf. Rem. 4.8. In order to further analyze (20), one will need to first compute the equivariant quasi-elliptic cohomology of representation 4-spheres of finite subgroups of $SU(2)$ (Prop. A.42) twisted by the resulting transgressed 3-class $\int_{S^1} \text{Cyc}(\tilde{\Gamma}_4)$. Of course, this is just one of the “twisted equivariant homotopy-groups” (rather: “-modules”, due to the non-trivial twist) of quasi-elliptic cohomology, which are bound to be of fundamental interest in their own right. We leave their computation to the quasi-elliptic community.

Outline.

In §2 we prove that the abstract cyclification construction (5) recovers GRH’s component model (Thm. 2.3). In §3 we prove that the abstract transgression operation (11) recovers traditional component formulas (Thm. 3.4). In §4 we compute the integral 4-class (transgressing to a 3-class) to be used in (20) for measuring M5-brane genera. In appendix A we compile some technical background in simplicial & geometric homotopy theory, for reference.

2 Cyclic inertia orbifolds

In §2.1 we recall the “extended” inertia orbifolds due to Ganter, Rezk & Huan (GRH);
in §2.2 we present a general abstract theory of cyclic inertia ∞ -groupoids;
in §2.3 we prove that the GRH construction models the abstract definition.

2.1 GRH’s extended inertia orbifold

Smooth loop and inertia stacks. Given an orbifold or, more generally, any smooth ∞ -groupoid $\mathcal{X} \in \text{SmthGrpd}_\infty$ (see Nota. A.48) it is well-known (albeit not always stated in the following model-independent stack-theoretic manner³) that:

(i) its *smooth loop stack* ([LU02, §3][BGNX07, §5]) is the mapping stack out of the smooth circle $S_{\text{coh}}^1 \in \text{SmthMnflid} \xrightarrow{y} \text{SmthGrpd}_\infty$:

$$\text{smooth loop stack } \mathcal{L}\mathcal{X} := \text{Map}(S_{\text{coh}}^1, \mathcal{X}) \in \text{SmthGrpd}_\infty \quad (21)$$

(ii) its *inertia stack* (e.g. [LU04, §4]) is the mapping stack out of the delooping groupoid of the integers $\mathbf{B}\mathbb{Z} \in \text{Grpd}_\infty \xrightarrow{\text{Disc}} \text{SmthGrpd}_\infty$:

$$\text{inertia stack } \Lambda\mathcal{X} := \text{Map}(\mathbf{B}\mathbb{Z}, \mathcal{X}) \in \text{SmthGrpd}_\infty. \quad (22)$$

Remark 2.1 (Components of inertia). In the special case of *good orbifolds*, or, more generally, of good diffeological orbispaces [SS20Orb, §4],

$$\mathcal{X} \text{ is good} \Leftrightarrow \exists_{\substack{G \in \text{Grp}(\text{Set}) \\ G \curvearrowright X \in G\text{Act}(\text{DfflgSpc})}} \mathcal{X} \simeq X // G, \quad (23)$$

the inertia stack is readily found (and well-known) to be equivalent to a disjoint union over the conjugacy classes $[g] \in G/\text{ad} G$ of the corresponding fixed loci $X^g \subset X$ by their residual action of the centralizer subgroups $C_g \subset G$, as follows⁴

$$G \curvearrowright X \in G\text{Act}(\text{DfflgSpc}) \text{ and } \mathcal{X} \simeq X // G \quad \Rightarrow \quad \Lambda\mathcal{X} \simeq \coprod_{[g] \in G/\text{ad} G} X^g // C_g. \quad (24)$$

inertia orbifold
skeletal presentation

Remark 2.2 (Cohesion knows about essentially constant loops). The difference and the relation between these two constructions (21) and (22) is brought out by the shape modality: The shape (107) of the cohesive circle is (Lemma 2.8) equivalently the delooping groupoid of the integers

$$\int S_{\text{coh}}^1 \simeq \mathbf{B}\mathbb{Z} \in \text{SmthGrpd}_\infty$$

so that we may understand the inertia stack (22) as being a loop stack itself, but for loops that are just the *bare shape* of a smooth circle:

$$\underbrace{\Lambda\mathcal{X} \simeq \text{Map}(\int S_{\text{coh}}^1, \mathcal{X}) \xrightarrow{\text{Map}(\eta_{\int}, \mathcal{X})} \text{Map}(S_{\text{coh}}^1, \mathcal{X}) \simeq \mathcal{L}\mathcal{X}}_{\text{inclusion of cohesively constant loops}} \quad (25)$$

inertia stack
 $\int S_{\text{coh}}^1$ -loop stack
shape unit
 S_{coh}^1 -loop stack
smooth loop stack

It is this cohesive relation that formalizes the traditional notion that the inertia stack is the restriction of the smooth loop stack to the essentially constant loops: The “bare shape” of a loop cannot traverse any non-constant path in a manifold, but it can still jump between the “twisted sectors” to which the point belongs that it is constantly sitting on.

³Regarding S_{coh}^1 as an object of SmthGrpd_∞ ensures that stacky maps out of it are modeled by simplicial maps out of any good open cover (cf. [SS21EPB, Ex. 3.3.41]), which is what gives rise to component constructions such as in [LU02, Def. 3.1.1].

⁴If $G \curvearrowright X$ is a proper action on a smooth manifold, then the fixed loci X^g are themselves smooth manifolds, so that the inertia stack is again a good orbifold. But the equivalence (24) holds more generally, as shown, for $\mathcal{X} \simeq X // G$ any *good diffeological orbi-space*, where X may be any *diffeological space* with smooth G -action, faithfully subsuming finite-dimensional smooth manifolds as well as infinite-dimensional Fréchet-manifolds. This is a convenient generalization, as the smooth loop stack construction (21) restricts to an endo-functor (2-functor) on diffeological orbispaces, where it may hence be iterated.

Cyclic loop spaces and cyclic homology. Closely related to (25) is the fact that for $\mathcal{X} = \int(X)$ the shape (107) of a topological space $X \in \text{TopSp} \xrightarrow{\text{Cdfflg}} \text{SmthGrpd}_\infty$ [SS20Orb, Ex. 3.18] its inertia ∞ -stack (22) is the shape (by the *smooth Oka principle*, cf. [SS21EPB, p. 7]) of the topological free loop space of X (e.g. [CO15]):

$$\Lambda \int X \simeq \int \text{Maps}(S_{\text{coh}}^1, X).$$

In this context it is a familiar idea ([Jo87, Thm. A], review in [Lo92, Cor. 7.3.14][Lo15, §3,4]) that associated to the free topological loop space is what we call its *cyclic loop space* [FSS16TD1, §3][BSS18, §2.2], namely the homotopy quotient (Borel construction) by the circle action which rigidly rotates the loops:

$$\begin{aligned} \text{Cyc}(X) &:= \text{Maps}(S_{\text{coh}}^1, X) // S_{\text{coh}}^1 \simeq \left(\text{Maps}(S_{\text{coh}}^1, X) \times ES_{\text{coh}}^1 \right) / S_{\text{coh}}^1, \\ H^\bullet(\text{Cyc}(X)) &\simeq \text{HC}_\bullet(C^\bullet(X)), \end{aligned}$$

whose ordinary cohomology (for any commutative coefficient ring R) is the cyclic homology of (the graded R -algebra of R -chains of) X .

Cyclic loop stacks. Similarly, it is clear that the loop stack (21) carries a canonical smooth action of the circle group $S_{\text{coh}}^1 \simeq U(1)$ by rotation of loops (component constructions are given in [LU02, §3.6], the general abstract construction follows by (31) below), so that, in view of (25), we may consider the further homotopy quotient by this action, which we will denote as

$$\text{Cyc}_{S_{\text{coh}}^1}(\mathcal{X}) := \text{Map}(S_{\text{coh}}^1, \mathcal{X}) // S_{\text{coh}}^1 \in \text{SmthGrpd}_\infty. \quad (26)$$

This fundamental construction has not received much attention (it is alluded to in [Ga07, §2.1][St15, p. 2]) until the recent introduction of modified orbifold loop groupoids in [Hua18a] (denoted “ $Loop^{\text{ext}}$ ” in Def. 2.5, 2.9 there) which we may understand as plausible models for the homotopy quotient in (26). Restricting these ad-hoc models to cohesively constant loops leads to a definition ([Hua18a, Def. 2.14] in slight variation of [Ga07, Def. 2.3], review in [Do19, p. 62][HS20, Def. 2.1]) of a variation of the inertia orbifold (24), as follows:

$$\Lambda_{S_{\text{coh}}^1}^{\text{GRH inertia orbifold}} \mathcal{X} := \coprod_{[g] \in G/\text{ad}} X^g // \Lambda_g, \quad \text{where } \Lambda_g := \frac{C_g \times \mathbb{R}}{\langle (g^{-1}, 1) \rangle}. \quad (27)$$

In [Hua18a] and followups ([Hua18b][HS20]), this definition is justified *a posteriori* by the fact that a completion of the orbifold K-theory of GRH’s inertia orbifold (27) yields a good model of the equivariant elliptic cohomology at the Tate curve of the original orbifold:

$$\text{KU}_{\text{orb}}(\Lambda_{S_{\text{coh}}^1}^{\text{GRH's extended inertia orbifold}} \mathcal{X}) \otimes_{\mathbb{Z}[q, q^{-1}]} \mathbb{Z}((q)) \simeq \text{Ell}_{\text{orb}}^{\text{orbifold Tate-elliptic cohomology}}(\mathcal{X}). \quad (28)$$

2.2 Cyclic inertia ∞ -Groupoids

A general theory of stacky transformation ∞ -groups. Our starting point is the observation [NSS12a][SS20Orb, §2.2] that the general theory of *transformation groups*, i.e. of geometric (topological-, Lie-, ...) groups acting as symmetries of geometric spaces ([Br72][tD79][tD87]), not only has a good generalization to groupal ∞ -stacks, but becomes conceptually more transparent in this generalization, when regarded systematically topic in *higher topos theory* ([TV05][Lu09][Re10]). Namely, [SS21EPB, Prop. 0.2.1]:

(i) for $\mathcal{T} \in \text{Grp}(\mathbf{H})$ a group object in any ∞ -topos \mathbf{H} (Ntn. A.45), the ∞ -category of \mathcal{T} -actions on ∞ -stacks is equivalent to the *slice ∞ -topos* over the \mathcal{T} -moduli ∞ -stack $\mathbf{B}\mathcal{T}$ ([SS21EPB, Prop. 3.2.76], going back to [NSS12a, §4.1] and [DDK80], see also Prop. A.34 below),

(ii) for $\mathcal{T}_1 \xrightarrow{\phi} \mathcal{T}_2$ any homomorphism of such ∞ -groups, the constructions of *restricted* and of *(co-)induced* ∞ -actions along ϕ are equivalently given by the base change adjoint triple (106) along ϕ ([SS21EPB, Ex. 3.2.78]):

⁵The quotient group Λ_g in (27) may be motivated (as explained in [Do19, (6.5)]) as that which implements the “rotation condition” proposed in [Ga13, Def. 2.6].

$$\begin{array}{ccc}
& \xrightarrow{\text{induced } \infty\text{-action}} & \\
\infty\text{-actions of } \mathcal{T}_1 \text{ on } \infty\text{-stacks} & \mathcal{T}_1 \text{Act}(\mathbf{H}) & \xleftarrow{\text{restricted } \infty\text{-action}} \mathcal{T}_2 \text{Act}(\mathbf{H}) & \infty\text{-actions of } \mathcal{T}_2 \text{ on } \infty\text{-stacks} \\
& \xleftarrow{\text{co-induced } \infty\text{-action}} & \\
& \xrightarrow{\text{left base change}} & \\
& \xrightarrow{(\mathbf{B}\phi)_!} & \\
\infty\text{-stacks over } \mathcal{T}_1\text{-moduli} & \mathbf{H}/\mathbf{B}\mathcal{T}_1 & \xleftarrow{\text{base change/pullback } (\mathbf{B}\phi)^*} \mathbf{H}/\mathbf{B}\mathcal{T}_2 & \infty\text{-stacks over } \mathcal{T}_2\text{-moduli} \\
& \xrightarrow{(\mathbf{B}\phi)_*} & \\
& \xrightarrow{\text{right base change}} &
\end{array}
\tag{29}$$

Cyclic ∞ -Stacks. Specializing (29) to the unique homomorphism from the trivial group $1 \xrightarrow{e} \mathcal{T}$, where $\mathbf{B}e = \text{pt}_{\mathbf{B}\mathcal{T}}$ is the base point inclusion into the \mathcal{T} -moduli ∞ -stack, we see that, in full generality, the construction of \mathcal{T} -cyclic ∞ -stacks

$$\text{Cyc}_{\mathcal{T}}(\mathcal{X}) := \text{Map}(\mathcal{T}, \mathcal{X}) // \mathcal{T} \in \mathbf{H}/\mathbf{B}\mathcal{T} \xrightarrow{(p_{\mathbf{B}\mathcal{T}})_!} \mathbf{H} \tag{30}$$

is equivalently the right base change along the point inclusion into the \mathcal{T} -moduli ∞ -stack (derived left base change back to the global context, as desired):

$$\begin{array}{ccccc}
& \xrightarrow{\text{free } \infty\text{-action}} & \mathcal{T} \wr (\mathcal{T} \times (-)) & \xrightarrow{\text{homotopy co-invariants}} & (-)_{\mathcal{T}} & \xrightarrow{\text{homotopy co-invariants}} & \mathbf{H} \\
\infty\text{-stacks } \mathbf{H} & \xleftarrow{\text{underlying } \infty\text{-stack}} & \mathcal{T} \text{Act}(\mathbf{H}) & \xleftarrow{\text{trivial } \infty\text{-action}} & \mathbf{H} & \xleftarrow{\text{trivial } \infty\text{-action}} & \mathbf{H} \\
& \xrightarrow{\text{co-free } \infty\text{-action}} & \mathcal{T} \wr [\mathcal{T}, -] & \xrightarrow{\text{homotopy invariants}} & (-)_{\mathcal{T}} & \xrightarrow{\text{homotopy invariants}} & \mathbf{H} \\
& & \mathcal{T} \wr \mathcal{X} & \xrightarrow{\text{left base change to absolute context}} & (p_{\mathbf{B}\mathcal{T}})_! & \xrightarrow{\text{left base change to absolute context}} & \mathbf{H} \\
& & \mathcal{X} // \mathcal{T} & \xrightarrow{\text{extension = base change}} & \text{Ext}_{\mathcal{T}} := (\text{pt}_{\mathbf{B}\mathcal{T}})^* & \xrightarrow{\text{extension = base change}} & \mathbf{H}/\mathbf{B}\mathcal{T} \\
& & & \xrightarrow{\text{cyclification = right base change}} & \text{Cyc}_{\mathcal{T}} := (\text{pt}_{\mathbf{B}\mathcal{T}})_* & \xrightarrow{\text{cyclification = right base change}} & \mathbf{H}/\mathbf{B}\mathcal{T} \\
& & & & \text{base point inclusion} & & \text{projection to base point} \\
& & & & \mathbf{B}\mathcal{T} & & \mathbf{B}\mathcal{T}
\end{array}
\tag{31}$$

In the top left and in many following diagrams we abbreviate our notation for mapping stacks to

$$[-, -] := \text{Map}(-, -).$$

Notice how this ∞ -group-theoretic adjunction (31) witnesses shaundamental aspects of loop stacks and their cyclification:

Object in:	Mapping ∞ -Stack Theory	Transf. ∞ -Group Theory	Slice ∞ -Topos Theory
$\text{Map}(\mathcal{T}, \mathcal{X})$	\mathcal{T} -loop ∞ -stack of \mathcal{X}	underlying ∞ -stack of co-free \mathcal{T} - ∞ -action co-induced by \mathcal{X}	comonadic descent of \mathcal{X} along base point into \mathcal{T} -moduli ∞ -stack
$\text{Map}(\mathcal{T}, \mathcal{X}) // \mathcal{T}$	\mathcal{T} -cyclic ∞ -stack of \mathcal{X}	homotopy quotient by co-free \mathcal{T} - ∞ -action co-induced by \mathcal{X}	right derived base change of \mathcal{X} along base point into \mathcal{T} -moduli ∞ -stack

Inertia ∞ -Stacks. Moreover, if \mathbf{H} is cohesive (Ntn. A.48) and thus equipped with a shape modality (107), then with \mathcal{T} also its shape $\int \mathcal{T}$ is canonically a group ∞ -stack ([SS20Orb, Prop. 3.4]) and the shape unit $\mathcal{T} - \eta_{\mathcal{T}}^{\int} \rightarrow \int \mathcal{T}$ is a homomorphism of group ∞ -stacks, so that we may consider \mathcal{T} -inertia ∞ -stacks and their \mathcal{T} -cyclification in full generality:

$$\begin{array}{ccc}
\mathcal{T}\text{-inertia } \infty\text{-stack} & (\int \mathcal{T}) \hookrightarrow \text{Map}((\int \mathcal{T}), \mathcal{X}) \xrightarrow[\text{along shape unit}]{\text{restricted } \infty\text{-action}} \mathcal{T} \hookrightarrow \text{Map}((\int \mathcal{T}), \mathcal{X}) & \\
& \downarrow \text{homotopy quotient} & \downarrow \text{homotopy quotient} \\
\text{cyclic } \mathcal{T}\text{-inertia } \infty\text{-stacks} & \text{Map}((\int \mathcal{T}), \mathcal{X}) // (\int \mathcal{T}) \xrightarrow{(\eta_{\int \mathcal{T}}^{\int})^*} \text{Map}((\int \mathcal{T}), \mathcal{X}) // \mathcal{T} &
\end{array} \tag{32}$$

Notice that we obtain a natural comparison morphism, as claimed in (4), from any cyclic \mathcal{T} -inertia stack (32) to the full \mathcal{T} -cyclification (30) by factoring the defining base change through the decomposition

$$\begin{array}{ccc}
* & \xrightarrow{\text{pt}_{\mathbf{B}\mathcal{T}}} \mathbf{B}\mathcal{T} & \xrightarrow{\mathbf{B}\eta_{\mathbf{B}\mathcal{T}}^{\int}} \mathbf{B}\int \mathcal{T} \\
& \searrow \text{pt}_{\mathbf{B}\int \mathcal{T}} & \nearrow \\
& &
\end{array}$$

and invoking the counit $\varepsilon : f^* f_* \rightarrow \text{id}$ of the right base change adjunction (106):

$$\begin{array}{ccccc}
\text{Cyc}_{\int \mathcal{T}}(\mathcal{X}) & & & & \text{Cyc}_{\mathcal{T}}(\mathcal{X}) \\
\parallel & & & & \parallel \\
\text{Map}(\int \mathcal{T}, \mathcal{X}) // \int \mathcal{T} & \text{Map}(\int \mathcal{T}, \mathcal{X}) // \mathcal{T} & \xrightarrow{\text{-----}} & \text{Map}(\mathcal{T}, \mathcal{X}) // \mathcal{T} & \\
\parallel & \parallel & & \parallel & \\
(\text{pt}_{\mathbf{B}\int \mathcal{T}})_*(\mathcal{X}) & \xleftarrow{\quad} (\mathbf{B}\eta_{\mathbf{B}\int \mathcal{T}}^{\int})^*(\text{pt}_{\mathbf{B}\int \mathcal{T}})_*(\mathcal{X}) \simeq (\mathbf{B}\eta_{\mathbf{B}\mathcal{T}}^{\int})^*(\mathbf{B}\eta_{\mathbf{B}\mathcal{T}}^{\int})_*(\text{pt}_{\mathbf{B}\mathcal{T}})_*(\mathcal{X}) \xrightarrow{\varepsilon_{(\text{pt}_{\mathbf{B}\mathcal{T}})_*(\mathcal{X})}} (\text{pt}_{\mathbf{B}\mathcal{T}})_*(\mathcal{X}) & & & \\
\downarrow & \text{(pb)} \downarrow & & \downarrow & \\
\mathbf{B}\int \mathcal{T} & \xleftarrow{\mathbf{B}\eta_{\mathbf{B}\mathcal{T}}^{\int}} \mathbf{B}\mathcal{T} & \xlongequal{\quad\quad\quad} & \mathbf{B}\mathcal{T} &
\end{array} \tag{33}$$

Theorem 2.3 (GRH's extended inertia groupoid models the cyclified orbifold). *For $\mathcal{X} \simeq X // G \in \text{SmthGrpd}_{\infty}$ any good orbifold (23), GRH's inertia orbifold (27) is equivalently the $\mathcal{T} := S_{\text{coh}}^1$ -cyclic inertia ∞ -stack in the general sense of (32):*

$$\Lambda_{S_{\text{coh}}^1}^{\mathcal{X}} \simeq \text{Map}(\int S_{\text{coh}}^1, \mathcal{X}) // S_{\text{coh}}^1 =: \underbrace{(\eta_{\mathbf{B}S_{\text{coh}}^1}^{\int})^*}_{\substack{\text{restrict action} \\ \text{from shape of circle} \\ \text{to full smooth circle}}} \text{Cyc}_{\int S_{\text{coh}}^1}(\mathcal{X}).$$

Proof. In view of (29), this follows from Prop. 2.6, discussed in detail in §2.3 below. \square

Remark 2.4 (Subtleties). While the idea that Thm. 2.3 should be true possibly motivated the definition (27), its proof requires some care (see the proof of Lemma 2.5 below). Notice that the approach via a shape modality on higher stacks is crucial in bringing out this result: Our diagram (31) shows at once that the traditional way of identifying the inertia stack inside the smooth loop stack as the S_{coh}^1 -fixed locus in a suitable groupoid presentation (following [LU02, Thm. 3.6.4]) is *not homotopy-meaningful*, as the *homotopy*-fixed locus (e.g. [SS21EPB, Ex. 3.2.78]) of every \mathcal{T} -loop ∞ -stack is just the original ∞ -stack:

$$(\text{Map}(\mathcal{T}, \mathcal{X}))^{\mathcal{T}} \simeq (p_{\mathbf{B}\mathcal{T}})_*(\text{pt}_{\mathbf{B}\mathcal{T}})_*(\mathcal{X}) \simeq (\text{id})_*(\mathcal{X}) \simeq \mathcal{X}, \tag{34}$$

since, by (31), it ends up computing the right base change of \mathcal{X} along the identity:

$$\begin{array}{ccc}
* & \xrightarrow{\text{pt}_{\mathbf{B}\mathcal{T}}} \mathbf{B}\mathcal{T} & \xrightarrow{p_{\mathbf{B}\mathcal{T}}} * \\
& \searrow \text{id} & \nearrow \\
& &
\end{array}$$

2.3 Reproducing GRH's extended inertia orbifold

Here we prove Theorem 2.3, that GRH's inertia orbifold construction is a model for an abstractly defined cyclic inertia orbifold: $\Lambda_{S_{\text{coh}}^1} \mathcal{X} \simeq \text{Map}(\mathcal{J}S_{\text{coh}}^1, \mathcal{X}) // S_{\text{coh}}^1$.

Lemma 2.5 (Comparison morphism from GRH's inertia orbifold to cyclic orbifold). *For $\mathcal{X} \simeq X // G$ a good orbifold, there is a comparison morphism, as shown in (37) and (38),*

$$\Lambda_{S_{\text{coh}}^1}(\mathcal{X}) \xrightarrow{\text{comp}_{\mathcal{X}}} \text{Cyc}_{\mathcal{J}S_{\text{coh}}^1}(\mathcal{X}) = \text{Map}(\mathcal{J}S_{\text{coh}}^1, \mathcal{X}) // \mathcal{J}S_{\text{coh}}^1 \quad (35)$$

from GRH's extended inertia orbifold (27) to the $\mathcal{J}S_{\text{coh}}^1$ -cyclification (32) of the inertia orbifold.

Proof. Shown in (37) and (38) (on the next two pages) is a morphism of simplicial presheaves. We need to see that:

- (i) this is well-defined as a morphism in $\Delta\text{Sh}(\text{CartSpc})$,
- (ii) under localization it presents a morphism in SmthGrpd_{∞} of the claimed form (35).

Regarding (i): On both sides of (37) and (38) we show the diagonal quotient $((-) \times W\mathcal{G})/\mathcal{G}$ (108) of the Cartesian product with a universal simplicial classifying space $W\mathcal{G}$ Def. A.20 by the diagonal action of a simplicial group \mathcal{G} , all extended to simplicial presheaves over CartSpc .

On the left of (37) and (38), the simplicial group $\mathcal{G} = C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet}$ is the cofibrant resolution (41) of GRH's centralizer group with its induced simplicial action on the fixed loci:

$$\begin{array}{ccc} X^g \times (C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet}) & \xrightarrow{(-) \cdot (-)} & X^g \\ (x, (h, r, \vec{n})) & \longmapsto & x \cdot h \end{array}$$

On the right of (37) and (38), the simplicial group is $\mathcal{G} = \mathbb{Z}^{\times \bullet}$, and its action is on the skeleton of the inertia hom-complex via (43):

$$\begin{array}{ccc} [\mathbf{B}\mathbb{Z} \times \Delta[\bullet], X // G]_{\text{skel}} \times \mathbb{Z}^{\times \bullet} & \longrightarrow & [\mathbf{B}\mathbb{Z} \times \Delta[\bullet], X // G]_{\text{skel}} \\ ((x, (\dots, h_1, h_0)), (\dots, n_1, n_0)) & \longmapsto & (x, (\dots, g^{n_1} \cdot h_1, g^{n_0} \cdot h_0)) \end{array} \quad (36)$$

The bulk of the diagrams (37) and (38) shows the face maps – using (80) – and the degeneracy maps – using (81) – of the resulting quotients on both sides, to check that the comparison morphism indeed respects these. For illustration of how these maps are obtained, we spell out the computation of the one on the bottom left of (37):

$$\begin{array}{c} \xrightarrow{d_0} \\ \begin{array}{ccccc} X^g \times (C_g \times \mathbb{R}) & \xrightarrow{\sim} & \frac{\overbrace{X^g}_{{(X^g)_1}} \times \overbrace{(C_g \times \mathbb{R} \times \mathbb{Z}) \times (C_g \times \mathbb{R})}^{W(C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet})_1}}{\underbrace{C_g \times \mathbb{R} \times \mathbb{Z}}_{{(C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet})_1}}} & \xrightarrow{\frac{d_0^{X^g} \times d_0^{W(\mathcal{G})}}{d_0^{\mathcal{G}}}} & \frac{\overbrace{X^g}_{{(X^g)_0}} \times \overbrace{C_g \times \mathbb{R}}^{W(C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet})_0}}{\underbrace{C_g \times \mathbb{R}}_{{(C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet})_0}}} & \xrightarrow{\sim} & X^g \\ (x, (h, r)) & \longmapsto & [x, (e, 0, 0), (h, r)] & \longmapsto & [x, (e, 0) \cdot (h, r)] = [x \cdot h, (e, 0)] & \longmapsto & x \cdot h \end{array} \end{array}$$

Proceeding this way, one checks (see the next two pages) that all parallel squares in (37) and (38) indeed commute.

smooth
∞-stacks

$$\Lambda_{S^1_{\text{coh}}}^{\text{GRH's inertia}}(X // G) := \coprod_{[g]} \left(X^g // ((C_g \times \mathbb{R}) // (g^{-1}, 1)) \right) \xrightarrow[\simeq_f]{\text{comparison morphism } \text{comp}_{X // G}} [fS^1_{\text{coh}}, X // G] // fS^1_{\text{coh}} =: \text{Cyc}_{fS^1_{\text{coh}}}(X // G) \quad \text{FSS-cyclification}$$

simplicial
sheaves

$$\coprod_{[g]} \left((X^g \times W(C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet})) // (C_g \times \mathbb{R} \times \mathbb{Z}^{\times \bullet}) \right) \longrightarrow \left([\mathbb{Z}^{\times \bullet}, X \times G^{\times \bullet}]_{\text{skel}} \times W(\mathbb{Z}^{\times \bullet}) \right) // \mathbb{Z}^{\times \bullet}$$

deg 4

$$\coprod_{[g]} \left(X^g \times (C_g \times \mathbb{R} \times \mathbb{Z}^3) \times (C_g \times \mathbb{R} \times \mathbb{Z}^2) \right) \xrightarrow[\substack{x \mapsto x \\ h_3 \mapsto h_3 \\ h_2 \mapsto h_2 \cdot g^{-n_{3,2}} \\ h_1 \mapsto h_1 \cdot g^{-n_{3,1} - n_{2,1}} \\ h_0 \mapsto h_0 \cdot g^{-n_{3,0} - n_{2,0} - n_1} \\ n_{(-)} \mapsto n_{(-)}}]{\text{comparison morphism}} \coprod_{[g]} (X^g \times C_g \times C_g \times C_g \times \mathbb{Z}^3 \times \mathbb{Z}^2 \times \mathbb{Z})$$

degeneracy maps

$$\begin{array}{ccc} \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (e, 0, 0, 0), \\ (h_2, r_2, n_{2,1}, n_{2,0}), \\ (h_1, r_1, n_1), \\ (h_0, r_0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, (h_2, r_2, 0, n_{2,1}, n_{2,0}), \\ (e, 0, 0, 0), (h_1, r_1, n_1), \\ (h_0, r_0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h_2, r_2, n_{2,1}, n_{2,0}), \\ (h_1, r_1, n_1), \\ (h_0, r_0) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (h_2, r_2, n_{2,1}, 0, n_{2,0}), \\ (h_1, r_1, 0, n_1), (e, 0, 0), \\ (h_0, r_0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, (h_2, r_2, n_{2,1}, n_{2,0}, 0), \\ (h_1, r_1, n_1, 0), \\ (h_0, r_0), (e, 0, 0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h_2, r_2, n_{2,1}, n_{2,0}), \\ (h_1, r_1, n_1), \\ (h_0, r_0) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} (x, e, h_2, h_1, h_0), \\ (0, 0, 0), (n_{2,1}, n_{2,0}), \\ n_1 \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} (x, h_2, e, h_1, h_0), \\ (0, n_{2,1}, n_{2,0}), \\ (0, 0), n_1 \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} (x, h_2, h_1, h_0), \\ (n_{2,1}, n_{2,0}, 0), \\ (n_1, 0), 0 \end{array} \right) \end{array} \end{array}$$

deg 3

$$\coprod_{[g]} \left(X^g \times (C_g \times \mathbb{R} \times \mathbb{Z}^2) \times (C_g \times \mathbb{R} \times \mathbb{Z}) \times (C_g \times \mathbb{R}) \right) \xrightarrow[\substack{x \mapsto x \\ h_2 \mapsto h_2 \\ h_1 \mapsto h_1 \cdot g^{-n_{2,1}} \\ h_0 \mapsto h_0 \cdot g^{-n_{2,0} - n_1} \\ n_{(-)} \mapsto n_{(-)}}]{\text{comparison morphism}} \coprod_{[g]} (X^g \times C_g \times C_g \times C_g \times \mathbb{Z}^2 \times \mathbb{Z})$$

degeneracy maps

$$\begin{array}{ccc} \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (e, 0, 0, 0), \\ (h_1, r_1, n), \\ (h_0, r_0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h_1, r_1, n), \\ (h_0, r_0) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (h_1, r_1, 0, n), \\ (e, 0, 0), \\ (h_0, r_0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h_1, r_1, n), \\ (h_0, r_0) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (h_1, r_1, n, 0), \\ (h_0, r_0, 0), \\ (e, 0) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h_1, r_1, n), \\ (h_0, r_0) \end{array} \right) \end{array} \end{array}$$

deg 2

$$\coprod_{[g]} (X^g \times (C_g \times \mathbb{R} \times \mathbb{Z}) \times (C_g \times \mathbb{R})) \xrightarrow[\substack{x \mapsto x \\ h_1 \mapsto h_1 \\ h_0 \mapsto h_0 \cdot g^{-n} \\ n \mapsto n}]{\text{comparison morphism}} \coprod_{[g]} (X^g \times C_g \times C_g \times \mathbb{Z})$$

degeneracy maps

$$\begin{array}{ccc} \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (e, 0, 0), (h, r) \\ \uparrow \\ (x, (h, r)) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h, r) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (h, r, 0), (e, 0) \\ \uparrow \\ (x, (h, r)) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ (h, r) \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} (x, e, h), 0 \\ \uparrow \\ (x, h) \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} (x, h, e), 0 \\ \uparrow \\ (x, h) \end{array} \right) \end{array} \end{array}$$

deg 1

$$\coprod_{[g]} (X^g \times C_g \times \mathbb{R}) \xrightarrow[\substack{x \mapsto x \\ h \mapsto h \\ n \mapsto n}]{\text{comparison morphism}} \coprod_{[g]} (X^g \times C_g)$$

degeneracy maps

$$\begin{array}{ccc} \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, (e, 0) \\ \uparrow \\ x \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ x \end{array} \right) \end{array} & \begin{array}{c} \uparrow \\ \left(\begin{array}{c} x, e \\ \uparrow \\ x \end{array} \right) \\ \uparrow \\ \left(\begin{array}{c} x, \\ x \end{array} \right) \end{array} \end{array}$$

deg 0

$$\coprod_{[g]} (X^g) \xrightarrow{x \mapsto x} \coprod_{[g]} (X^g)$$

(38)

Regarding (ii): It remains to show that the morphism of simplicial presheaves shown on the previous two pages, in (37) and (38), indeed represents a morphism of SmothGrpd_∞ of the form (35). That the action on the right is the correct one is the content of Lemma 2.9. The simplicial Borel constructions on both sides of (37) and (38) present the respective homotopy quotients by [NSS12a, §3.5] (recalled as [SS21EPB, Lem. 3.2.73]). \square

Proposition 2.6 (Comparison morphism is pullback of shape unit). *The comparison morphism from Lemma 2.5 sits in a homotopy-Cartesian square as follows (thus implying Thm. 2.3):*

$$\begin{array}{ccc} \Lambda_{S^1_{\text{coh}}}(X//G) & \xrightarrow{\text{comp}_{X//G}} & \text{Cyc}_{\mathcal{J}S^1_{\text{coh}}}(X//G) \\ \downarrow & \text{(pb)} & \downarrow \\ \mathbf{BS}^1_{\text{coh}} & \xrightarrow{\eta_{S^1_{\text{coh}}}} & \mathcal{J}\mathbf{BS}^1_{\text{coh}} \end{array} \Leftrightarrow \Lambda_{S^1_{\text{coh}}}(X//G) \simeq (\eta_{S^1_{\text{coh}}}^{\mathcal{J}})^* \text{Cyc}_{\mathcal{J}S^1_{\text{coh}}}(X//G). \quad (39)$$

Proof. A glance at the component maps in (37) and (38) reveals that the comparison morphism fails to be a degreewise isomorphism of simplicial presheaves only in that the elements $r \in \mathbb{R}$ on the left hand side are forgotten. Since pullbacks of simplicial diagrams are computed objectwise, this means that the presentation of comp on simplicial presheaves factors through a dashed isomorphism of simplicial presheaves in the following diagram:

$$\begin{array}{ccc} \prod_{[g]} \left((X^g \times W(C_g \times \mathbb{R} \times \mathbb{Z}^{\times\bullet})) / (C_g \times \mathbb{R} \times \mathbb{Z}^{\times\bullet}) \right) & & \\ \downarrow \wr & \searrow \text{comp}_{X//G} & \\ \left([\mathbb{Z}^{\times\bullet}, X \times G^{\times\bullet}]_{\text{skel}} \times W(\mathbb{R} \times \mathbb{Z}^{\times\bullet}) \right) / (\mathbb{R} \times \mathbb{Z}^{\times\bullet}) & \longrightarrow & \left([\mathbb{Z}^{\times\bullet}, X \times G^{\times\bullet}]_{\text{skel}} \times W(\mathbb{Z}^{\times\bullet}) \right) / \mathbb{Z}^{\times\bullet} \\ \downarrow & \text{(pb)} & \downarrow \in \text{Fib} \\ \overline{W}(\mathbb{R} \times \mathbb{Z}^{\times\bullet}) & \xrightarrow{\overline{W}((r, \vec{n}) \mapsto \vec{n})} & \overline{W}(\mathbb{Z}^{\times\bullet}) \end{array} \quad (40)$$

Here:

- (i) the morphism on the right is a fibration, since $[\mathbb{Z}^{\times\bullet}, X \times G^{\times\bullet}]_{\text{skel}}$ is the nerve of a groupoid and hence a Kan complex (over each $U \in \text{CartSpc}$) and because $((-)\times W(\mathbb{Z}^{\times\bullet})) / (\mathbb{Z}^{\times\bullet})$ is a right Quillen functor (by Prop. A.34) and hence preserves Kan fibrations;
- (ii) the object in the bottom left represents the delooping $\mathbf{BS}^1_{\text{coh}}$, by Lemma 2.8;
- (iii) the morphism on the bottom represents the shape unit on $\mathbf{BS}^1_{\text{coh}}$ by the same argument as in Lemma 2.8.

This implies that the diagram exhibits the claimed homotopy pullback (39). \square

We conclude by proving the remaining Lemmas used in the above arguments.

Lemma 2.7 (Cofibrant resolution of circle group). *For $g \in G$, we have the following cofibrant replacements of the Lie group Λ_g (27) in $\Delta\text{PSh}(\text{CartSpc})_{\text{proj, loc}}$:*

$$\begin{array}{ccc} \emptyset \xrightarrow{\in \text{Cof}} \mathbb{R} \times \mathbb{Z}^{\times\bullet} & \xrightarrow{\sim} & \mathbb{R}^{\times S^1_{\text{coh}}} \xrightarrow{\in \text{W}} S^1_{\text{coh}}, \\ (r, \vec{n}) & \longmapsto & (r, (r+n_1), (r+n_1+n_2), \dots) \end{array} \quad (41)$$

$$\begin{array}{ccc} \emptyset \xrightarrow{\in \text{Cof}} C_g \times \mathbb{R} \times \mathbb{Z}^{\times\bullet} & \xrightarrow{\sim} & (C_g \times \mathbb{R})^{\times \Lambda_g} \xrightarrow{\in \text{W}} \Lambda_g. \\ (h, r, -\vec{n}) & \longmapsto & ((h, r), (g^{n_1} \cdot h, r+n_1), (g^{n_1+n_2} \cdot h, r+n_1+n_2), \dots) \end{array}$$

Proof. The first statement is Example A.56 and the second follows analogously. \square

Lemma 2.8 (Shape unit of the smooth circle). *A presentation for the shape unit of the smooth circle is given by*

$$\begin{array}{ccc} \text{smooth} & S^1_{\text{coh}} & \xrightarrow{\eta_{S^1_{\text{coh}}}} & \mathcal{J}S^1_{\text{coh}} \simeq \mathbf{B}\mathbb{Z} & \in \text{SmothGrpd}_\infty \\ \infty\text{-stacks} & & & & \uparrow \text{Loc}_W \\ \text{simplicial} & \mathbb{R} \times \mathbb{Z}^{\times\bullet} & \xrightarrow{(r, \vec{n}) \mapsto \vec{n}} & \mathbb{Z}^{\times\bullet} & \in \Delta\text{PSh}(\text{CartSpc})_{\text{proj, loc}} \\ \text{presheaves} & & & & \end{array} \quad (42)$$

Proof. By Lemma 2.7, the simplicial presheaf $\mathbb{R} \times \mathbb{Z}^{\times \bullet}$ is a cofibrant resolution of the circle, and by Prop. A.53 its image under shape is given by

$$\varinjlim (\mathbb{R} \times \mathbb{Z}^{\times \bullet}) \simeq \mathbb{Z}^{\times \bullet} \in (\Delta \text{Set}_{\text{Qu}})_{\text{fib}} \xrightarrow{\text{const}} (\Delta \text{Sh}(\text{CartSpc}))_{\text{loc}}^{\text{proj}}_{\text{fib}}.$$

This being fibrant means that the $(\varinjlim \dashv \text{const})$ -adjunction unit, which is as shown in (42), is already the derived unit. \square

Lemma 2.9 (Canonical $\int S_{\text{coh}}^1$ -action on inertia orbifold). *The canonical action (by Prop. A.41) of $\int S_{\text{coh}}^1 \simeq \mathbf{B}\mathbb{Z}$ on any inertia stack (25) – in particular on the inertia orbifold (24) of a good orbifold $\mathcal{X} \simeq X // G$ – is presented (under Prop. A.34) by the following simplicial action of $\mathbf{B}\mathbb{Z} \simeq \mathbb{Z}^{\times \bullet}$:*

$$\begin{aligned} \text{Map}(\mathbf{B}\mathbb{Z} \times \Delta[\bullet], \mathcal{X}) \times \mathbb{Z}^{\times \bullet} &\longrightarrow \text{Map}(\mathbf{B}\mathbb{Z} \times \Delta[\bullet], \mathcal{X}) \\ ((x, (h_k, \dots, h_2, h_1)), (n_k, \dots, n_2, n_1)) &\longmapsto (x, (g^{n_k} \cdot h_k, \dots, g^{n_2} \cdot h_2, g^{n_1} \cdot h_1)). \end{aligned} \quad (43)$$

Proof. We show the case $k = 1$; the general case is directly analogous. Here we unwind formula (102) as follows:

$$\begin{array}{ccccccc} \mathbf{B}\mathbb{Z} \times \Delta[1] & \xrightarrow{\text{id} \times \text{diag}} & \mathbf{B}\mathbb{Z} \times \Delta[1] \times \Delta[1] & \xrightarrow{\text{id} \times n \times \text{id}} & \mathbf{B}\mathbb{Z} \times \mathbf{B}\mathbb{Z} \times \Delta[1] & \xrightarrow{+\times \text{id}} & \mathbf{B}\mathbb{Z} \times \Delta[1] & \xrightarrow{1 \square \xrightarrow{x} g^h \square} & X // G \\ ((0, 1), [0, 1, 1]) & \mapsto & ((0, 1), [0, 1, 1][0, 1, 1]) & \mapsto & ((0, 1), (n, 0), [0, 1, 1]) & \mapsto & ((n, 1), [0, 1, 1]) & \xrightarrow{\quad} & \\ ((1, 0), [0, 0, 1]) & \mapsto & ((1, 0), [0, 0, 1][0, 0, 1]) & \mapsto & ((1, 0), (0, n), [0, 0, 1]) & \mapsto & ((1, n), [0, 0, 1]) & \xrightarrow{\quad} & x \square \xrightarrow{g^n} h \square \end{array}$$

$$\begin{array}{ccc} \begin{array}{ccc} (\bullet, [0]) & \xrightarrow{(0, [0, 1])} & (\bullet, [1]) \\ \downarrow & \searrow^{((0, 1), [0, 1, 1])} & \downarrow \\ (1, [0, 0]) & & (1, [1, 1]) \\ \downarrow & \searrow^{((1, 0), [0, 0, 1])} & \downarrow \\ (\bullet, [0]) & \xrightarrow{(0, [0, 1])} & (\bullet, [1]) \end{array} & \longmapsto & \begin{array}{ccc} (\bullet, [0]) & \xrightarrow{(n, [0, 1])} & (\bullet, [1]) \\ \downarrow & \searrow^{((n, 1), [0, 1, 1])} & \downarrow \\ (1, [0, 0]) & & (1, [1, 1]) \\ \downarrow & \searrow^{((1, n), [0, 0, 1])} & \downarrow \\ (\bullet, [0]) & \xrightarrow{(n, [0, 1])} & (\bullet, [1]) \end{array} \end{array} \quad \xrightarrow{\quad} \quad \begin{array}{ccc} x & \xrightarrow{g^n \cdot h} & h \cdot x \\ g \downarrow & & \downarrow g' \\ x & \xrightarrow{g^n \cdot h} & h \cdot x \end{array}$$

On the left above we show the non-degenerate 2-cell in the Cartesian product (Ex. A.5) whose image will pick up the generating data in the 1-cell of the hom-complex. Chasing this 2-cell through the formula (102) yields the result, as shown. \square

3 Transgression as cyclification

A simple but archetypical special case of orbifold cohomology is *group cohomology*, which may be understood as given by homotopy classes of maps of ∞ -groupoids of the form $\mathbf{B}G \rightarrow \mathbf{B}^n A$ (Ex. A.17). Here we discuss the technical detail of how cyclification of orbifolds leads to transgression in group cohomology, by proving the equivalence expressed by the top horizontal arrow in the diagram (3).

The following component Definition 3.1 is (up to an irrelevant global sign which we omit) due to [ARZ07, Def. 4.1], and maybe independently due to [Wi08, §1.3.3] where it is motivated by a geometric picture similar to that made precise by Thm. 3.4 below. Notice that it is tedious (albeit straightforward) to explicitly check that the component formula (44) really makes sense, in that it satisfies the cocycle condition; but this is implied by our more abstract characterization in Thm. 3.4 below.

Definition 3.1 (Transgression formula for discrete group cocycles). For $A \in \text{AbGrp}$, $n \in \mathbb{N}$ and $c : G^{\times n+1} \rightarrow A$ a group cocycle $[c] \in H_{\text{grp}}^{n+1}(G; A)$, obtain an n -cocycle on the inertia groupoid $\Delta \mathbf{B}G$ (Ntn. 3.3)

$$[\text{tr}(c)] \in H^n(\Delta \mathbf{B}G; A)$$

by setting:

$$\begin{aligned} \text{tr}(c) & \left(\gamma \xrightarrow{g_{n-1}} \text{Ad}_{n-1}(\gamma) \xrightarrow{g_{n-2}} \cdots \xrightarrow{g_1} \text{Ad}_1(\gamma) \xrightarrow{g_0} \text{Ad}_0(\gamma) \right) \\ & := \sum_{0 \leq j \leq n} (-1)^j \cdot c \left(g_{n-1}, \dots, g_{n-j}, \text{Ad}_j(\gamma), g_{n-j-1}, \dots, g_0 \right), \end{aligned} \quad (44)$$

where we abbreviate

$$\text{Ad}_j(\gamma) := \text{Ad}_{(g_{n-1} \cdots g_j)}(\gamma) := (g_{n-1} \cdots g_0)^{-1} \cdot \gamma \cdot (g_{n-1} \cdots g_0). \quad (45)$$

Remark 3.2 (Group cohomology). Since the inertia groupoid of the delooping of a finite group decomposes as a disjoint union of delooped centralizer groups C_g indexed over the conjugacy classes $[g] \in G/\text{ad}G$

$$\Lambda \mathbf{B}G \simeq \coprod_{[g] \in G/\text{ad}G} \mathbf{B}C_g \simeq \mathbf{B}G \sqcup \coprod_{[g] \neq [e]} \mathbf{B}C_g, \quad (46)$$

the formula (44) restricts to transgression maps to the group cohomology of all these groups, in particular to $C_e = G$ itself – as highlighted on the right of (46). Often only this leading component of the full transgression map is considered (e.g., implicitly so in [DW90, p. 14]).

We now prove (Thm. 3.4 below) that the transgression formula (44) abstractly arises from looping/cyclification according to (3). (Something similar is alluded to in [Wi08, §1.3.3].) First, we record the following Lemma 3.3 in simplicial homotopy theory, which is elementary but requires some care; in stating this we make heavy use of constructions and facts in simplicial homotopy theory that are recalled/introduced in appendix A.1:

Lemma 3.3 (Evaluation map on nerve of inertia groupoid). *On non-degenerate cells (cf. Prop. A.4), the evaluation map (61) on the inertia function complex (Ntn. A.26) is given by*

$$\begin{aligned} (S_{\min}^1)_{n+1} \times \text{Hom}(S_{\min}^1 \times \Delta[n+1], \overline{W}G) & \xrightarrow{\text{ev}} (\overline{W}G)_{n+1} \\ (s_{(0, \dots, \widehat{k}, \dots, n)}^\ell, s_k(\gamma; g_{n-1}, \dots, g_0)) & \longmapsto (g_{n-1}, \dots, g_{n-k}, \text{Ad}_k(\gamma), g_{n-(k+1)}, \dots, g_0). \end{aligned} \quad (47)$$

Here, on the left, $\ell \in (S_{\min}^1)$ denotes the non-degenerate cell in the minimal simplicial circle (Ntn. A.24) and $(\gamma; g_{n-1}, \dots, g_0)$ (Ntn. A.26) denotes a sequence of n composable morphisms in the inertia groupoid.

Proof. Using formula (70) for the components of the evaluation map, and then unwinding as in Ntn. A.26:

$$\begin{array}{c} (s_{(1, \dots, \widehat{k}, \dots, n+1)}^\ell, \mathbf{t}_{n+1}) \\ \cap \\ (S \times \Delta[n+1])_{n+1} \\ \downarrow \\ s_k(\gamma; g_{n-1}, \dots, g_0) = \\ (\gamma; g_{n-1}, \dots, g_{n-k}, e; g_{n-(k+1)}, \dots, g_0) \\ \downarrow \\ (\overline{W}G)_{n+1} \end{array} \quad \begin{array}{c} \begin{array}{ccccccc} (*, 0) & \xrightarrow{(\text{id}_*, [0,1])} & (*, 1) & \xrightarrow{(\text{id}_*, [1,2])} & \cdots & \xrightarrow{(\text{id}_*, [k-1,k])} & (*, k) & \xrightarrow{\text{id}_{(*,k)}} & (*, k) & \longrightarrow & \cdots & \xrightarrow{(\text{id}_*, [n-1,n])} & (*, n) \\ \downarrow & & \downarrow & & & & \downarrow & \searrow & \downarrow & & & & \downarrow \\ (\ell, \text{id}_0) & & (\text{id}_*, [0,1]) & & & & (\ell, \text{id}_k) & & (\ell, \text{id}_k) & & \cdots & & (\ell, \text{id}_n) \\ \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & & & \downarrow \\ (*, 0) & \xrightarrow{(\text{id}_*, [0,1])} & (*, 1) & \xrightarrow{(\text{id}_*, [1,2])} & \cdots & \longrightarrow & (*, k) & \xrightarrow{\text{id}_{(*,k)}} & (*, k) & \xrightarrow{\quad} & \cdots & \xrightarrow{(\text{id}_*, [n-1,n])} & (*, n) \end{array} \\ \downarrow \\ \begin{array}{ccccccc} \bullet & \xrightarrow{g_{n-1}} & \bullet & \xrightarrow{g_{n-2}} & \cdots & \xrightarrow{g_{n-k}} & \bullet & \xrightarrow{e} & \bullet & \xrightarrow{g_{n-(k+1)}} & \cdots & \xrightarrow{g_0} & \bullet \\ \downarrow & & \downarrow & & & & \downarrow & \searrow & \downarrow & & & & \downarrow \\ \bullet & \xrightarrow{g_{n-1}} & \bullet & \xrightarrow{g_{n-2}} & \cdots & \xrightarrow{g_{n-k}} & \bullet & \xrightarrow{e} & \bullet & \xrightarrow{g_{n-(k+1)}} & \cdots & \xrightarrow{g_0} & \bullet \\ & & \downarrow & & & & \downarrow & & \downarrow & & & & \downarrow \\ & & \text{Ad}_1(\gamma) & & & & \text{Ad}_k(\gamma) & & \text{Ad}_k(\gamma) & & \cdots & & \text{Ad}_n(\gamma) \end{array} \end{array}$$

Theorem 3.4 (Transgression in discrete group cohomology via looping). *The transgression formula of Def. 3.1 expresses equivalently the operation of applying the free homotopy loop functor $\text{Map}(\mathbf{B}\mathbb{Z}, -)$ to group cocycles understood as maps $\mathbf{B}G \rightarrow \mathbf{B}^{n+1}A$ (cf. Ex. A.17), composed with projection onto the resulting shifted coefficients (via Ex. A.40):*

$$\begin{array}{ccc}
H_{\text{grp}}^{n+1}(G; A) \simeq \pi_0 \text{Map}(\mathbf{B}G, \mathbf{B}^{n+1}A) & \xrightarrow{\text{Map}(\mathbf{B}\mathbb{Z}, -)} & \pi_0 \text{Map}(\Lambda \mathbf{B}G, \overbrace{\mathbf{B}^n A \times \mathbf{B}^{n+1}A}^{\text{by Ex. A.40}}) \\
& \searrow^{\text{transgression}} & \downarrow^{\text{pr}_1} \\
& & \pi_0 \text{Map}(\Lambda \mathbf{B}G, \mathbf{B}^n A) \simeq H^n(\Lambda \mathbf{B}G; A) \\
& & \downarrow^{\pi_0 \text{Map}(\mathbf{B}G \rightarrow \Lambda \mathbf{B}G, \mathbf{B}^n A)} \quad \downarrow \\
& & \pi_0 \text{Map}(\mathbf{B}G, \mathbf{B}^n A) \simeq H_{\text{grp}}^n(G; A).
\end{array}$$

Proof. In outline, this follows by regarding (62) the functor $\text{Map}(\mathbf{B}\mathbb{Z}, -)$ as pre-composition with the evaluation map and further precomposing this, under the free abelian group functor, with the Eilenberg-Zilber map (Prop. A.19): the component formula for the evaluation map, from Lemma 3.3, fed through the Eilenberg-MacLane formula (78) produces the signed summands that appear in (44).

In detail, we have the following sequence of natural transformations between homotopy classes⁶ of morphisms:

$$\begin{aligned}
H_{\text{grp}}^{n+1}(G; A) &= \pi_0 \text{Map}(\mathbf{B}G, \mathbf{B}^{n+1}A) && \text{Ex. A.17} \\
&\simeq \text{Ho}(\Delta \text{Set})\left(\overline{\mathbf{W}}G; \text{DK}(A[n+1])\right) && \text{Ex. A.22} \\
&\xrightarrow{\text{Map}(\mathbf{B}\mathbb{Z}, -)} \text{Ho}(\Delta \text{Set})\left(\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G), \text{Map}(S_{\min}^1, \text{DK}(A[n+1]))\right) && \text{\& Ex. A.16} \\
&\simeq \text{Ho}(\Delta \text{Set})\left(S_{\min}^1 \times \text{Map}(S_{\min}^1, \overline{\mathbf{W}}G), \text{DK}(A[n+1])\right) && \text{Ex. A.25} \\
&\simeq \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet} \circ \mathbb{Z}[S_{\min}^1 \times \text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)], A[n+1]\right) && (59) \\
&\simeq \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet}\left(\mathbb{Z}[S_{\min}^1] \otimes \mathbb{Z}[\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)]\right), A[n+1]\right) && (76) \\
&\xrightarrow[\nabla^*]{\simeq} \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet}\left(\mathbb{Z}[S_{\min}^1]\right) \otimes N_{\bullet}\left(\mathbb{Z}[\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)]\right), A[n+1]\right) && (77) \\
&\simeq \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet}\left(\mathbb{Z}[\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)]\right), \text{Map}\left(N_{\bullet}\left(\mathbb{Z}[S_{\min}^1]\right), A[n+1]\right)\right) && (78) \\
&\simeq \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet}\left(\mathbb{Z}[\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)]\right), A[n] \oplus A[n+1]\right) && (79) \\
&\xrightarrow{\text{pr}_1} \text{Ho}(\text{Ch}^{\geq 0}(\text{AbGrp}))\left(N_{\bullet}\left(\mathbb{Z}[\text{Map}(S_{\min}^1, \overline{\mathbf{W}}G)]\right), A[n]\right) \\
&= \pi_0 \text{Map}(\Lambda \mathbf{B}G, \mathbf{B}^n A) = H^m(\Lambda \mathbf{B}G; A) && (76).
\end{aligned}$$

Noticing here (see (62)) that the composite of the application of the functor $\text{Map}(\mathbf{B}\mathbb{Z}, -)$ with the adjointness relation is equivalent to pre-composition with the evaluation map, the net effect of this composite of transformations in degree $n+1$ is the pre-composition of the cocycle map in that degree, regarded as a homomorphism of abelian groups

$$\tilde{c}_{n+1} : \mathbb{Z}[G^{n+1}] \simeq \mathbb{Z}[(\overline{\mathbf{W}}G)_{n+1}] \longrightarrow A,$$

with the composite of the Eilenberg-Zilber- and the evaluation map:

$$\begin{aligned}
\mathbb{Z}[(S_{\min}^1)_1] \otimes \mathbb{Z}[\text{Hom}(S_{\min}^1 \times \Delta[n], \overline{\mathbf{W}}G)] &\xrightarrow{\nabla} \mathbb{Z}[(S_{\min}^1)_{n+1} \times \text{Hom}(S_{\min}^1 \times \Delta[n+1], \overline{\mathbf{W}}G)] \xrightarrow{\mathbb{Z}[\text{ev}]} \mathbb{Z}[(\overline{\mathbf{W}}G)_{n+1}] \\
\ell \otimes (\gamma, g_{n-1}, \dots, g_0) &\xrightarrow{(78)} \sum_{0 \leq j \leq n} (-1)^j \cdot (s_{(0, \dots, \widehat{k}, \dots, n)} \ell, s_k(\gamma, g_{n-1}, \dots, g_0)) \xrightarrow{\text{Lem. 3.3}} \sum_{0 \leq j \leq n} (-1)^j \cdot (g_{n-1}, \dots, g_{n-k}, \text{Ad}_k(\gamma), g_{n-(k+1)}, \dots, g_0)
\end{aligned} \tag{48}$$

followed by restriction to the coefficient of ℓ (86) on the left. This manifestly yields the formula (44). \square

⁶The collection of facts from model category theory needed for the routine verification that the underlying naive sequence of natural transformations passes to homotopy classes in each step (on p. 20) may be found reviewed [FSS20Cha, §A].

4 Integral 4-classes of equivariant 4-Cohomotopy

Here we prove that the integral 4-class $\tilde{\Gamma}_4$ which underlies any tangentially twisted 4-Cohomotopy cocycle *at an ADE-singularity* is that which classifies the ‘‘Platonic’’ 2-group extensions of [EG17]. These 4-classes (equivalently in their incarnation as 3-classes after cyclification) are the twists of quasi-elliptic cohomology which are ‘‘predicted’’ by *Hypothesis H* in the sense explained at the end of §1.

Notation 4.1 (Canonical representation 4-sphere of finite subgroup of $\mathrm{Sp}(1)$). Given a finite subgroup $G \xrightarrow{i} \mathrm{Sp}(1)$ (Prop. A.42), we write

$$G \wr \mathbb{H} := \mathrm{res}_i(\mathrm{Sp}(1) \wr \mathbb{H}) \in G \mathrm{Act}(\mathrm{VectorSpaces}_{\mathbb{R}})$$

for the linear representation which is the restriction along i of the defining linear action of $\mathrm{Sp}(1)$ on the real vector space $\mathbb{H} \simeq_{\mathbb{R}} \mathbb{R}^4$ underlying the algebra of quaternions. Accordingly, we write

$$G \wr \int S^{\mathbb{H}} \in G \mathrm{Act}(\mathrm{TopSpc}) \xrightarrow{\mathrm{Sing}} G \mathrm{Act}(\Delta \mathrm{Set})$$

for the shape of the corresponding representation sphere.

Theorem 4.2 (Equivariant integral characteristic classes of ADE-equivariant 4-Cohomotopy). *For $G \xrightarrow{i} \mathrm{Sp}(1) \simeq \mathrm{SU}(2) \simeq \mathrm{Spin}(3)$ a finite subgroup (Prop. A.42), the group of equivariant integral characteristic 4-classes of ADE-equivariant 4-Cohomotopy is*

$$H_G^4(S^{\mathbb{H}}, \mathbb{Z}) \simeq \mathbb{Z} \oplus \mathbb{Z}_{|G|}.$$

Proof. By Lemma 4.3 with Lemma 4.4. □

The following Lemma 4.3 may be taken to be the definition of ordinary equivariant cohomology, or else of a standard fact of Bredon cohomology with invariant coefficients. We include the following proof just to showcase how this fits into the singular-cohesive formalism of [SS20Orb][SS21EPB]:

Lemma 4.3 (Equivariant integral characteristic classes of equivariant Cohomotopy). *The equivariant integral characteristic classes of equivariant Cohomotopy are naturally identified with the ordinary cohomology of the homotopy quotient of the sphere.*

Proof. We have the following sequence of natural equivalences of hom- ∞ -groupoids:

$$\begin{aligned} \left\{ \begin{array}{ccc} \int \gamma(S^V // G) & \dashrightarrow & \gamma(\mathbf{B}^4 \mathbb{Z} // G) \\ & \searrow & \swarrow \\ & \gamma \mathbf{B}G & \end{array} \right\} &\simeq \left\{ \begin{array}{ccc} \int \gamma(S^V // G) & \dashrightarrow & \gamma(\mathbf{B}^4 \mathbb{Z} \times \mathbf{B}G) \\ & \searrow & \swarrow \\ & \gamma \mathbf{B}G & \end{array} \right\} & \text{trivial action on coefficients} \\ &\simeq \left\{ \begin{array}{ccc} \int \gamma(S^V // G) & \dashrightarrow & (\gamma \mathbf{B}^4 \mathbb{Z}) \times (\gamma \mathbf{B}G) \\ & \searrow & \swarrow \\ & \gamma \mathbf{B}G & \end{array} \right\} & \text{since } \gamma \text{ is right adjoint} \\ &\simeq \left\{ \int \gamma(S^V // G) \dashrightarrow \mathbf{B}^4 \mathbb{Z} \right\} & \text{by base change (106)} \\ &\simeq \left\{ \cup \int \gamma(S^V // G) \dashrightarrow \mathbf{B}^4 \mathbb{Z} \right\} & \text{by singular cohesion } \cup \dashv \gamma \\ &\simeq \left\{ \int \cup \gamma(S^V // G) \dashrightarrow \mathbf{B}^4 \mathbb{Z} \right\} & \text{by [SS20Orb, Lem. 3.67]} \\ &\simeq \left\{ \int \cup (S^H // G) \dashrightarrow \mathbf{B}^4 \mathbb{Z} \right\} & \text{by singular cohesion} \\ &\simeq \left\{ \int (S^V // G) \dashrightarrow \mathbf{B}^4 \mathbb{Z} \right\} & \text{since } S^V // G \text{ is smooth.} \quad \square \end{aligned}$$

Lemma 4.4 (Integral 4-cohomology of homotopy quotient of 4-sphere by ADE-group). *Let $G \xrightarrow{i} \mathrm{Sp}(1)$ be a finite subgroup (Prop. A.42). Then the integral 4-cohomology of the homotopy quotient of its canonical representation sphere (Nota. 4.1) is*

$$H^4((\mathbb{J}S^{\mathbb{H}}) // G, \mathbb{Z}) \simeq \mathbb{Z} \oplus (\mathbb{Z}/|G|), \quad (49)$$

where the first summand is the cohomology of the fiber, and the second is the group cohomology of G , in that we have a split short exact sequence in cohomology

$$\begin{array}{ccccccc} 0 & \longleftarrow & \overbrace{H^4(S^4; \mathbb{Z})}^{\mathbb{Z}} & \xleftarrow{q^*} & H^4(\mathbb{J}S^{\mathbb{H}} // G; \mathbb{Z}) & \xleftarrow{p^*} & \overbrace{H^4(BG; \mathbb{Z})}^{\mathbb{Z}/|G|} \longleftarrow 0 \\ & & \mathbb{J}S^4 & \xrightarrow{q} & \mathbb{J}(S^{\mathbb{H}} // G) & \xrightarrow{\rho} & BG \end{array} \quad (50)$$

induced by the homotopy fiber sequence of the Borel construction (Lem. 4.3), shown at the bottom in (50).

Proof. Observe that the fundamental group $\pi_1(BG) \simeq G$ of the base space of the fibration (50) acts trivially on the cohomology

$$H^n(\mathbb{J}S^4, \mathbb{Z}) \simeq \begin{cases} \mathbb{Z} & \text{for } n \in \{0, 4\} \\ 0 & \text{otherwise} \end{cases}$$

of the fiber space. This is because:

- we have a G -equivariant isomorphism between the representation sphere of \mathbb{H} and the unit sphere in $\mathbb{R} \oplus \mathbb{H}$ (by stereographic projection, e.g. [MP04, p. 2]): $S^{\mathbb{H}} \simeq_G S(\mathbb{R} \oplus \mathbb{H})$;
- the action of $\mathrm{Sp}(1)$ on $\mathbb{H} \simeq_{\mathbb{R}} \mathbb{R}^4$ is through the defining action of $\mathrm{SO}(4)$ (since the quaternions are a normed division algebra, so that left multiplication by unit-norm quaternions $q \in \mathrm{Sp}(1) = S(\mathbb{H})$ preserves the norm), whence the action of $\mathrm{Sp}(1)$ on $\mathbb{R} \oplus \mathbb{H} \simeq_{\mathbb{R}} \mathbb{R}^5$ is through $\mathrm{SO}(5)$ (e.g. [HSS18, Rem. A.8]);
- by the Hopf degree theorem and the de Rham theorem, the generator of $\mathbb{Z} \simeq H^4(S^4, \mathbb{Z})$ is represented by the standard volume form on S^4 , which is evidently preserved by the $\mathrm{SO}(5)$ -action (e.g. [BSS18, p. 31]).

Therefore (e.g. [HatSS, Thm. 1.14]), we have a cohomological Serre spectral sequence of the form

$$E_2^{p,q} = H^p(BG; H^q(S^4, \mathbb{Z})) \Rightarrow H^{p+q}(S^{\mathbb{H}} // G; \mathbb{Z}). \quad (51)$$

From Prop. A.43 with Ex. A.17, it follows that its second page looks as follows:

$$E_2^{\bullet, \bullet} = \begin{array}{cccccccc} & & S^4 & & & & & & \\ & & \vdots & & \vdots & & \vdots & & \vdots \\ & & 0 & & 0 & & 0 & & 0 & \dots \\ & & \mathbb{Z} & & 0 & & G^{\mathrm{ab}} & & 0 & \dots \\ & & 0 & & 0 & & 0 & & 0 & \dots \\ & & 0 & & 0 & & 0 & & 0 & \dots \\ & & 0 & & 0 & & 0 & & 0 & \dots \\ & & \mathbb{Z} & & 0 & & G^{\mathrm{ab}} & & 0 & \dots \\ & & & & & & & & & BG \end{array}$$

This shows that every differential on this and on every following page has zero domain or zero codomain, so that the spectral sequence collapses already on this page: $E_{\infty}^{\bullet, \bullet} \simeq E_2^{\bullet, \bullet}$. By its convergence (51), this means that we have a short exact sequence of the form (50). Since $\mathrm{Ext}(\mathbb{Z}, -) = 0$, the claim (49) follows. \square

Proposition 4.5 (Shifted integral characteristic 4-class for equivariant 4-Cohomology ([FSS19HypH, §3.4])). *There exists an integral class*

$$\tilde{\Gamma}_4 := \frac{1}{2}\chi_4 + \frac{1}{4}p_1 \in H^4(\mathbb{J}S^{\mathbb{H}} // \mathrm{Sp}(2); \mathbb{Z}) \quad (52)$$

whose rational image is the sum of half the Euler class with 1/4th of the Pontrjagin class, hence whose restriction to the 4-sphere fiber is the volume class

$$\begin{array}{ccc} H^4(\mathbb{J}S^4; \mathbb{Z}) & \ni & [\text{vol}_{S^4}] \\ \uparrow & & \uparrow \\ H^4(\mathbb{J}(S^{\mathbb{H}} // \text{Sp}(2)); \mathbb{Z}) & \ni & \tilde{\Gamma}_4 \end{array} \quad (53)$$

Proposition 4.6 (Universal shifted integral 4-flux restricts to generator on ADE-singularity). *For a finite subgroup of $\text{Sp}(1)$ (Prop. A.42), embedded as the left diagonal entry into $\text{Sp}(2)$*

$$\begin{array}{ccccccc} & & & & j & & \\ & & & & \downarrow & & \\ G & \xleftarrow{i} & \text{Sp}(1) & \xleftarrow{l} & \text{Sp}(1) \times \text{Sp}(1) & \xrightarrow{\quad} & \text{Sp}(2) \\ q & \mapsto & q & \mapsto & (q, 1) & \mapsto & \begin{pmatrix} q & 0 \\ 0 & 1 \end{pmatrix} \end{array} \quad (54)$$

the pullback of $\tilde{\Gamma}_4$ (52) along $\mathbb{J}(S^{\mathbb{H}} // j)$ is the element $(1, [1])$ according to Lem. 4.4:

$$\begin{array}{ccc} \mathbb{Z} \oplus \mathbb{Z}_{|G|} \simeq H^4(\mathbb{J}S^{\mathbb{H}} // G; \mathbb{Z}) & \xleftarrow{Bj^*} & H^4(\mathbb{J}(S^{\mathbb{H}} // \text{Sp}(2)); \mathbb{Z}) \\ (1, [1]) & \xleftarrow{\quad} & \tilde{\Gamma}_4 \end{array} \quad (55)$$

Proof. Consider the corresponding morphism of Borel constructions (96) and observe the following pullbacks of cohomology generators through this diagram:

$$\begin{array}{ccccc} (1, [0]) & \xleftarrow{\text{Lem. 4.4}} & & & [\text{vol}_{S^4}] \\ & & S^4 & \xlongequal{\quad} & S^4 \\ & & \downarrow & & \downarrow \\ (0, [1]) & \mathbb{J}(S^{\mathbb{H}} // G) & \xrightarrow{\mathbb{J}(S^{\mathbb{H}} // j)} & \mathbb{J}(S^{\mathbb{H}} // \text{Sp}(1)) & \tilde{\Gamma}_4 \\ \uparrow \text{Lem. 4.4} & \downarrow & & \downarrow & \uparrow (53) \\ [1] & BG & \xrightarrow{Bi} & B\text{Sp}(1) & \\ & \xleftarrow{\text{Prop. A.44}} & & & \frac{1}{4}p_1 \end{array} \quad (56) \quad \square$$

Remark 4.7 (Background integral 4-flux at flat ADE-singularity). Noticing that $G \curvearrowright S^{\mathbb{H}}$ has a fixed point (in fact two), hence that $S^{\mathbb{H}} // G \xrightarrow{\rho} BG$ has a section, this means that in the vicinity of any G -orbifold-singularity $\mathcal{X} \simeq \mathbb{R}^n // G$, hence with $\mathbb{J}\mathcal{X} \simeq * // G$, there is integral *background* C-flux of value $[1] \in H^4(BG; \mathbb{Z})$

$$\begin{array}{ccccc} & & (0, [1]) & & \\ & & \downarrow & & \\ * // G & \xrightarrow{\quad} & S^{\mathbb{H}} // \text{Sp}(2) & \xrightarrow{\tilde{\Gamma}_4} & \mathbf{B}^4\mathbb{Z} \\ \downarrow \wr & \text{unique cocycle in} & \downarrow \rho & & \\ \mathbf{B}G & \xrightarrow{\quad} & \mathbf{B}\text{Sp}(2) & & \end{array} \quad (57)$$

equivariant 4-Cohomotopy

This is hence the background value of “M-theoretic discrete torsion”, in the terminology of [Sh03][Se01], see also [dB+02, §4.6]. The 2-gerbe over $* // G$ classified by this background 4-flux is (the delooping of) the universal *Platonic 2-group-extension* of G , in the terminology of [EG17]. However, the proper definition of brane charge *localized* at a singularity (i.e. disregarding charges that are “escaping to infinity”) is given by the cohomology/cohomotopy of the *one-point compactification* of the transverse space $\mathbb{R}^n // G$ to the brane [SS20Tad, (8)][SS22Conf, §2.1][SS21MF, (12)]. For flat M5-branes at an ADE-singularity this is again the 4-sphere orbifold $S^{\mathbb{H}} // G$ [SS20Tad], and the unit integral brane charge of a single one of these is (55):

$$\begin{array}{ccccc} & & (1, [1]) & & \\ & & \downarrow & & \\ S^{\mathbb{H}} // G & \xrightarrow{\quad} & S^{\mathbb{H}} // \text{Sp}(2) & \xrightarrow{\tilde{\Gamma}_4} & \mathbf{B}^4\mathbb{Z} \\ \downarrow & \text{unit cocycle in} & \downarrow \rho & & \\ \mathbf{B}G & \xrightarrow{\quad} & \mathbf{B}\text{Sp}(2) & & \end{array} \quad (58)$$

equivariant 4-Cohomotopy

Remark 4.8 (Conclusion). With this we have finished compiling the ingredients for the construction/application indicated at the end of §1: The unit integral background 4-flux in M-theory localized at an ADE-singularity, as predicted by *Hypothesis H*, is (58); and its double dimensional reduction (16) to 3-flux according to (16) is, by (11), the corresponding transgression class. This is the twist for ADE-equivariant quasi-elliptic cohomology to be used in (20) for measuring M-brane charge in equivariant quasi-elliptic cohomology and thus potentially relating to the elliptic genus of the M5-brane.

A Appendix: Technical background

For use in the main text, here we recall and reference some technical background:

- on basic facts of simplicial homotopy theory (appendix A.1)
- on basic notions in cohesive ∞ -topos theory (appendix A.2).

A.1 Some simplicial homotopy theory

After recalling the combinatorics of products of simplicial sets in streamlined form, here we compile some technical background on simplicial groups and their actions and prove some useful facts that do not seem to be easily citable from the literature. For more along these lines see [SS21EPB, §3.1.2], whose notation we follow. For general background on simplicial homotopy theory see [GJ99][Re22, §1].

Categories and simplicial sets. Just to set up our notation:

Notation A.1 (Mapping objects).

- Generally, $\text{Hom}(-, -)$ is to denote hom-sets in a given category, while $\text{Map}(-, -)$ denotes internal hom-objects (for cartesian closed categories), defined to yield natural bijections:

$$\text{Hom}(X \times Y, Z) \xleftarrow[\sim]{(-)} \text{Hom}(X, \text{Map}(Y, Z)). \quad (59)$$

- In particular, for a pair of (small) categories⁷ $\mathcal{X}, \mathcal{Y} \in \text{Cat}$, the notation $\text{Hom}(\mathcal{C}, \mathcal{D})$ denotes the plain set of functors between them, while $\text{Map}(\mathcal{X}, \mathcal{Y})$ denotes the category of such functors with natural transformations between them, schematically:

$$\text{Hom}(\mathcal{X}, \mathcal{Y}) = \left\{ \mathcal{X} \xrightarrow{F} \mathcal{Y} \right\}, \quad \text{Map}(\mathcal{X}, \mathcal{Y}) = \left\{ \mathcal{X} \begin{array}{c} \xrightarrow{F} \\ \Downarrow \phi \\ \xrightarrow{F'} \end{array} \mathcal{Y} \right\}. \quad (60)$$

- We write

$$\text{ev}_X^A := \text{id}_{\widetilde{\text{Map}}(X, A)} : X \times \text{Map}(X, A) \longrightarrow A \quad (61)$$

for the *evaluation map* on these mapping objects, i.e. the adjunct (59) of the identity on $\text{Map}(X, A)$.

Notice that the naturality of (59) implies that adjunct of the application of the mapping object functor to a morphism equals the pre-composition of that morphism with the evaluation map:

$$f : A \rightarrow B \quad \vdash \quad \widetilde{\text{Map}}(X, f) = f \circ \text{ev}_X^A : X \times \text{Map}(X, A) \longrightarrow B. \quad (62)$$

Notation A.2 (Notation for simplicial sets). We write, essentially as usual:

- $[n] := \{0 \rightarrow 1 \rightarrow \dots \rightarrow n\}$ for the category free on a sequence of n composable morphisms, for $n \in \mathbb{N}$; so that functors $[n] \rightarrow \mathcal{C}$ may be identified with paths of n composable morphisms in \mathcal{C} ;

- $\bullet d_n^i : [n] \rightarrow [n+1] \quad 0 \Rightarrow 1 \Rightarrow \dots \Rightarrow i-1 \xrightarrow{\quad} i \xrightarrow{\quad} i+1 \Rightarrow \dots \Rightarrow n \Rightarrow n+1,$

⁷To be pedantic, in the present context by a “category” we mean, as is usual, a *strict* category, i.e. with a fixed set of objects (which is more information than retained in the equivalence class). It is (only) on these strict categories that the simplicial nerve (Ntn. A.2) is defined.

- $s_{n+1}^i : [n+1] \rightarrow [n]$ $0 \rightrightarrows 1 \rightrightarrows \dots \rightrightarrows i$
 \parallel
 $i \rightrightarrows \dots \rightrightarrows n-1 \rightrightarrows n$,
- $l_n : [n] \xrightarrow{\sim} [n]$ $0 \rightrightarrows 1 \rightrightarrows \dots \rightrightarrows n-1 \rightrightarrows n$,

for the *co-face*-, the *co-degeneracy*-, and the *identity*-functors between these categories, shown on the right as paths of composable edges in the respective codomain (as such used around Lem. 3.3, cf. *Figure NDS*);

- $\Delta \hookrightarrow \text{Cat}$ for the full subcategory of the 1-category of (strict) categories on those of the form $[n]$, $n \in \mathbb{N}$;
- $\Delta\text{Set} = \text{PSh}(\Delta) = \text{Hom}(\Delta^{\text{op}}, \text{Set})$ for the category of simplicial sets;
- $N : \text{Cat}^{\text{small}} \xrightarrow{\mathcal{C} \mapsto ([n] \mapsto \text{Hom}([n], \mathcal{C}))} \Delta\text{Set}$ for the simplicial nerve functor;
- $\Delta[n] := N[n] \in \Delta\text{Set}$ for the standard simplicial simplices.

Example A.3 (Product categories). For \mathcal{C}, \mathcal{D} a pair of small categories, their *product category* $\mathcal{C} \times \mathcal{D}$ has morphisms forming commuting squares as follows, for f any morphism in \mathcal{C} and g any morphism in \mathcal{D} :

$$\begin{array}{ccc}
 (c_1, d_1) & \xrightarrow{(f, \text{id}_{d_1})} & (c_2, d_1) \\
 (\text{id}_{c_1}, g) \downarrow & \searrow (f, g) & \downarrow (\text{id}_{c_2}, g) \\
 (c_1, d_2) & \xrightarrow{(f, \text{id}_{d_2})} & (c_2, d_2)
 \end{array}$$

This implies that natural transformations between functors $\mathcal{C} \rightarrow \mathcal{D}$ are in bijection to functors $\mathcal{C} \times [1] \rightarrow \mathcal{D}$, hence that the simplicial nerve of functor categories (60) is given by:

$$N\text{Map}(\mathcal{C}, \mathcal{D}) : [n] \longmapsto \text{Hom}(\mathcal{C} \times [n], \mathcal{D}). \quad (63)$$

Products and mapping complexes of simplicial sets. Famously, the cartesian product of simplicial sets, despite its evident component-wise construction,

$$X, Y \in \Delta\text{Set} \vdash \begin{cases} X \times Y \in \Delta\text{Set} \\ (X \times Y)_n = X_n \times Y_n \\ s_k^{X \times Y} = (s_k^X, s_k^Y), \quad d_k^{X \times Y} = (d_k^X, d_k^Y), \end{cases} \quad (64)$$

has a remarkably rich combinatorics, which we briefly recall as Prop. A.4. The phenomena dually induced on the mapping complexes (these we recall as Prop. A.6 below and use as Lemma 3.3 in the main text) may not have received comparable attention yet; in any case this is what drives the proof of Thm. 3.4 in the main text.

The content of the following Prop. A.4 is classical (it is at least implicit in the ‘‘Eilenberg-MacLane formula’’ [EML53, (5.3)], as reviewed for instance in [Fr20, §5.2.1 & §B.6], with exposition in [Fr12, §5]) but we will have need for the following concise formulation (cf. also [Kerodon, Ntn. 2.5.7.2]):

Proposition A.4 (Non-degenerate simplices in product simplicial sets). *For $p, q \in \mathbb{N}$, the non-degenerate $p+q$ -simplices in the Cartesian product of the standard p -simplex with the standard q -simplex,*

$$\Delta[p+q] \xrightarrow{(\sigma^{\text{hor}}, \sigma^{\text{ver}})} \Delta[p] \times \Delta[q]$$

are precisely (cf. *Figure NDS*):

- as pairs of paths of edges $(\sigma^{\text{hor}}[k, k+1], \sigma^{\text{ver}}[k, k+1])$, those for which at each step precisely one of the two edges in the pair is degenerate and the other a generating edge;

– equivalently, the sequences of step numbers at which one or the other is non-trivial, these being the “ (p, q) -shuffles” (μ, ν) of elements $(0, 1, 2, \dots, p + q - 1)$:

$$\begin{array}{ccc} & \text{steps } k \text{ at which} & \\ \sigma^{\text{hor}} \text{ is non-trivial:} & & \sigma^{\text{ver}} \text{ is non-trivial:} \\ \mu_0 < \dots < \mu_{p-1} & & \nu_0 < \dots < \nu_{q-1}, \end{array} \quad (65)$$

– from which the given non-degenerate cell $(\sigma^{\text{hor}}, \sigma^{\text{ver}})$ is recovered as:

$$\begin{aligned} \sigma^{\text{hor}} = s_{p+q}^{\nu} & := [p+q] \xrightarrow{s_{p+q}^{\nu_0}} [p+q-1] \xrightarrow{s_{p+q-1}^{\nu_1}} [p+q-2] \rightarrow \dots \rightarrow [p+1] \xrightarrow{s_{p+1}^{\nu_{q-1}}} [p] \\ \sigma^{\text{ver}} = s^{\mu} & := [p+q] \xrightarrow{s_{p+q}^{\mu_0}} [p-1+q] \xrightarrow{s_{p-1+q}^{\mu_1}} [p-2+q] \rightarrow \dots \rightarrow [1+q] \xrightarrow{s_{1+q}^{\mu_{p-1}}} [q]. \end{aligned} \quad (66)$$

For $X, Y \in \Delta\text{Set}$, the non-degenerate $p+q$ -simplices in $X \times Y$ (64) are of the form


$$\left(s_{\nu}^X(x_p), s_{\mu}^Y(y_q) \right) : \Delta[p+q] \xrightarrow{\text{diag}} \begin{array}{ccc} \Delta[p+q] & \xrightarrow{s_{p+q}^{\nu}} & \Delta[p] \\ \times & & \times \\ \Delta[p+q] & \xrightarrow{s_{p+q}^{\mu}} & \Delta[q] \end{array} \xrightarrow{\begin{array}{c} x_p \\ y_q \end{array}} \begin{array}{c} X \\ \times \\ Y \end{array} \quad (67)$$

for $x_p \in X_p$ and $y_q \in Y_q$ non-degenerate cells and (μ, ν) a (p, q) -shuffle.


Proof. Generally, since the degeneracy maps in a product $X \times Y$ of simplicial sets are the pairs of the separate degeneracy maps, $s_k^{X \times Y} = (s_k^X, s_k^Y)$ (64), a cell in the product is non-degenerate precisely if its two components are not both in the image of some s_k , for the same k . This implies generally that non-degenerate paths in products are those that do not have steps where both components are degenerate. On the other hand, since the particular paths in question have length $p+q$, a moment of reflection shows that if they had a step with both components non-trivial, then the previous constraint could not be satisfied in all other steps. This implies the first claim above. From this, the next two formulations follow by inspection. \square

Figure NDS. The non-degenerate simplices in the product $\Delta[p] \times \Delta[q]$ (seen as in Ex. A.3) are (according to Prop. A.4) given by those paths of $(p+q)$ steps for which each step is a unit step either horizontally or vertically. The lists of steps (counted starting at 0) going horizontally, $\mu = (\mu_0 < \mu_1 < \dots, \mu_{p-1})$, or going vertically, $\nu = (\nu_0 < \nu_1 < \dots, \nu_{p-1})$ form jointly a (p, q) -*(un-)shuffle* permutation of $(0, 1, \dots, p+q)$, which bijectively encodes the respective non-degenerate cell according to the formula (66).

(Beware that the dashed diagonal arrow shown on the right is a stand-in for any remaining *zig-zag* path and not for any actual diagonal steps.)



Example A.5. The complete set of non-degenerate cells in the simplicial square $\Delta[1] \times \Delta[1]$ is the following, with the paths (according to *Figure NDS*) highlighted which correspond to the two non-degenerate 2-simplices:



The following makes explicit the solution to the universal property (59) of mapping objects in simplicial sets.

Proposition A.6 (Simplicial mapping complexes, e.g. [GJ99, §I.5]). For $X, A \in \Delta\text{Set}$:

(i) their mapping complex $\text{Map}(X, A) \in \Delta\text{Set}$ is

$$\text{Map}(X, A) : [n] \longmapsto \text{Hom}(X \times \Delta[n], A) \quad (68)$$

in that with this formula we have, for $S \in \Delta\text{Set}$, the required natural isomorphisms (cf. Ntn. A.1)

$$\text{Hom}(S \times X, A) \simeq \text{Hom}(S, \text{Map}(X, A)); \quad (69)$$

(ii) the corresponding evaluation map (61) is given by naive evaluation on given cells σ_n but paired with the identity n -cell ι_n :

$$\begin{array}{ccc} (S \times \text{Map}(S, A))_n & \underset{(68)}{\simeq} & S_n \times \text{Hom}(S \times \Delta[n], A) & \xrightarrow{(\text{ev}_S^A)_n} & A_n \\ & & \underset{(\sigma_n, f)}{\longmapsto} & & \underset{f(\sigma_n, \iota_n)}{\longrightarrow} \end{array} \quad (70)$$

Remark A.7 (Self-enrichment of simplicial sets). By (69), the construction of simplicial mapping complexes is functorial in both arguments:

$$\text{Map}(-, -) : \Delta\text{Set}^{\text{op}} \times \Delta\text{Set} \longrightarrow \Delta\text{Set}$$

and has a natural, associative and unital composition operation

$$\text{Map}(X, Y) \times \text{Map}(Y, Z) \xrightarrow{\circ} \text{Map}(X, Z),$$

which on 0-cells reduces to the ordinary composition of morphisms. This gives ΔSet the structure of an ΔSet -enriched category (e.g. [Ke82][Hir02, §9.1.1]).

Proposition A.8 (Properties of the simplicial nerve). *The simplicial nerve $N : \text{Cat}^{\text{small}} \rightarrow \Delta\text{Set}$ (Ntn. A.2)*

(i) is fully faithful:

$$\text{Hom}(\mathcal{C}, \mathcal{D}) \xrightarrow[\sim]{N_{\mathcal{C}, \mathcal{D}}} \text{Hom}(N\mathcal{C}, N\mathcal{D}); \quad (71)$$

(ii) sends product categories (Ex. A.3) to products (64) of their nerves

$$N(\mathcal{C} \times \mathcal{D}) \simeq (N\mathcal{C}) \times (N\mathcal{D}); \quad (72)$$

(iii) sends functor categories (60) to simplicial function complexes (68):

$$N\text{Map}(\mathcal{C}, \mathcal{D}) \simeq \text{Map}(N\mathcal{C}, N\mathcal{D}) \quad (73)$$

(all these being natural isomorphisms).

Proof. The first statement is classical, the (elementary) proof is spelled out e.g. in [Re22, Prop. 4.10] [Kerodon, Prop. 1.2.2.1]. The second statement is immediate from the two definitions, as both are given by pairs of all structure morphisms. From this the third statement is obtained as the following sequence of natural isomorphisms, for $n \in \mathbb{N}$:

$$\begin{aligned} (N\text{Map}(\mathcal{C}, \mathcal{D}))_n &\simeq \text{Hom}(\mathcal{C} \times [n], \mathcal{D}) && \text{by (63)} \\ &\simeq \text{Hom}(N(\mathcal{C} \times [n]), N\mathcal{D}) && \text{by (71)} \\ &\simeq \text{Hom}(N\mathcal{C} \times \Delta[n], N\mathcal{D}) && \text{by (72)} \\ &\simeq (\text{Hom}(N\mathcal{C}, N\mathcal{D}))_n && \text{by (68)}. \end{aligned} \quad \square$$

Homotopy theory of simplicial Sets. We make use of basic (simplicial) model category theory. A standard account is [Hir02], a concise overview of all the ingredients that we need is in [FSS20Cha, §A].

Notation A.9 (Homotopy theory of simplicial sets, e.g. [GJ99, §I.11]). We write

$$\Delta\text{Set}_{\text{Qu}} \in \text{MdlCat} \quad (74)$$

for the classical Kan-Quillen model category structure on simplicial sets.

Notation A.10 (Homotopy theory of reduced simplicial sets, e.g. [GJ99, §V Prop. 6.12]). We write

$$\Delta\text{Set}_{\geq 1, \text{inj}} \in \text{MdlCat} \quad (75)$$

for the model category of reduced simplicial sets (those $S \in \Delta\text{Set}$ with a single vertex, $S_0 = *$) whose weak equivalences and cofibrations are those of the underlying $\Delta\text{Set}_{\text{Qu}}$ (74).

Lemma A.11 (Fibrant reduced simplicial sets are Kan complexes [GJ99, §V, Lem. 6.6]). *While the forgetful functor from (75) to (74)*

$$\Delta\text{Set}_{\geq 1, \text{inj}} \xrightarrow{\text{undrlng}} \Delta\text{Set}_{\text{Qu}}$$

does not preserve all fibrations, it does preserve fibrant objects.

Simplicial Groups.

Notation A.12 (Homotopy theory of simplicial groups, e.g. [Qu67, §II 3.7][GJ99, §V]). We write

$$\text{Grp}(\Delta\text{Set})_{\text{proj}} \in \text{MdlCat}$$

for the model category of simplicial groups ([May67, §17][Cu71, §3]), whose weak equivalences and fibrations are those of the underlying $\Delta\text{Set}_{\text{Qu}}$ (74).

Lemma A.13 (Simplicial groups are Kan complexes, e.g. [Mo54, Thm. 3][May67, Thm. 17.1][Cu71, Lem. 3.1]). *The underlying simplicial set of any simplicial group is a Kan complex.*

Proposition A.14 (Quillen equivalence between simplicial groups and reduced simplicial sets [GJ99, §V, Prop. 6.3]). *The simplicial classifying space construction (Def. A.20) is the right adjoint of a Quillen equivalence between the projective model structure on simplicial groups (Nota. A.12) and the injective model structure on reduced simplicial sets (Nota A.10):*

$$\text{Grp}(\Delta\text{Set})_{\text{proj}} \begin{array}{c} \xleftarrow{\simeq_{\text{Qu}}} \\ \xrightarrow{\overline{W}(-)} \end{array} (\Delta\text{Set}_{\geq 1})_{\text{inj}} .$$

Abelian simplicial groups and chain complexes.

Notation A.15 (Dold-Kan correspondence – e.g. [GJ99, §III.2][SS03, §2]). We write:

- $(\text{AbGrp}(\Delta\text{Set}), \otimes)$ for the category of simplicial abelian groups equipped with the degree-wise tensor product of abelian groups;
- $(\text{Ch}^{\geq 0}(\text{AbGrp}), \otimes)$ for the category of connective chain complexes of abelian groups, equipped with its natural tensor product;
- the Dold-Kan correspondence (reviewed in our context in [FSS20Cha, A.63]):

$$\begin{array}{ccccc} \text{Ch}^{\geq 0}(\text{AbGrp}) & \xleftarrow{N_{\bullet}} & \text{AbGrp}(\Delta\text{Set}) & \xleftarrow{\mathbb{Z}[-]} & \Delta\text{Set}, \\ & \xrightarrow{\Gamma} & & \xrightarrow{\text{undrlng}} & \\ & & & & \uparrow \\ & & & & \text{DK} \end{array} \quad (76)$$

where $N_{\bullet}(-)$ forms (“normalized”) chain complexes of non-degenerate cells, and

where $\mathbb{Z}[-]$ denotes the (degreewise) free abelian group functor;

- notice here that $N_{\bullet}(-)$ respects the tensor product only up to homotopy (this is the content of Prop. A.19 below), while $\mathbb{Z}[-]$ is strong monoidal:

$$X, Y : \Delta\text{Set} \quad \vdash \quad \mathbb{Z}[X \times Y] \simeq \mathbb{Z}[X] \otimes \mathbb{Z}[Y]; \quad (77)$$

- the Eilenberg-MacLane formula [EML53, (5.3)] for the Eilenberg-Zilber map [EZ53] (review in [Lo92, §1.6]):

$$A, B \in \text{AbGrp}(\Delta\text{Set}) \quad \vdash \quad \begin{array}{ccc} N_{\bullet}(A) \otimes N_{\bullet}(B) & \xrightarrow{\nabla_{A,B}} & N_{\bullet}(A \otimes B) \\ a_p \otimes b_q & \longmapsto & \sum_{(\mu, \nu)} \text{sgn}(\mu, \nu) \cdot (s_{\nu}(a_p)) \otimes (s_{\mu}(b_q)), \end{array} \quad (78)$$

where the sum is over all (p, q) -shuffles (μ, ν) from (65) and $(s_{\nu}(-), s_{\mu}(-))$ is according to (67).

Example A.16 (Shifted abelian groups). For $A \in \text{AbGrp}(\text{Set})$ and $n \in \mathbb{N}$, the image under the Dold-Kan construction (76) of the chain complex that is concentrated on A in degree n is a model for the n -fold delooping of A :

$$\mathbf{B}^n A \simeq \text{DK}(A[n]) \in \text{Ho}(\Delta\text{Set}_{\text{Qu}}).$$

Example A.17 (Incarnations of group cohomology). For $G, A \in \text{Grp}(\text{Set}) \hookrightarrow \text{Grp}(\Delta\text{Set}) \xrightarrow{\text{Loc}_W} \text{Grp}(\text{Grpd}_\infty)$ with A abelian, the following abelian groups are all naturally isomorphic, for all $n \in \mathbb{N}$:

$$H_{\text{grp}}^{n+1}(G, A) \simeq \pi_0 \Delta\text{Set} \left(\overline{WG}, \text{DK}(A[n]) \right) \simeq H^{n+1}(BG, A) \simeq \pi_0 \text{Grpd}_\infty(BG, \mathbf{B}^{n+1}A).$$

group cohomology
homotopy classes of homs of delooped simplicial groups
cohomology of classifying space
homotopy classes of homs of delooped ∞ -groups

Example A.18 (Normalized chains on minimal simplicial circle). The normalized chains complex of the free simplicial abelian group (76) on the minimal simplicial circle (Ntn. A.24) is

$$N_\bullet(\mathbb{Z}[S_{\text{min}}^1]) = N_\bullet \left(\begin{array}{c} \mathbb{Z}[\{*, \ell\}] \\ \downarrow \quad \uparrow \quad \downarrow \\ \mathbb{Z}[\{*\}] \end{array} \right) = \begin{array}{c} \mathbb{Z}[\{\ell\}] \\ \downarrow_{\partial=0} \\ \mathbb{Z}[\{*\}] \end{array} = \mathbb{Z} \oplus \mathbb{Z}[1]. \quad (79)$$

Proposition A.19 (Eilenberg-Zilber/Alexander-Whitney deformation retraction). *The Eilenberg-Zilber map $\nabla_{A,B}$ (78) on normalized chain complexes is a homotopy equivalence, in fact, it has a deformation retraction (given by the Alexander-Whitney map $\Delta_{A,B}$):*

$$\begin{array}{ccccccc} & & \text{id} & & & & \\ & \swarrow & \parallel & \searrow & & & \\ N_\bullet(A) \otimes N_\bullet(B) & \xrightarrow{\nabla_{A,B}} & N_\bullet(A \otimes B) & \xrightarrow{\Delta_{A,B}} & N_\bullet(A) \otimes N_\bullet(B) & \xrightarrow{\nabla_{A,B}} & N_\bullet(A \otimes B) \\ & & & & \downarrow & & \\ & & & & \text{id} & & \end{array}$$

For the case of unnormalized chain complexes (where we have just an homotopy equivalence) this is due to [EZ53], the case for normalized chain complexes is due to [EML54, Thm. 2.1a], both are reviewed in [May67, Cor. 29.10]. Explicit description of the homotopy operator is in [GDR99].

Universal principal simplicial complex.

Definition A.20 (Universal principal simplicial complex [Kan58, Def. 10.3][GJ99, p. 269]). Let $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$. (i) Its *standard universal principal complex* is the simplicial set

$$W\mathcal{G} \in \Delta\text{Set}$$

whose

- component sets are

$$(W\mathcal{G})_n := \mathcal{G}_n \times \mathcal{G}_{n-1} \times \cdots \times \mathcal{G}_0,$$

- face maps are given by

$$d_i(\gamma_n, \gamma_{n-1}, \cdots, \gamma_0) := \begin{cases} (d_i(\gamma_n), d_{i-1}(\gamma_{n-1}), \cdots, d_0(\gamma_{n-i}) \cdot \gamma_{n-i-1}, \gamma_{n-i-2}, \cdots, \gamma_0) & \text{for } 0 < i < n \\ (d_n(\gamma_n), d_{n-1}(\gamma_{n-1}), \cdots, d_1(\gamma_1)) & \text{for } i = n, \end{cases} \quad (80)$$

- degeneracy maps are given by

$$s_i(\gamma_n, \gamma_{n-1}, \cdots, \gamma_0) := (s_i(\gamma_n), s_{i-1}(\gamma_{n-1}), \cdots, s_0(\gamma_{n-i}), e, \gamma_{n-i-1}, \cdots, \gamma_0), \quad (81)$$

- and equipped with the left \mathcal{G} -action (Ex. A.29) given by

$$\begin{array}{ccc} \mathcal{G} \times W\mathcal{G} & \longrightarrow & W\mathcal{G} \\ (h_n, (\gamma_n, \gamma_{n-1}, \cdots, \gamma_0)) & \longmapsto & (h_n \cdot \gamma_n, \gamma_{n-1}, \cdots, \gamma_0). \end{array} \quad (82)$$

(ii) Its standard *simplicial delooping* or *simplicial classifying complex* $\overline{W}\mathcal{G}$ is the quotient by that action (82):

$$W\mathcal{G} \xrightarrow{q_{W\mathcal{G}}} \overline{W}\mathcal{G} := (W\mathcal{G})/\mathcal{G}. \quad (83)$$

Example A.21 (Low-dimensional cells of universal simplicial principal complex). Unwinding the definition (80) of the face maps of $W\mathcal{G}$ (Def. A.20) shows that its 1-simplices are of the form

$$(W\mathcal{G})_1 = \left\{ d_1(g_1) \xrightarrow{(g_1, g_0)} d_0(g_1) \cdot g_0 \mid \begin{array}{l} g_0 \in \mathcal{G}_0 \\ g_1 \in \mathcal{G}_1 \end{array} \right\}$$

and its 2-simplices are of this form:

$$(W\mathcal{G})_2 = \left\{ \begin{array}{ccc} & d_0 d_2(g_2) \cdot d_1(g_1) = d_1 d_0(g_2) \cdot d_1(g_1) & \\ \begin{array}{c} \nearrow (d_1(g_2), d_1(g_1)) \\ \searrow (d_0(g_2), g_1, g_0) \end{array} & \begin{array}{c} \Downarrow (g_2, g_1, g_0) \\ \Downarrow \end{array} & \\ d_1 d_2(g_2) & \xrightarrow{(d_1(g_2), d_0(g_1), g_0)} & d_0 d_0(g_2) \cdot d_0(g_1) \cdot g_0 \\ = d_1 d_1(g_2) & & = d_0 d_1(g_2) \cdot d_0(g_1) \cdot g_0 \end{array} \mid \begin{array}{l} g_0 \in \mathcal{G}_0, \\ g_1 \in \mathcal{G}_1, \\ g_2 \in \mathcal{G}_2 \end{array} \right\}.$$

Example A.22 (Universal principal simplicial complex for ordinary group G). If

$$G \in \text{Grp} \hookrightarrow \text{Grp}(\Delta\text{Set})$$

is an ordinary discrete group, regarded as a simplicial group (hence the functor constant on G on the opposite simplex category), then the standard model of its universal principal complex (Def. A.20) is isomorphic to the nerve of the action groupoid of the right multiplication action of G on itself:

$$WG = N(G \times G \rightrightarrows G). \quad (84)$$

$$(WG)_2 = \left\{ \begin{array}{ccc} & g_2 g_1 & \\ \begin{array}{c} \nearrow (g_2, g_1) \\ \searrow (g_2, g_1, g_0) \end{array} & \begin{array}{c} \Downarrow (g_2, g_1, g_0) \\ \Downarrow \end{array} & \\ g_2 & \xrightarrow{(g_2, g_1 g_0)} & g_2 g_1 g_0 \end{array} \mid g_0, g_1, g_2 \in G \right\}$$

Accordingly, the standard simplicial delooping (83) of an ordinary group is isomorphic to the simplicial nerve of its delooping groupoid:

$$\overline{W}G \simeq N(\underbrace{G \rightrightarrows *}_{=: \mathbf{B}G}) \in \Delta\text{Set}. \quad (85)$$

Proposition A.23 (Basic properties of standard simplicial principal complex [GJ99, §V, Lem. 4.1, 4.6, Cor. 6.8]). For $\mathcal{G} \in \text{Grp}(\text{SimplSets})$, its standard universal principal complex (Def. A.20) has the following properties:

- (i) $W\mathcal{G}$ is contractible;
- (ii) $W\mathcal{G}$ and $\overline{W}\mathcal{G}$ are Kan complexes;

Proof. That $\overline{W}\mathcal{G}$ is Kan fibrant follows as the combination of Lem. A.13, Prop. A.14, and Lem. A.11. This implies that $W\mathcal{G}$ is Kan fibrant since $W\mathcal{G} \xrightarrow{q} \overline{W}\mathcal{G}$ is a Kan fibration (99) (by Prop. A.14, see Ex. A.35). \square

Inertia groupoids.

Notation A.24 (Minimal simplicial circle). We write

$$S_{\min}^1 := \Delta[1]/\partial\Delta[1] \in \Delta\text{Set},$$

and denote its unique non-degenerate 1-cell by

$$\ell := [0, 1] \in \Delta[1] \twoheadrightarrow S_{\min}^1. \quad (86)$$

Simplicial group actions.

Notation A.27 (Simplicial group actions). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, we denote

(i) by

$$\mathbf{B}\mathcal{G} \in \Delta\text{SetEnrCat}, \quad \mathbf{B}\mathcal{G}(*, *) := \mathcal{G}, \quad (89)$$

the simplicial groupoid with a single object $*$, with \mathcal{G} as its unique hom-object and with composition “ \circ ” given by the *reverse* of the group product “ \cdot ”

$$\begin{array}{ccc} \mathcal{G} \times \mathcal{G} & \xrightarrow{\circ} & \mathcal{G} \\ (g_n, h_n) & \longmapsto & h_n \cdot g_n \end{array} \quad (90)$$

(ii) the category of \mathcal{G} -actions on simplicial sets by:

$$\mathcal{G} \text{ Act}(\Delta\text{Set}) := \text{Fnctr}_{\Delta}(\mathbf{B}\mathcal{G}, \Delta\text{Set}), \quad (91)$$

identified with the category of ΔSet -enriched functors from the delooping (89) of \mathcal{G} to ΔSet (via Rem. A.7).

Remark A.28 (Simplicial group actions are from the left). The convention (90) for the delooping $\mathbf{B}\mathcal{G}$ (89) implies that the simplicial \mathcal{G} -actions (91) are *left* actions:

$$\begin{array}{ccc} \mathbf{B}\mathcal{G} & \xrightarrow{\mathcal{G} \zeta X} & \Delta\text{Set} \\ \left. \begin{array}{c} \bullet \\ \downarrow g_1 \\ \bullet \\ \downarrow g_2 \\ \bullet \end{array} \right\} g_1 \circ g_2 := g_2 \cdot g_1 & & \left. \begin{array}{c} X \\ \downarrow g_1 \cdot (-) \\ X \\ \downarrow g_2 \cdot (-) \\ X \end{array} \right\} (-) \cdot (-) \end{array} \quad \mathcal{G} \times X \xrightarrow{(-) \cdot (-)} X$$

$$(g_n, h_n) \longmapsto g_n \cdot x_n$$

Example A.29 (Universal principal simplicial complex in \mathcal{G} -actions). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, the universal principal simplicial complex $W\mathcal{G}$ (Def. A.20) becomes an object of (91) by the formula (82).

$$\mathcal{G} \zeta W\mathcal{G} \in \mathcal{G} \text{ Act}(\Delta\text{Set}). \quad (92)$$

Making explicit the following elementary Ex. A.30 serves to straighten out a web of conventions about (simplicial) group actions.

Example A.30 (Simplicial group canonically acting on itself). Any $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$ becomes an object of the category of simplicial \mathcal{G} -actions (91) in three canonical ways:

$$\begin{array}{ccc} \mathcal{G} \times \mathcal{G} \xrightarrow[\text{multiplication action}]{\text{left}} \mathcal{G}, & \mathcal{G} \times \mathcal{G} \xrightarrow[\text{multiplication action}]{\text{right inverse}} \mathcal{G}, & \mathcal{G} \times \mathcal{G} \xrightarrow[\text{action}]{\text{adjoint/conjugation}} \mathcal{G}. \\ (g_n, h_n) \longmapsto g_n \cdot h_n & (g_n, h_n) \longmapsto h_n \cdot g_n^{-1} & (g_n, h_n) \longmapsto g_n \cdot h_n \cdot g_n^{-1} \end{array} \quad (93)$$

The first two are isomorphic in $\mathcal{G} \text{ Act}(\Delta\text{Set})$ via the inversion operation:

$$\begin{array}{ccc} (g_n, h_n) & \xrightarrow{\quad} & g_n \cdot h_n \\ \downarrow & \begin{array}{ccc} \mathcal{G} \times \mathcal{G} & \xrightarrow{\text{left multiplication}} & \mathcal{G} \\ \text{id} \times (-)^{-1} \downarrow \wr & & \downarrow \wr (-)^{-1} \\ \mathcal{G} \times \mathcal{G} & \xrightarrow{\text{right inverse multiplication}} & \mathcal{G} \end{array} & \downarrow \\ (g_n, h_n^{-1}) & \xrightarrow{\quad} & h_n^{-1} \cdot g_n^{-1} \end{array} \quad (94)$$

Homotopy theory of simplicial group actions.

Notation A.31 (Model category of simplicial group actions ([DDK80, §2][Gui06, §5][GJ99, §V Thm. 2.3])). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, we have on the category of \mathcal{G} -actions (91) the projective model structure (the *coarse-* or *Borel-equivariant model structure*) whose fibrations and weak equivalences are those of the underlying $\Delta\text{Set}_{\text{Qu}}$ (Ntn. A.49), which we denote as:

$$\mathcal{G} \text{ Act}(\Delta\text{Set})_{\text{proj}} := \text{Fnctr}_{\Delta}(\mathbf{B}\mathcal{G}, \Delta\text{Set})_{\text{proj}}. \quad (95)$$

Lemma A.32 (Cofibrations of simplicial group actions [DDK80, Prop. 2.2 (ii)][Gui06, Prop. 5.3][GJ99, §V Lem. 2.4]). *The cofibrations of $\mathcal{G} \text{ Act}(\Delta\text{Set})_{\text{proj}}$ (95) are the monomorphisms such that the \mathcal{G} -action on the simplices not in their image is free.*

Lemma A.33 (Equivariant equivalence of simplicial universal principal complexes). *For $\mathcal{H} \xrightarrow{i} \mathcal{G}$ a simplicial subgroup inclusion, the induced inclusion*

$$W\mathcal{H} \xrightarrow[\in W]{W(i)} W\mathcal{G} \in \mathcal{H} \text{ Act}(\Delta\text{Set})_{\text{proj}}$$

of their standard simplicial principal complexes (Def. A.20) equipped with their canonical \mathcal{H} -action (82) is a weak equivalence in the Borel-equivariant model structure (95)

Proof. The underlying simplicial sets of both are contractible, by Prop. A.23, so that underlying any equivariant morphism between them is an simplicial weak homotopy equivalence. \square

The following Prop. A.34 is the model category theoretic avatar of the slice characterization of ∞ -group actions in (31):

Proposition A.34 (Quillen equivalence between Borel model structure and the slice over classifying complex). *For any $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$ there is a simplicial adjunction*

$$\begin{array}{c} \text{homotopy fiber} \\ (-) \times_{\overline{W}\mathcal{G}} W\mathcal{G} \end{array} \dashv \begin{array}{c} \text{Borel construction} \\ ((-) \times W\mathcal{G})/\mathcal{G} \end{array} \quad (96)$$

between the Borel model structure (95) and the slice model structure of $\Delta\text{Set}_{\text{Qu}}$ (74) over the simplicial classifying complex $\overline{W}\mathcal{G}$ (83), hence a natural isomorphism of hom-complexes

$$\mathcal{G} \text{ Act}(\Delta\text{Set})((-) \times_{\overline{W}\mathcal{G}} W\mathcal{G}, (-)) \simeq \Delta\text{Set}_{/\overline{W}\mathcal{G}}\left((-), ((-) \times W\mathcal{G})/\mathcal{G}\right) \in \Delta\text{Set}, \quad (97)$$

which is a Quillen equivalence:

$$\mathcal{G} \text{ Act}(\Delta\text{Set})_{\text{proj}} \begin{array}{c} \xleftarrow{(-) \times_{\overline{W}\mathcal{G}} W\mathcal{G}} \\ \xrightarrow{((-) \times W\mathcal{G})/\mathcal{G}} \end{array} (\Delta\text{Set}_{\text{Qu}})_{/\overline{W}\mathcal{G}}. \quad (98)$$

Proof. The plain adjunction in (98) is [DDK80, Prop. 2.3]. The simplicial enrichment (97), hence the natural bijections

$$\text{Hom}\left(\left((-) \times_{\overline{W}\mathcal{G}}\right) \times \Delta[k], (-)\right) \simeq \text{Hom}\left((-) \times \Delta[k], ((-) \times W\mathcal{G})/\mathcal{G}\right) \in \text{Set},$$

are left somewhat implicit in [DDK80, Prop. 2.4], but follows readily from the plain adjunction via the natural isomorphism

$$\left((-) \times_{\overline{W}\mathcal{G}} W\mathcal{G}\right) \times \Delta[k] \simeq \left((-) \times \Delta[k]\right) \times_{\overline{W}\mathcal{G}} W\mathcal{G},$$

which, in turn, follows from the pasting law (Fact A.46):

$$\begin{array}{ccc} (X \times_{\overline{W}\mathcal{G}} W\mathcal{G}) \times \Delta[k] \simeq (X \times \Delta[k]) \times_{\overline{W}\mathcal{G}} W\mathcal{G} & \longrightarrow & X \times \Delta[k] \\ \downarrow & \text{(pb)} & \downarrow \text{pr}_1 \\ X \times_{\overline{W}\mathcal{G}} W\mathcal{G} & \longrightarrow & X \\ \downarrow & \text{(pb)} & \downarrow \\ W\mathcal{G} & \longrightarrow & \overline{W}\mathcal{G}. \end{array} \quad \square$$

Example A.35 (Coprojections out of Borel construction are Kan fibrations). For $\mathcal{G} \curvearrowright X \in \mathcal{G} \text{ Act}(\Delta\text{Set})$ such that the underlying simplicial set X is a Kan complex, hence such that

$$\mathcal{G} \curvearrowright X \xrightarrow{\in \text{Fib}} * \in \mathcal{G} \text{ Act}(\Delta\text{Set})_{\text{proj}},$$

the projection from the Borel construction (96) to the simplicial classifying space (Def. A.20) is a Kan fibration, due to the right Quillen functor property (98):

$$\begin{array}{ccc} X & \xrightarrow{\text{fib}(q)} & (W\mathcal{G} \times X)/\mathcal{G} \\ & & \downarrow q \in \text{Fib} \\ & & \overline{W}\mathcal{G} \end{array} = \left(W\mathcal{G} \times \left(\begin{array}{c} X \\ \downarrow \\ * \end{array} \right) \right) / \mathcal{G}.$$

The fiber of this fibration, hence the *homotopy fiber*, is clearly X .

In the special case where $\mathcal{G} \curvearrowright X = \mathcal{G} \curvearrowright \mathcal{G}_L$ is the multiplication action of the simplicial group on itself (Exp. A.30), this construction reduces to the *universal principal simplicial bundle* (83):

$$\begin{array}{ccc} \mathcal{G} & \longrightarrow & W\mathcal{G} \\ & & \downarrow q \in \text{Fib} \\ & & \overline{W}\mathcal{G} \end{array} = \left(W\mathcal{G} \times \left(\begin{array}{c} \mathcal{G}_L \\ \downarrow \\ * \end{array} \right) \right) / \mathcal{G} \quad (99)$$

Notation A.36 (Homotopy quotient of simplicial group actions). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, we denote the right derived functor of the *Borel construction* right Quillen functor (98) by:

$$(-) // \mathcal{G} := \mathbb{R}((-) \times W\mathcal{G}) / \mathcal{G} : \text{Ho}(\mathcal{G} \text{ Act}(\Delta\text{Set})_{\text{proj}}) \longrightarrow \text{Ho}(\Delta\text{Set}_{\text{Qu}}).$$

Applied to the point $\mathcal{G} \curvearrowright *$ we also write

$$\mathbf{B}\mathcal{G} := * // \mathcal{G}. \quad (100)$$

As an example:

Proposition A.37 (*G*-Sets in the homotopy theory over *BG*). For $G \in \text{Grp}(\text{Set}) \hookrightarrow \text{Grp}(\Delta\text{Set})$, we have an equivalence between *G*-actions on sets and 0-truncated objects in the homotopy theory slice over *BG*:

$$\begin{array}{ccc} G \text{ Act}(\text{Set}) & \simeq & ((\text{Grpd}_\infty)_{/BG})_{\leq 0} \longrightarrow (\text{Grpd}_\infty)_{/BG} \\ G \curvearrowright X & \longmapsto & (X // G \rightarrow BG) \end{array}$$

Proof. Since $(X \times G \rightrightarrows X) \rightarrow (G \rightrightarrows *)$ is clearly a Kan fibration with fiber X , the latter is the homotopy fiber of $X // G \xrightarrow{B} G$. With this, the statement follows by Prop. A.34. \square

Free loop space of classifying spaces.

Proposition A.38 (Free loop space of simplicial classifying space). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, the function complex from the simplicial classifying space (Def. A.20) of the additive group of integers to that of \mathcal{G} is equivalent to the homotopy quotient (Nota. A.36) of the conjugation action of \mathcal{G} on itself (Ex. A.30)

$$\text{Map}(\overline{W}\mathbb{Z}, \overline{W}\mathcal{G}) \simeq \mathcal{G} //_{\text{ad}} \mathcal{G} \in \text{Ho}(\Delta\text{Set}_{\text{Qu}}).$$

For topological groups this statement is folklore but rarely argued in detail. A point-set topology argument is spelled out in [Gr07, §A], and the idea of the following abstract argument is in [KSS09, Lem. 9.1], which we adapt to simplicial groups:

Proof. Consider the following commuting diagram in ΔSet :

$$\begin{array}{ccc}
\frac{W\mathcal{G} \times \mathcal{G}_{\text{ad}}}{\mathcal{G}} & \xrightarrow{\quad} & \frac{W\mathcal{G} \times W\mathcal{G} \times \mathcal{G}_{\text{ad}}}{\mathcal{G} \times \mathcal{G}} & \xrightarrow{\quad} & \frac{W\mathcal{G}}{\mathcal{G}} \times W\mathcal{G} \\
\downarrow & \text{(pb)} & \downarrow \in \text{Fib} & \xrightarrow{[\bar{g}, (\bar{g}, h)] \mapsto ([\bar{g}], (\bar{g}^{-1}, (h; \bar{g}')))} & \downarrow \text{pr}_1 \in W \\
\frac{W\mathcal{G}}{\mathcal{G}} & \xrightarrow{\quad} & \frac{W\mathcal{G} \times W\mathcal{G}}{\mathcal{G} \times \mathcal{G}} & \xrightarrow{[\bar{g}, (\bar{g}, e)] \mapsto [\bar{g}]} & \frac{W\mathcal{G}}{\mathcal{G}} \\
& \text{diag} & & \in W & \\
& & & \text{diag} &
\end{array} \tag{101}$$

where

$$\mathcal{G}_{\text{ad}} \in (\mathcal{G} \times \mathcal{G}) \text{Act}(\Delta\text{Set}) \xrightarrow{\text{res}_{\text{diag}/\mathcal{G}}} \mathcal{G} \text{Act}(\Delta\text{Set})$$

denotes the conjugation action on $\mathcal{G} = \text{undrlng}(\mathcal{G}_{\text{ad}})$

$$\begin{array}{ccc}
\mathcal{G}_{\text{ad}} \times (\mathcal{G} \times \mathcal{G}) & \longrightarrow & \mathcal{G}_{\text{ad}} \\
(g_k, (h'_k, h_k)) & \longmapsto & h'_k \cdot g_k \cdot h_k^{-1}
\end{array}$$

and its restriction along the diagonal to $h' = h$. Here

- the right vertical morphism is a Kan fibration by Lemma A.13 and Prop. A.34 (see Ex. A.35);
- all objects are fibrant (are Kan complexes), by Prop. A.23 and by the previous item;
- the horizontal morphism on the top right is a weak equivalence, by 2-out-of-3, as it is the left inverse to the triangular composite of morphisms, which is a weak equivalence by Prop. A.23.

Therefore (as recalled in [FSS20Cha, Def. A.24]), the pullback diagram in (101) represents the homotopy pullback that characterizes $[\overline{W\mathbb{Z}}, \overline{W\mathcal{G}}]$, according to Example A.25. \square

As an immediate consequence:

Proposition A.39 (Free loop space of classifying space of simplicial abelian group).

Let $\mathcal{A} \in \text{AbGrp}(\Delta\text{Set}) \xrightarrow{\text{Disc}} \text{Grp}(\text{SmthGrpd}_\infty)$ be a simplicial abelian group. Then

$$\text{Map}(\mathbf{B}\mathbb{Z}, \mathbf{B}\mathcal{A}) \simeq \mathcal{A} \times \mathbf{B}\mathcal{A} \quad \text{and} \quad \text{Cyc}_{\text{JS}^1_{\text{coh}}}(\mathbf{B}\mathcal{A}) \simeq \mathbf{B}^2\mathbb{Z} \times \mathcal{A} \times \mathbf{B}\mathcal{A}.$$

Proof. This is given by the following sequence of equivalences:

$$\begin{aligned}
\text{Map}(\mathbf{B}\mathbb{Z}, \mathbf{B}\mathcal{A}) &\simeq \frac{\mathcal{A}_{\text{ad}} \times W\mathcal{A}}{\mathcal{A}} && \text{by Prop. A.38} \\
&\simeq \frac{\mathcal{A}_{\text{triv}} \times W\mathcal{A}}{\mathcal{A}} && \text{since each } \mathcal{A}_n \text{ is abelian} \\
&\simeq \mathcal{A} \times \frac{W\mathcal{A}}{\mathcal{A}} \\
&\simeq \mathcal{A} \times \overline{W\mathcal{A}} && \text{by (83)} \\
&\simeq \mathcal{A} \times \mathbf{B}\mathcal{A} && \text{by (100)}.
\end{aligned} \quad \square$$

Example A.40 (Free loop space of Eilenberg-MacLane space). For $A \in \text{AbGrp}(\text{Set})$ with $\mathbf{B}^n A \simeq K(A, n)$ for $n \in \mathbb{N}$ (Ex. A.16), Prop. A.39 yields:

$$\text{Map}(\mathbf{B}\mathbb{Z}, \mathbf{B}^{n+1}A) \simeq \mathbf{B}^n A \times \mathbf{B}^{n+1}A.$$

Free and co-free simplicial actions.

Proposition A.41 (Free and co-free simplicial actions). For $\mathcal{G} \in \text{Grp}(\Delta\text{Set})$, the forgetful functor that sends \mathcal{G} -actions (91) to their underlying simplicial set has both a left adjoint (“free action”) and a right adjoint (“co-free action”), both of which are Quillen adjunctions with respect to the classical model structure on ΔSet and the projective model structure on $\mathcal{G} \text{Act}(\Delta\text{Set})$ (95):

$$\begin{array}{ccc}
& \xrightarrow{\mathcal{G} \times (-)} & \\
\Delta\text{Set}_{\text{Qu}} & \xleftarrow{\perp_{\text{Qu}}} \text{undrlng} \text{ --- } \mathcal{G} \text{Act}(\Delta\text{Set}_{\text{Qu}})_{\text{proj}} & \\
& \xrightarrow{\perp_{\text{Qu}}} & \\
& \xrightarrow{[\mathcal{G}, -]} &
\end{array}$$

Here the underlying object of the co-free action on $X \in \Delta\text{Set}$ is the simplicial function complex (68)

$$\text{Map}(\mathcal{G}, X) = \text{Hom}(\mathcal{G} \times \Delta[\bullet], X) \in \Delta\text{Set}$$

and its \mathcal{G} -action

$$\mathcal{G} \times \text{Map}(\mathcal{G}, X) \xrightarrow{(-)\cdot(-)} \text{Map}(\mathcal{G}, X)$$

in degree $m \in \mathbb{N}$ is the function $\text{Hom}(\Delta[n], \mathcal{G}) \times \text{Hom}(\mathcal{G} \times \Delta[n], X) \rightarrow \text{Hom}(\mathcal{G} \times \Delta[n], X)$ given by

$$(\Delta[n] \xrightarrow{g_n} \mathcal{G}, \mathcal{G} \times \Delta[n] \xrightarrow{\phi} X) \mapsto (\mathcal{G} \times \Delta[n] \xrightarrow{\text{id} \times \text{diag}} \mathcal{G} \times \Delta[n] \times \Delta[n] \xrightarrow{\text{id} \times g_n \times \text{id}} \mathcal{G} \times \mathcal{G} \times \Delta[n] \xrightarrow{(-)\cdot(-) \times \text{id}} \mathcal{G} \times \Delta[n] \xrightarrow{\phi} X). \quad (102)$$

Proof. The undrlng-functor preserves fibrations and weak equivalences by definition of the projective model structure (95), and it preserves cofibrations by Lemma A.32. Therefore it is both a left and a right Quillen adjoint as soon as it is a left and a right adjoint functor at all.

The left adjoint free action is straightforward. Also the right adjoint cofree action follows the same idea as that of the cofree action on topological G -spaces, only that for simplicial sets one cannot argue point-wise as in point-set topology, but needs the more abstract formula (102). The structure of this formula manifestly gives a simplicial homomorphism, and by focusing on its images of the unique non-degenerate top-degree cell, $\iota_n \in (\Delta[n])_n$ (Ntn. A.2), right-adjointness is seen essentially by the standard argument:

It is sufficient to check that we have a hom-isomorphism of the form

$$\left\{ P \xrightarrow{\phi_{(-)}} [\mathcal{G}, X] \right\} \xleftrightarrow{\widetilde{(-)}} \left\{ \text{undrlng}(P) \xrightarrow{\widetilde{\phi}_{(-)}} X \right\} \quad (103)$$

as a bijection natural in $P \in \mathcal{G} \text{Act}(\Delta\text{Set})$ and $X \in \Delta\text{Set}$. So, for

$$\phi_{(-)} : p_n \mapsto (\phi_{p_n} : \mathcal{G} \times \Delta[n] \rightarrow X)$$

on the left in (103), define its candidate adjunct to be

$$\widetilde{\phi}_{(-)} : p_n \mapsto \phi_{p_n}(\mathbf{e}_n, p_n) \in X_n, \quad (104)$$

where $\mathbf{e}_n \in \mathcal{G}_n$ denotes the neutral element in degree $n \in \mathbb{N}$. It is clear that this assignment is a natural transformation in P and X , hence it remains to show that $\widetilde{\phi}$ uniquely determines $\phi_{(-)}$.

To that end, observe for any $g_n \in \mathcal{G}_n$ the following sequence of identifications:

$$\begin{aligned} \phi_{p_n}(g_n, \iota_n) &= \phi_{p_n}(\mathbf{e}_n \cdot g_n, \iota_n) && \text{by the unit law in } \mathcal{G}_n \\ &= (g_n \cdot \phi_{p_n})(\mathbf{e}_n, \iota_n) && \text{by (102)} \\ &= \phi_{g_n \cdot p_n}(\mathbf{e}_n, \iota_n) && \text{by equivariance of } \phi \\ &= \widetilde{\phi}_{g_n \cdot p_n} && \text{by (104)}. \end{aligned}$$

This shows that the morphisms $\phi_{(-)}$ and $\widetilde{\phi}_{(-)}$ uniquely determine each other, establishing the bijection (103) and hence the claimed adjunction. \square

A.1.1 Group cohomology

Proposition A.42 (Finite subgroups of $\text{Sp}(1)$ [Klein1884]). *The finite subgroups $G \hookrightarrow \text{Spin}(3) \simeq \text{SU}(2) \simeq \text{Sp}(1)$ are given, up to conjugacy, by the following classification (where $n \in \mathbb{N}$):*

Label	Finite subgroup of $\text{SU}(2)$	Name of group
\mathbb{A}_n	\mathbb{Z}_{n+1}	Cyclic
\mathbb{D}_{n+4}	$2\mathbb{D}_{n+2}$	Binary dihedral
\mathbb{E}_6	$2\mathbb{T}$	Binary tetrahedral
\mathbb{E}_7	$2\mathbb{O}$	Binary octahedral
\mathbb{E}_8	$2\mathbb{I}$	Binary icosahedral

For pointers to modern proofs, see [HSS18, Rem. A.9].

Proposition A.43 (Integral group cohomology of finite subgroups of $SU(2)$). *The integral group cohomology (Ex. A.17) of the finite subgroups $G \xrightarrow{i} SU(2)$ (Prop. A.42) is as follows:*

$$H_{\text{grp}}^n(G, \mathbb{Z}) \simeq \begin{cases} \mathbb{Z} & \text{for } n = 0 \\ G/[G, G] & \text{for } n = 2 \bmod 4 \\ \mathbb{Z}/|G| & \text{for } n = 0 \bmod 4, n \geq 1; \\ 0 & \text{otherwise.} \end{cases}$$

This is summarized in [EG17, p. 12]. Detailed computation for the three exceptional cases is given in [TZ08, §4]. The vanishing of $H_{\text{grp}}^3(G, \mathbb{Z}) \simeq H_{\text{grp}}^2(G, U(1))$ is also made explicit in [FHHP00, Cor. 3.1].

Proposition A.44 (Pullback of 2nd Chern class to ADE-subgroup). *For $G \xrightarrow{i} SU(2) \simeq \text{Spin}(3)$ a finite subgroup (Prop. A.42) the induced pullback in degree-4 integral group cohomology (Ex. A.17) takes the generator (e.g. [MiSt74][CV98, Lem. 2.1])*

$$c_2 = 1 \in \mathbb{Z} \simeq H^4(BSU(2); \mathbb{Z}) \quad \text{equivalently} \quad \frac{1}{4}p_1 = 1 \in \mathbb{Z} \simeq H^4(B\text{Spin}(3); \mathbb{Z})$$

to a generator (via Prop. A.43)

$$[1] \in \mathbb{Z}/|G| \simeq H^4(G; \mathbb{Z})$$

in that the pullback is the quotient projection $\mathbb{Z} \xrightarrow{q} \mathbb{Z}/|G|$:

$$\begin{array}{ccccccc} H_{\text{grp}}^4(\text{Sp}(1); \mathbb{Z}) & \simeq & H^4(B\text{Sp}(1); \mathbb{Z}) & \simeq & \mathbb{Z} & \ni & \frac{1}{2}p_1 \\ \downarrow i^* & & \downarrow Bi^* & & \downarrow q & & \downarrow \\ H_{\text{grp}}^4(G; \mathbb{Z}) & \simeq & H^4(BG; \mathbb{Z}) & \simeq & \mathbb{Z}/|G| & \ni & [1] \end{array} \quad (105)$$

Proof. This is essentially the statement of [EG17, Prop. 4.1], whose proof is analogous to Lemma 3.1 there, where the analogue of (105) is the top left square of the commuting diagram on p. 11. \square

A.2 Notions of cohesive ∞ -topos theory

In the main text we make free use of basic notions from ∞ -topos theory ([TV05][Lu09][Re10]), and of cohesive ∞ -topos theory [SSS12, §3.1][Sc13] as laid out and developed in [SS20Orb][SS21EPB]. For reference, here we list some key notation and facts.

We first list abstract notions in ∞ -toposes and then recall, in some cases, their presentation by model categories of simplicial presheaves.

Notation A.45 (∞ -Toposes). Given any ∞ -topos \mathbf{H} , we write:

- $(-)//\mathcal{G} : \mathcal{G} \text{ Act}(\mathbf{H}) \rightarrow \mathbf{H}$
for the homotopy quotient construction by ∞ -actions of smooth ∞ -groups $\mathcal{G} \in \text{Grp}(\mathbf{H})$
- $\mathbf{B}\mathcal{G} \simeq *//\mathcal{G} \in \mathbf{H}$
for the delooping (“moduli stack”) of any group ∞ -stack $\mathcal{G} \in \text{Grp}(\text{SmthGrpd}_\infty)$;
- $\mathbf{B}^{n+1}\mathcal{A} = \mathbf{B}(\mathbf{B}^n\mathcal{A})$
for the iterative delooping of $(n+1)$ -fold commutative ∞ -groups

We denote homotopy Cartesian squares by putting the label “(pb)” at their center.

Fact A.46 (Homotopy pasting law, [Hir02, Pro. 13.3.15][Lu09, Lem. 4.4.2.1]). *Given a pasting diagram of homotopy commutative squares*

$$\begin{array}{ccc} \longrightarrow & \longrightarrow & \\ \downarrow & \downarrow & \downarrow \\ \longrightarrow & \longrightarrow & \downarrow \\ & & \text{(pb)} \end{array}$$

where the right square is homotopy cartesian (is a homotopy pullback), then the left square is so if and only if the total rectangle is. (NB: In 1-categories this restricts to the pasting law for plain Cartesian/pullback squares.)

Fact A.47 (Fundamental theorem of ∞ -topos theory, [Lu09, §6.3.5]). *Given an ∞ -topos \mathbf{H} , then:*

- (i) *for every object $\mathcal{X} \in \mathbf{H}$ the slice ∞ -category $\mathbf{H}/_{\mathcal{X}}$ is again an ∞ -topos;*
- (ii) *for every morphism $\mathcal{X} \xrightarrow{f} \mathcal{Y}$ in \mathbf{H} there is an induced base change adjoint triple:*

$$\mathbf{H}/_{\mathcal{X}} \begin{array}{c} \xrightarrow{f_!} \\ \perp \\ \xleftarrow{f^*} \\ \perp \\ \xrightarrow{f_*} \end{array} \mathbf{H}/_{\mathcal{Y}}, \quad (106)$$

where, in terms of \mathbf{H} , f^* is given by pullback along f and $f_!$ by postcomposition with f .

Notation A.48 (Cohesive/smooth ∞ -toposes). We write:

- $\text{SmthGrpd}_{\infty} := \text{Sh}_{\infty}(\text{CartSpc})$
for the ∞ -topos over the site of Cartesian spaces (equivalently over all smooth manifolds) with smooth functions between them and with respect to differentiably good open covers, presented by the projective local model structure on simplicial presheaves over this site (cf. Def. A.50).
- $\text{SmthMnfd} \hookrightarrow \text{SmthOrbfd} \hookrightarrow \text{SmthGrpd}_{\infty}$
for the full inclusion of the $(2, 1)$ -category of orbifolds regarded as differentiable stacks, among these the *good orbifolds* are the global homotopy quotients $X // G \in \text{SmthGrpd}_{\infty}$, for $G \curvearrowright X$ a smooth action of a discrete group on a smooth manifold X .
- $\text{Map}(\mathcal{X}, \mathcal{A}) \in \text{SmthGrpd}_{\infty}$
for the stack of maps between $\mathcal{X}, \mathcal{A} \in \text{SmthGrpd}_{\infty}$ (their internal hom).

$$\begin{array}{ccc} \text{SmthGrpd}_{\infty} & \xrightarrow{\text{shape modality } \int} & \text{SmthGrpd}_{\infty} \\ \parallel \dashv \! \! \! \dashv & \searrow \text{Disc} & \nearrow \text{Disc} \\ & \text{Grpd}_{\infty} & \end{array} \quad (-) \xrightarrow[\eta_{(-)}^{\int}]{\text{shape unit}} \int(-), \quad \int \circ \int \xrightarrow{\sim} \int \quad \text{shape idempotency} \quad (107)$$

for the *shape monad* sending any stack to the homotopy type of its fat geometric realization [SS20Orb, Ex. 3.18], re-embedded as a geometrically discrete ∞ -stack.

- $S_{\text{coh}}^1 \in \text{DTopSp} \hookrightarrow \text{DfflgSpc} \hookrightarrow \text{SmthGrpd}_{\infty}$.
for the *cohesive circle*, i.e. the circle with its usual structure of a topological space⁸, so that
 $\int S_{\text{coh}}^1 \simeq \mathbf{B}\mathbb{Z} \in \text{Grpd}_1 \hookrightarrow \text{Grpd}_{\infty} \xrightarrow{\text{Disc}} \text{SmthGrpd}_{\infty}$
denotes the homotopy type underlying the circle (cf. Lemma 2.8).

Simplicial presheaves. The possibly earliest reference on the homotopy theory of simplicial presheaves is [Br73], which is still highly recommended reading. A comprehensive modern monograph is [Ja15]. The cohesive example over the site CartSpc originates with [FStS12, App.]. All facts that we need here are concisely reviewed and referenced in [FSS20Cha, §A] and [SS21EPB, §3.2].

Notation A.49 (Model categories of simplicial presheaves). Given a simplicial site (\mathcal{C}, J) , i.e. a small simplicially enriched category (Rem. A.7) equipped with a Grothendieck pre-topology (coverage) on its homotopy category, we write:

- (i) $\Delta\text{PSh}(\mathcal{C}) \in \text{Cat}_{\Delta}$ for the simplicial category of simplicial presheaves on \mathcal{C} ;
- (ii) $\Delta\text{PSh}(\mathcal{C})_{\text{inj}/} \in \text{CombMdlCat}_{\Delta, \text{prop}}$ for the global injective or projective model category structure, respectively;
- (iii) $\Delta\text{PSh}(\mathcal{C}, J)_{\text{inj}/} \in \text{CombMdlCat}_{\Delta, \text{prop}}$ for the J -local injective or projective model category structure, whose weak equivalences are the J -stalk-wise weak equivalences in $\Delta\text{Set}_{\text{Qu}}$ (the “hypercomplete” local model structure).

Definition A.50 (Smooth ∞ -groupoids). We write

$$\text{SmthGrpd}_{\infty} := \text{Loc}_{\mathbf{W}}(\Delta\text{PSh}(\text{CartSp})_{\text{proj}/}) \simeq \text{Loc}_{\mathbf{W}_{\text{loc}}}(\Delta\text{PSh}(\text{SmthMnfd})_{\text{proj}/}) \in 2\text{Ho}(\text{Topos}_{\infty}).$$

⁸We may alternatively regard $S_{\text{coh}}^1 \in \text{SmthMnfd} \hookrightarrow \text{DfflgSpc} \hookrightarrow \text{SmthGrpd}_{\infty}$ with its standard structure of a smooth manifold. All constructions in the main text remain valid, only that in this case they pass through differentiable stacks instead of topological stacks.

for the ∞ -topos presented by the local model structure (Nota. A.49) with respect to differentiably good open covers of Cartesian spaces.

The point of working over the site CartSpc instead of over the (hypercompletely) equivalent site of all smooth manifolds is that it allows for more efficient computations (ultimately related to the fact that smooth manifolds are themselves already glued from Cartesian spaces, hence that the inclusion $\text{CartSpc} \hookrightarrow \text{SmthMnfd}$ exhibits a dense subsite). Namely, over CartSpc

- (i) the simplicial delooping \overline{W} (Def. A.20) of every homotopy-sheaf of simplicial groups is itself a homotopy-sheaf, in that it is fibrant in the local model structure (Ntn. A.31); ([SS21EPB, Prop. 3.3.30][Pa22, Prop. 4.13])
- (ii) cofibrant replacement is still nicely tractable, namely given by (differentiably) *good open covers*.

In the main text we mainly appeal to this second fact, and so in the remainder we spell this out further:

Proposition A.51 (Dugger’s cofibrancy recognition [Du01b, Cor. 9.4]). *Let \mathcal{S} be a 1-site. A sufficient condition for $X \in \text{SimplPSh}(\mathcal{S})_{\text{proj}}^{\text{loc}}$ (Nota. A.49) to be projectively cofibrant is that in each simplicial degree k , the component presheaf $X_k \in \text{PSh}(\mathcal{S})$ is*

- (i) a coproduct $X_k \simeq \coprod_{i_k} U_{i_k}$ of representables $U_{i_k} \in \mathcal{S} \xrightarrow{y} \text{PSh}(\mathcal{S})$;
- (ii) whose degenerate cells split off as a disjoint summand: $X_k \simeq N_k \coprod \text{im}(\sigma)$ for some N_k .

Example A.52 (Basic examples of projectively cofibrant simplicial presheaves). Let \mathcal{S} be a 1-site. Examples of projectively cofibrant simplicial presheaves over \mathcal{S} include:

- (i) every representable $U \in \mathcal{S} \xrightarrow{y} \text{SimplPSh}(\mathcal{S})$;
- (ii) every constant simplicial presheaf $S \in \text{SimpSets} \xrightarrow{\text{const}} \text{SimplPSh}(\mathcal{S})$;

and in joint generalization of these two cases:

- (iv) every product $U \times S$ of a representable with a simplicial set.

In all cases, the defining lifting property is readily checked. Alternatively, these follow with Prop. A.51.

Proposition A.53 (Quillen functor for shape modality on smooth ∞ -groupoids). *The shape-monad (107) on smooth ∞ -groupoids (Def. A.50)*

$$\text{SmthGrpd}_\infty \begin{array}{c} \xrightarrow{\text{Shp}} \\ \xleftarrow{\text{Disc}} \end{array} \text{Grpd}_\infty$$

is equivalently the left derived functor of the colimit operation on simplicial presheaves over Cartesian spaces, regarded as functors $\text{CartSpc}^{\text{op}} \rightarrow \Delta\text{Set}$, in that the following is a Quillen adjunction:

$$\Delta\text{PSh}(\text{CartSpc})_{\text{proj}}^{\text{loc}} \begin{array}{c} \xrightarrow{\text{lim}} \\ \xleftarrow{\text{const}} \end{array} \Delta\text{Set}_{\text{Qu}} \quad (108)$$

Moreover, on a simplicial presheaf satisfying Dugger’s cofibrancy condition (Prop. A.51)

$$\emptyset \xrightarrow{\in \text{Cof}} \coprod_{i_\bullet \in \mathbf{I}_\bullet} \mathbb{R}^{n_{i_\bullet}} \xrightarrow{\in \mathbf{W}} X \in \Delta\text{PSh}(\text{CartSpc})_{\text{proj}}^{\text{loc}} \quad (109)$$

the shape is given by the simplicial set obtained by contracting all copies of Cartesian spaces to the point:

$$\int X \simeq \coprod_{i_\bullet \in \mathbf{I}_\bullet} * \in \Delta\text{Set}_{\text{Qu}}.$$

Proof. First observe that the colimit over a representable functor is the point (e.g. [SS20Orb, Lem. 2.40])

$$\lim_{\rightarrow} \mathbb{R}^n := \lim_{\rightarrow} y(\mathbb{R}^n) \simeq * \in \text{Set} \hookrightarrow \Delta\text{Set}, \quad (110)$$

so that the colimit of a simplicial presheaf of the form (109) is the simplicial set obtained by replacing all copies of Cartesian spaces by a point:

$$\begin{aligned} \lim_{\rightarrow} \left(\coprod_{i_\bullet \in \mathbf{I}_\bullet} \mathbb{R}^{n_{i_\bullet}} \right) &\simeq \left(\lim_{\rightarrow} \mathbb{R}^{n_{i_\bullet}} \right) && \text{since colimits commute with coproducts} \\ &\simeq \coprod_{i_\bullet \in \mathbf{I}_\bullet} * && \text{by (110)}. \end{aligned} \quad (111)$$

Next, it is clear that (108) is a simplicial Quillen adjunction for the *global* projective model structure. To show that it is also Quillen for the local model structure it is hence sufficient, by [Lu09, Cor. A.3.7.2], to see that the right adjoint preserves fibrant objects. By adjunction this is equivalent to the statement that for $\{U_i \hookrightarrow X\}$ a differentiably good open cover, with $U := \coprod_i U_i$, we have a simplicial weak homotopy equivalence

$$\lim_{\rightarrow} y(U^{\times_{\check{X}}}) \xrightarrow{\in W} * . \quad (112)$$

But, by (111), the left hand side of (112) is the simplicial set obtained by contracting summands of the Čech nerve of the good cover to the point. Therefore, since any Cartesian space is contractible, the *nerve theorem* ([MC67, Thm. 2], review in [HatAT, Prop. 4G.3]) implies (112). With this, the last statement follows from the fact that left derived functors may be computed on any cofibrant resolution:

$$\begin{aligned} \int X &\simeq (\mathbb{L}\lim)(X) && \text{by (108)} \\ &\simeq \lim_{\rightarrow} \left(\prod_{i_{\bullet} \in \mathcal{I}_{\bullet}} \mathbb{R}^n \right) && \text{by Prop. A.51} \\ &\simeq \prod_{i_{\bullet} \in \mathcal{I}_{\bullet}} * && \text{by (111)}. \end{aligned} \quad \square$$

Example A.54 (Good open covers are projectively cofibrant resolutions of smooth manifolds).

Any $X \in \text{SmthMnfd} \xrightarrow{y} \Delta\text{PSh}(\text{CartSpc})$ admits a *differentially good open cover* ([FStS12, Prop. A.1]), namely an open cover such that all non-empty finite intersections of patches are *diffeomorphic* to an open ball, and hence to $\mathbb{R}^{\dim(X)}$:

$$\{U_i \simeq \mathbb{R}^{\dim(X)} \hookrightarrow X\}_{i \in I}, \quad \text{s.t.} \quad \forall_{\substack{k \in \mathbb{N} \\ i_0, i_1, \dots, i_k \in I}} U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_k} \underset{\text{diff}}{\simeq} \mathbb{R}^{\dim(X)} \quad \text{if non-empty} . \quad (113)$$

By Dugger's recognition (Prop. A.51) this means that the corresponding Čech nerve is projectively cofibrant; moreover, its canonical morphism to X is clearly a stalkwise weak equivalence, so that it provides a cofibrant resolution of X in the local model structure (Def. A.50):

$$\emptyset \xrightarrow{\in \text{Cof}} U^{\times_{\check{X}}} \xrightarrow{\in W_{\text{loc}}} X \in \Delta\text{PSh}(\text{CartSpc})_{\text{proj, loc}}, \quad \text{for } U := \coprod_i U_i .$$

Proposition A.55 (Shape of smooth manifolds is their homotopy type). *For $X \in \text{SmthMnfd} \xrightarrow{y} \text{SmthGrpd}_{\infty}$ (Def. A.50) their shape is their standard homotopy type.*

Example A.56 (Standard cofibrant resolution of the smooth circle). Considering the smooth circle as the quotient of the real numbers by the integers

$$\mathbb{Z} \xleftarrow{i} \mathbb{R} \xrightarrow{p} S^1 \in \text{SmthMnfd} \xrightarrow{y} \Delta\text{PSh}(\text{CartSpc})_{\text{proj, loc}},$$

Dugger's recognition (Prop. A.51) shows that the Čech nerve of p constitutes a cofibrant resolution of the circle in the projective local model structure (Def. A.50)

$$\emptyset \xrightarrow{\in \text{Cof}} \mathbb{R} \times \mathbb{Z}^{\times_{\bullet}} \xrightarrow{\sim} \mathbb{R}^{\times_{S^1}} \xrightarrow{\in W} S^1 .$$

$$(r, \vec{n}) \quad \longmapsto \quad (r, (r+n_1), (r+n_1+n_2), \dots)$$

Proposition A.57 (Presentation of ∞ -topos by simplicial presheaves). *Let \mathcal{C} be a 1-site. Then the Čech/stalk-local injective or projective model category structure on simplicial presheaves over \mathcal{C} presents the topological/hypercomplete ∞ -topos over \mathcal{C} in that there is an equivalence of homotopy categories*

$$\text{Ho} \left(\Delta\text{PSh}(\mathcal{C})_{\text{inj/proj, loc}} \right) \xrightarrow[\text{Ho}(L)]{\sim} \text{Ho}(\text{Sh}_{\infty}(\mathcal{C}))$$

and for all cofibrant X and fibrant A in $\Delta\text{PSh}(\mathcal{C})$ an equivalence of hom- ∞ -groupoids

$$\Delta\text{PSh}(\mathcal{C})(X, A) \simeq \text{Sh}_{\infty}(L(X), L(A)) . \quad (114)$$

Proposition A.58 (∞ -Yoneda lemma). *For $\mathcal{S} \in \text{Categories}_{\infty}$ we have a fully faithful embedding*

$$\begin{array}{ccc} \mathcal{S} & \xrightarrow{y} & \text{PSh}_{\infty}(\mathcal{S}) \\ U & \longmapsto & \mathcal{S}(-, U) \end{array} \quad (115)$$

and a natural equivalence for $U \in \mathcal{C}$ and $X \in \text{PSh}_{\infty}(\mathcal{S})$:

$$\text{PSh}_{\infty}(y(U), X) \simeq X(U) . \quad (116)$$

References

- [Ac99] B. Acharya, *M theory, Joyce Orbifolds and Super Yang-Mills*, Adv. Theor. Math. Phys. **3** (1999), 227-248, [arXiv:hep-th/9812205].
- [ARZ07] A. Adem, Y. Ruan, and B. Zhang, *A Stringy Product on Twisted Orbifold K-theory*, Morfismos **11** (2007), 33-64, [arXiv:math/0605534].
- [AHHKRW15] M. Alim, B. Haghighat, M. Hecht, A. Klemm, M. Rauch, T. Wotschke, *Wall-crossing holomorphic anomaly and mock modularity of multiple M5-branes*, Comm. Math. Phys. **339** (2015) 773–814 [arXiv:1012.1608][doi:10.1007/s00220-015-2436-3]
- [Al20] L. Alfonsi, *Global Double Field Theory is Higher Kaluza-Klein Theory*, Fortsch. Phys. **68** (2020), [doi:10.1002/prop.202000010], [arXiv:1912.07089].
- [Al85] O. Alvarez, *Topological quantization and cohomology*, Comm. Math. Phys. **100** (1985), 279-309, [euclid:1103943448].
- [BGNX07] K. Behrend, G. Ginot, B. Noohi, and P. Xu, *String topology for stacks*, Astérisque **343** (2012), [numdam:AST_2012__343_R1_0], [arXiv:0712.3857].
- [BBP19] D. Berwick-Evans, P. Boavida de Brito, and D. Pavlov, *Classifying spaces of infinity-sheaves*, [arXiv:1912.10544].
- [BM15] A. Blumberg and M. Mandell, *The homotopy theory of cyclotomic spectra*, Geom. Topol. **19** (2015), 3105-3147, [doi:10.2140/gt.2015.19.3105], [arXiv:1303.1694].
- [BO05] M. Bökstedt and I. Ottosen, *A spectral sequence for string cohomology*, Topology **44** (2005), 1181-1212, [doi:10.1016/j.top.2005.04.006], [arXiv:math/0411571].
- [BEM04] P. Bouwknegt, J. Evslin and V. Mathai, *T-Duality: Topology Change from H-flux*, Commun. Math. Phys. **249** (2004) 383-415 [arXiv:hep-th/0306062] [doi:10.1007/s00220-004-1115-6]
- [BSS18] V. Braunack-Mayer, H. Sati, and U. Schreiber, *Gauge enhancement for Super M-branes via Parameterized stable homotopy theory*, Comm. Math. Phys. (2019), 197–265, [doi:10.1007/s00220-019-03441-4], [arXiv:1805.05987].
- [Br72] G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972, [ISBN:9780080873596].
- [Br73] K. S. Brown, *Abstract Homotopy Theory and Generalized Sheaf Cohomology*, Transactions of the Amer. Math. Soc., **186** (1973) 419-458 [jstor:1996573]
- [Br93] J.-L. Brylinski, *Loop Spaces, Characteristic Classes, and Geometric Quantization*, Birkhäuser (1993), [doi:10.1007/978-0-8176-4731-5].
- [BuSc05] U. Bunke and T. Schick, *On the topology of T-duality*, Rev. Math. Phys. **17** (2005), 77-112, [doi:10.1142/S0129055X05002315], [arXiv:math/0405132].
- [CV98] M. Čadek and J. Vanžura, *On 4-fields and 4-distributions in 8-dimensional vector bundles over 8-complexes*, Colloq. Math. **76** (1998), 213-228, [bwnjournal-article-cmv76z2p213bwm].
- [Cal1] D. Carchedi, *Categorical Properties of Topological and Differentiable Stacks*, PhD Dissertation, Utrecht U. (2011), [dSPACE:1874/208971].
- [Cha05] D. Chataur, *A bordism approach to string topology*, Intern. Math. Res. Not. **2005** (2005) 46, [doi:10.1155/IMRN.2005.2829], [arXiv:math/0306080].
- [CO15] D. Chataur and A. Oancea, *Basics on free loop spaces*, Chapter I in: *Free Loop Spaces in Geometry and Topology*, IRMA Lectures in Mathematics and Theoretical Physics **24**, Europ. Math. Soc. 2015, [ISBN:978-3-03719-153-8], [arXiv:1110.0405].
- [Cu71] E. B. Curtis, *Simplicial homotopy theory*, Adv. Math. **6** (1971), 107-209, [doi:10.1016/0001-8708(71)90015-6].
- [dB+02] J. de Boer, R. Dijkgraaf, K. Hori, A. Keurentjes, J. Morgan, D. Morrison, and S. Sethi, *Triples, Fluxes, and Strings*, Adv. Theor. Math. Phys. **4** (2002), 995-1186, [doi:10.4310/ATMP.2000.v4.n5.a1], [arXiv:hep-th/0103170].
- [DW90] R. Dijkgraaf and E. Witten, *Topological Gauge Theories and Group Cohomology*, Commun. Math. Phys. **129** (1990), 393-429, [euclid:cmp/1104180750].

- [DHVW85] L. Dixon, J. Harvey, C. Vafa, and E. Witten, *Strings on orbifolds*, Nucl. Phys. B **261** (1985), 678-686, [doi:10.1016/0550-3213(85)90593-0].
- [Do19] T. Dove, *Twisted Equivariant Tate K-Theory*, [arXiv:1912.02374].
- [DDK80] E. Dror, W. Dwyer, and D. Kan, *Equivariant maps which are self homotopy equivalences*, Proc. Amer. Math. Soc. **80** 4 (1980), 670-672, [jstor:2043448].
- [Du94] M. Duff, *Kaluza-Klein Theory in Perspective*, in *The Oskar Klein Centenary*, World Scientific (1994), [doi:10.1142/2816], [arXiv:hep-th/9410046].
- [DHIS87] M. Duff, M. Howe, T. Inami, and K. Stelle, *Superstrings in $D = 10$ from Supermembranes in $D = 11$* , Phys. Lett. **B191** (1987), 70-74, [doi:10.1016/0370-2693(87)91323-2].
- [Du01b] D. Dugger, *Universal homotopy theories*, Adv. Math. **164** (2001), 144-176, [arXiv:math/0007070], [doi:10.1006/aima.2001.2014].
- [EML53] S. Eilenberg and S. MacLane, *On the Groups $H(\Pi, n)$, I*, Ann. Math. **58** (1953), 55-106, [doi:10.2307/1969820].
- [EML54] S. Eilenberg and S. MacLane, *On the Groups $H(\Pi, n)$, II: Methods of Computation*, Ann. Math. **60** (1954), 49-139, [doi:10.2307/2372629].
- [EZ53] S. Eilenberg and J. Zilber, *On Products of Complexes*, Amer. J. Math. **75** (1953), 200-204, [jstor:2372629].
- [EG17] N. Epa and N. Ganter, *Platonic and alternating 2-groups*, High. Struc. **1** (2017), 122-146, [arXiv:1605.09192].
- [FHHP00] B. Feng, A. Hanany, Y.-H. He, and N. Prezas, *Discrete Torsion, Non-Abelian Orbifolds and the Schur Multiplier*, J. High Energy Phys. **0101** (2001), 033, [arXiv:hep-th/0010023].
- [FStS12] D. Fiorenza, J. Stasheff, and U. Schreiber, *Čech cocycles for differential characteristic classes*, Adv. Theor. Math. Phys. **16** (2012), 149-250, [arXiv:1011.4735].
- [FSS12CS] D. Fiorenza, H. Sati, and U. Schreiber, *Extended higher cup-product Chern-Simons theories*, J. Geom. Phys. **74** (2013), 130-163, [doi:10.1016/j.geomphys.2013.07.011], [arXiv:1207.5449].
- [FSS16WZW] D. Fiorenza, H. Sati and U. Schreiber, *Super Lie n -algebra extensions, higher WZW models and super p -branes with tensor multiplet fields*, Int. J. Geom. Meth. Mod. Phys. **12** (2015) 1550018 [doi:10.1142/S0219887815500188], [arXiv:1308.5264].
- [FSS17Sph] D. Fiorenza, H. Sati and U. Schreiber, *Rational sphere valued supercocycles in M -theory and type IIA string theory*, J. Geom. Phys., **114** (2017), 91-108, [doi:10.1016/j.geomphys.2016.11.024], [arXiv:1606.03206].
- [FSS16TDI] D. Fiorenza, H. Sati, and U. Schreiber, *T -Duality from super Lie n -algebra cocycles for super p -branes*, Adv. Theor. Math. Phys. **22** (2018), 1209-1270, [doi:10.4310/ATMP.2018.v22.n5.a3], [arXiv:1611.06536].
- [FSS19Rat] D. Fiorenza, H. Sati and U. Schreiber, *The rational higher structure of M -theory*, Proc. of Higher Structures in M -Theory 2018, Fortschr. Phys. **67** 8-9 (2019), [doi:10.1002/prop.201910017], [arXiv:1903.02834].
- [FSS20HTDI] D. Fiorenza, H. Sati, and U. Schreiber, *Higher T -duality of super M -branes*, Adv. in Math. and Theor. Phys. **24** (2020), 621-708, [doi:10.4310/ATMP.2020.v24.n3.a3], [arXiv:1803.05634].
- [FSS19HypH] D. Fiorenza, H. Sati, and U. Schreiber, *Twisted Cohomotopy implies M -theory anomaly cancellation on 8-manifolds*, Comm. Math. Phys. **377** (2020), 1961-2025, [doi:10.1007/s00220-020-03707-2], [arXiv:1904.10207].
- [FSS20Cha] D. Fiorenza, H. Sati, and U. Schreiber, *The character map in non-abelian cohomology*, World Scientific (2023) [doi:10.1142/13422] [arXiv:2009.11909].
- [FSS21Str] D. Fiorenza, H. Sati and U. Schreiber *Twisted cohomotopy implies twisted String structure on $M5$ -branes* J. Math. Phys. **62** 042301 (2021), [doi:10.1063/5.0037786], [arXiv:2002.11093].
- [FSS22GS] D. Fiorenza, H. Sati and U. Schreiber, *Twistorial Cohomotopy Implies Green-Schwarz anomaly cancellation*, Rev. Math. Phys. **34** 05 (2022) 2250013, [doi:10.1142/S0129055X22500131], [arXiv:2008.08544].

- [Fr12] G. Friedman, *An elementary illustrated introduction to simplicial sets*, Rocky Mount. J. Math. **42** 2 (2012) 353-423, [doi:10.1216/RMJ-2012-42-2-353], [arXiv:0809.4221].
- [Fr20] G. Friedman, *Singular Intersection Homology*, Cambridge Univ. Press (2020), [doi:10.1017/9781316584446].
- [GSY07] D. Gaiotto, A. Strominger, and X. Yin, *The M5-Brane Elliptic Genus: Modularity and BPS States*, J. High Energy Phys. **08** (2007) 070, [arXiv:hep-th/0607010].
- [GY07] D. Gaiotto and X. Yin, *Examples of M5-Brane Elliptic Genera*, J. High Energy Phys. **11** (2007) 004, [arXiv:hep-th/0702012].
- [Ga07] N. Ganter, *Stringy power operations in Tate K-theory*, [arXiv:math/0701565].
- [Ga13] N. Ganter, *Power operations in orbifold Tate K-theory*, Homology Homotopy Appl. **15** (2013), 313-342, [arXiv:1301.2754].
- [Gi23] G. Giotopoulos, H. Sati, and U. Schreiber, *Field Theory via Higher Geometry I: Smooth Sets of Fields* (2023, in preparation)
- [GJ99] P. Goerss and R. F. Jardine, *Simplicial homotopy theory*, Birkhäuser (1999, 2009), [doi:10.1007/978-3-0346-0189-4].
- [GDR99] R. Gonzalez-Diaz and P. Real, *A Combinatorial Method for Computing Steenrod Squares*, J. Pure Appl. Alg. **139** (1999) 89-108, [arXiv:math/0110308].
- [Gr07] K. Gruher, *String Topology of Classifying Spaces*, PhD thesis, Stanford 2007, [proquest:304826261], [ncatlab.org/nlab/files/Gruher_FreeLoopSpaceOfClassifyingSpace.pdf]
- [Gui06] B. Guillou, *A short note on models for equivariant homotopy theory*, 2006, [ncatlab.org/nlab/files/GuillouModelsForEquivariantHomotopyTheory.pdf]
- [GPPV21] S. Gukov, D. Pei, P. Putrov, C. Vafa, *4-manifolds and topological modular forms*, J. High Energ. Phys. **2021** 84 (2021) [arXiv:1811.07884][doi:10.1007/JHEP05(2021)084]
- [HatAT] A. Hatcher, *Algebraic topology* [https://pi.math.cornell.edu/~hatcher/AT/ATpage.html]
- [HatSS] A. Hatcher, *Spectral sequences in algebraic topology I: The Serre spectral sequence* [https://pi.math.cornell.edu/~hatcher/SSAT/SSch1.pdf].
- [Hir02] P. Hirschhorn, *Model Categories and Their Localizations*, Math. Surveys and Monographs bf 99 Amer. Math. Soc. (2002) [ISBN:978-0-8218-4917-0].
- [Ho08] S. Hollander, *A homotopy theory for stacks*, Israel J. Math. **163** (2008), 93-124, [arXiv:math/0110247], [doi:10.1007/s11856-008-0006-5].
- [Hua18a] Z. Huan, *Quasi-Elliptic Cohomology I*, Adv. Math. **337** (2018), 107-138, [arXiv:1805.06305], [doi:10.1016/j.aim.2018.08.007].
- [Hua18b] Z. Huan, *Quasi-theories*, [arXiv:1809.06651].
- [HS20] Z. Huan and M. Spong, *Twisted Quasi-elliptic cohomology and twisted equivariant elliptic cohomology*, [arXiv:2006.00554].
- [HY22] Z. Huan and M. B. Young, *Twisted Real quasi-elliptic cohomology*, [arXiv:2210.07511].
- [HSS18] J. Huerta, H. Sati, and U. Schreiber, *Real ADE-equivariant (co)homotopy of super M-branes*, Commun. Math. Phys. **371** (2019), 425-524, [doi:10.1007/s00220-019-03442-3], [arXiv:1805.05987].
- [Ja15] J. F. Jardine *Local homotopy theory* Monographs in Mathematics, Springer, New York (2015), [doi:10.1007/978-1-4939-2300-7].
- [Jo87] J. D. S. Jones, *Cyclic homology and equivariant homology*, Invent. Math. **87** (1987), 403-423, [doi:10.1007/BF01389424].
- [Kan58] D. Kan, *On homotopy theory and c.s.s. groups*, Ann. Math. **68** (1958), 38-53, [jstor:1970042].
- [Ke82] M. Kelly, *Basic concepts of enriched category theory*, Lon. Math. Soc. Lec. Note Series **64**, Cambridge Univ. Press, 1982, [ISBN:9780521287029]; reprinted in Th. App. of Cat. **10** (2005), 1-136, [tac:tr10].
- [KM04] N. Kitchloo and J. Morava, *Thom Spectra for Loopgroup representations*, [arXiv:math/0404541].
- [Klein1884] F. Klein, *Vorlesungen über das Ikosaeder und die Auflösung der Gleichungen vom fünften Grade*, 1884, translated as *Lectures on the Icosahedron and the Resolution of Equations of Degree Five* by George Morrice 1888, [archive.org/details/cu31924059413439].

- [KSS09] J. R. Klein, C. Schochet, and S. B. Smith, *Continuous trace C^* -algebras, gauge groups and rationalization*, *J. Top. Anal.* **01** (2009), 261-288, [doi:10.1142/S179352530900014X], [arXiv:0811.0771].
- [KS04] I. Kriz and H. Sati, *M Theory, Type IIA Superstrings, and Elliptic Cohomology*, *Adv. Theor. Math. Phys.* **8** (2004), 345-395, [doi:10.4310/ATMP.2004.v8.n2.a3], [arXiv:hep-th/0404013].
- [KS05] I. Kriz and H. Sati, *Type IIB String Theory, S-Duality, and Generalized Cohomology*, *Nucl.Phys. B* **715** (2005) 639-664, [doi:10.1016/j.nuclphysb.2005.02.016], [arXiv:hep-th/0410293].
- [Le10] E. Lerman, *Orbifolds as stacks?*, *Enseign. Math.* **56** 3-4 (2010), 315-363, [doi:10.4171/LEM/56-3-4], [arXiv:0806.4160].
- [Lo92] J.-L. Loday, *Cyclic Homology*, Grundlehren **301** Springer, Berlin, 1992, [doi:10.1007/978-3-662-21739-9].
- [Lo15] J.-L. Loday, *Free loop space and homology*, in: *Free Loop Spaces in Geometry and Topology*, IRMA Lectures in Mathematics and Theoretical Physics **24**, EMS 2015 [arXiv:1110.0405], [ISBN:978-3-03719-153-8].
- [LPSS95] H. Lu, C. N. Pope, E. Sezgin, and K. S. Stelle, *Stainless super p-branes*, *Nucl. Phys. B* **456** (1995), 669-698, [doi:10.1016/0550-3213(95)00524-4], [arXiv:hep-th/9508042].
- [LU02] E. Lupercio, and B. Uribe, *Loop groupoids, gerbes, and twisted sectors on orbifolds*, in: A. Adem, J. Morava and Y. Ruan (eds.), *Orbifolds in Mathematics and Physics*, Madison, WI, 2001, in: *Contemp. Math.* **310**, Amer. Math. Soc., Providence, RI (2002), 163-184, [ams:conm-310], [arXiv:math/0110207].
- [LU04] E. Lupercio and B. Uribe, *Inertia orbifolds, configuration spaces and the ghost loop space*, *Quart. J. Math.* **55** (2004), 185-201, [doi:10.1093/qmath/hag053], [arXiv:math/0210222].
- [LU06] E. Lupercio and B. Uribe, *Holonomy for Gerbes over Orbifolds*, *J. Geom.Phys.* **56** (2006), 1534-1560, [doi:10.1016/j.geomphys.2005.08.006], [arXiv:math/0307114].
- [Lu09] J. Lurie, *Higher Topos Theory*, *Annals of Mathematics Studies* 170, Princeton University Press, 2009, [pup:8957].
- [MP04] W. Marzantowicz and C. Prieto, *The unstable equivariant fixed point index and the equivariant degree*, *J. Lond. Math. Soc.* **69** (2004), 214-230, [doi:10.1112/S0024610703004721].
- [MaSa04] V. Mathai and H. Sati, *Some Relations between Twisted K-theory and E_8 Gauge Theory*, *J. High Energ. Phys.* **2004** 03 (2004) 016, [doi:10.1088/1126-6708/2004/03/016], [arXiv:hep-th/0312033].
- [May67] P. May, *Simplicial Objects in Algebraic Topology*, Univ. Chicago Press, 1967, [ISBN:9780226511818], [ncatlab.org/nlab/files/May_SimplicialObjectsInAlgebraicTopology.pdf]
- [May72] P. May, *The Geometry of Iterated Loop Spaces*, Springer, Berlin, 1972, [doi:10.1007/BFb0067491].
- [MC67] M. C. McCord, *Homotopy type comparison of a space with complexes associated with its open covers*, *Proc. Amer. Math. Soc.* **18** (1967), 705-708, [jstor:2035443].
- [Me15] L. Menichi, *Rational homotopy – Sullivan models*, in: *Free Loop Spaces in Geometry and Topology*, IRMA Lect. Math. Theor. Phys., Europ. Math. Soc., (2015), [doi:10.4171/153], [arXiv:1308.6685].
- [MiSt74] J. Milnor and J. D. Stasheff, *Characteristic Classes*, *Ann. Math. Stud.* **76**, Princeton University Press (1974), [ISBN:9780691081229].
- [Mo54] J. Moore, *Homotopie des complexes monoideaux, I*, *Sem. Henri Cartan*, 1954-55, [numdam:SHC_1954-1955__7_2_A8_0].
- [NS18] T. Nikolaus and P. Scholze, *On topological cyclic homology*, *Acta Math.* **221** (2018), 203-409, [doi:10.2140/gt.2015.19.3105], [arXiv:1707.01799].
- [NSS12a] T. Nikolaus, U. Schreiber, and D. Stevenson, *Principal ∞ -bundles – General theory*, *J. Homotopy Rel. Struc.* **10** 4 (2015), 749-801, [doi:10.1007/s40062-014-0083-6], [arXiv:1207.0248].
- [Pa22] D. Pavlov, *Numerable open covers and representability of topological stacks*, *Topology Appl.* **318** 108203 (2022) 1-28 [arXiv:2203.03120][doi:10.1016/j.topol.2022.108203]
- [Qu67] D. Quillen, *Homotopical Algebra*, *Lecture Notes in Mathematics* **43**, Springer, Berlin, 1967, [doi:10.1007/BFb0097438].
- [Re10] C. Rezk, *Toposes and homotopy toposes*, lecture notes (2010), [https://faculty.math.illinois.edu/~rezk/homotopy-topos-sketch.pdf]

- [Re15] C. Rezk, *Elliptic cohomology and elliptic curves*, Felix Klein Lectures, Bonn (2015), [www.hcm.uni-bonn.de/fkl-rezk]
- [Re22] C. Rezk, *Introduction to quasicategories*, lecture notes (2022), [ncatlab.org/nlab/files/Rezk-IntroToQuasicategories.pdf]
- [RV20] E. Riehl and D. Verity, *Infinity category theory from scratch*, Higher Structures **4** (2020) 1, [arXiv:1608.05314].
- [Ro09] J. Rosenberg, *Topology, C^* -algebras, and string duality*, Reg. Conf. Ser. Math. **111**, Amer. Math. Soc. (2009), [doi:10.1090/cbms/111].
- [SS20Tad] H. Sati and U. Schreiber, *Equivariant Cohomology implies orientifold tadpole cancellation*, J. Geom. Phys. **156** (2020) 103775, [doi:10.1016/j.geomphys.2020.103775], [arXiv:1909.12277].
- [SS20Orb] H. Sati and U. Schreiber, *Proper Orbifold Cohomology*, [arXiv:2008.01101].
- [SS21EPB] H. Sati and U. Schreiber, *Equivariant principal ∞ -bundles*, [arXiv:2112.13654].
- [SS21MF] H. Sati and U. Schreiber, *M/F-Theory as Mf -theory*, Rev. Math. Phys. (2023, in print) [arXiv:2103.01877].
- [SS22Conf] H. Sati and U. Schreiber, *Differential Cohomology implies intersecting brane observables*, Adv. Theor. Math. Phys. **26** 4 (2022) [doi:10.4310/ATMP.2022.v26.n4.a4] [arXiv:1912.10425]
- [SSS12] H. Sati, U. Schreiber, and J. Stasheff, *Twisted differential string and fivebrane structures*, Commun. Math. Phys. **315** (2012), 169-213, [doi:10.1007/s00220-012-1510-3], [arXiv:0910.4001].
- [SV21] H. Sati and A. A. Voronov, *Mysterious triality and rational homotopy theory*, Comm. Math. Phys. (2023) [doi:10.1007/s00220-023-04643-7] [arXiv:2111.14810].
- [Sc13] U. Schreiber, *Differential cohomology in a cohesive infinity-topos*, [arXiv:1310.7930].
- [Sc16] U. Schreiber, *Super Lie n -algebra of Super p -branes*, lecture notes (2016-2017), [ncatlab.org/schreiber/show/Super+Lie+n-algebra+of+Super+p-branes].
- [Sc18] U. Schreiber, *Introduction to Higher Supergeometry*, lecture at *Higher Structures in M-Theory* (2018), [ncatlab.org/schreiber/show/Introduction+to+Higher+Supergeometry].
- [SS03] S. Schwede and B. Shipley, *Equivalences of monoidal model categories*, Algebr. Geom. Topol. **3** (2003), 287-334, [euclid.agt/1513882376], [arXiv:math/0209342].
- [Se01] S. Seki, *Discrete Torsion and Branes in M-theory from Mathematical Viewpoint*, Nucl. Phys. **B606** (2001) 689-698, [doi:10.1016/S0550-3213(01)00245-0], [arXiv:hep-th/0103117].
- [Sh03] E. Sharpe, *Analogues of Discrete Torsion for the M-Theory Three-Form*, Phys. Rev. **D68** (2003) 126004, [doi:10.1103/PhysRevD.68.126004], [arXiv:hep-th/0008170].
- [St15] N. Stapleton, *Transchromatic twisted character maps*, J. Hom. Relat. Struct. **10** (2015), 29–61, [doi:10.1007/s40062-013-0040-9], [arXiv:1304.5194].
- [St67] N. Steenrod, *A convenient category of topological spaces*, Michigan Math. J. **14** (1967), 133–152, [euclid.mmj/1028999711].
- [Sto19] A. Stoffel, *Dimensional reduction and the equivariant Chern character*, Algebr. Geom. Topol. **19** (2019), 109-150, [doi:10.2140/agt.2019.19.109], [arXiv:1703.00314].
- [TV05] B. Toën and G. Vezzosi, *Homotopical Algebraic Geometry I: Topos theory*, Adv. Math. **193** (2005), 257-372, [doi:10.1016/j.aim.2004.05.004], [arXiv:math/0207028].
- [tD79] T. tom Dieck, *Transformation Groups and Representation Theory*, Lecture Notes in Mathematics **766**, Springer, 1979, [doi:10.1007/BFb0085965].
- [tD87] T. tom Dieck, *Transformation Groups*, de Gruyter, Berlin, 1987, [doi:10.1515/9783110858372].
- [TZ08] S. Tomoda and P. Zvengrowski, *Remarks on the cohomology of finite fundamental groups of 3-manifolds*, Geom. Topol. Monogr. **14** (2008), 519-556, [doi:10.2140/gtm.2008.14.519], [arXiv:0904.1876].
- [VPB85] M. Vigué-Poirrier, D. Burghelea, *A model for cyclic homology and algebraic K-theory of 1-connected topological spaces*, J. Diff. Geom. **22** (1985), 243-253, [euclid:1214439821].
- [Wa22] K. Waldorf, *Geometric T-duality: Buscher rules in general topology* [arXiv:2207.11799]
- [Wi08] S. Willerton, *The twisted Drinfeld double of a finite group via gerbes and finite groupoids*, Algebr. Geom. Topol. **8** (2008), 1419-1457, [doi:10.2140/agt.2008.8.1419], [arXiv:math/0503266].

- [Wit88] E. Witten, *The index of the Dirac operator in loop space*, in *Elliptic curves and modular forms in algebraic topology*, Lecture Notes in Math. **1326** (1988), 161–181, Springer, Berlin, [doi:10.1007/BFb0078045].
- [Wi95] E. Witten, *String Theory Dynamics In Various Dimensions*, Nucl. Phys. **B443** (1995), 85-126, [doi:10.1016/0550-3213(95)00158-0], [arXiv:hep-th/9503124].